

2-2012

Randomized controlled trial of community-based dancing to modify disease progression in Parkinson disease

Ryan P. Duncan

Washington University School of Medicine in St. Louis

Gammon M. Earhart

Washington University School of Medicine in St. Louis

Follow this and additional works at: http://digitalcommons.wustl.edu/pt_facpubs

Recommended Citation

Duncan, Ryan P. and Earhart, Gammon M., "Randomized controlled trial of community-based dancing to modify disease progression in Parkinson disease" (2012). *Physical Therapy Faculty Publications*. Paper 29.
http://digitalcommons.wustl.edu/pt_facpubs/29

This Article is brought to you for free and open access by the Program in Physical Therapy at Digital Commons@Becker. It has been accepted for inclusion in Physical Therapy Faculty Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Randomized Controlled Trial of Community-Based Dancing to Modify Disease Progression in Parkinson Disease

Ryan P. Duncan, PT, MPT¹ and Gammon M. Earhart, PhD, PT¹⁻³

Running Head: Community-Based Dance in PD

¹ Washington University in St. Louis School of Medicine, Program in Physical Therapy

² Washington University in St. Louis School of Medicine, Department of Anatomy & Neurobiology

³ Washington University in St. Louis School of Medicine, Department of Neurology

Corresponding Author:

Gammon M. Earhart, PhD, PT
Washington University School of Medicine
Program in Physical Therapy
Campus Box 8502
4444 Forest Park Blvd.
St. Louis, MO 63108
314-286-1425
earhartg@wusm.wustl.edu

Text word count: 4671

Number of tables: 2

Number of figures: 5

This manuscript has been accepted for publication in *Neurorehabilitation and Neural Repair*. Documentation confirming acceptance is included as the last page of this document.

Abstract

Background: Tango dancing has been effective in improving measures of physical function in people with Parkinson disease (PD). However, all previous studies were institution-based, tested participants on medication, and employed short-term interventions.

Objective: To determine the effects of a 12-month community-based tango program for individuals with PD on disease severity and physical function.

Methods: Sixty-two participants were randomly assigned to a twice weekly, community-based Argentine Tango program or a Control group (no intervention). Participants were assessed off anti-Parkinson medication at baseline, 3, 6, and 12 months. Our primary outcome measure was the Movement Disorders Society-Unified Parkinson Disease Rating Scale 3 (MDS-UPDRS-3). Secondary outcome measures were the MDS-UPDRS-1, MDS-UPDRS-2, MiniBESTest balance test; Freezing of Gait Questionnaire (FOG_Q); six minute walk (6MWT); gait velocity for comfortable forward, fast as possible forward, dual task, and backward walking; and Nine Hole Peg Test (9HPT).

Results: Groups were not different at baseline. Overall, the Tango group improved while the Control group showed little change on most measures. For the MDS-UPDRS-3 there was no significant change in the Control group from baseline to 12 months, whereas the Tango group had a reduction of 28.7% (12.8 points). There were significant group by time interactions for MDS-UPDRS-3, MiniBESTest, FOG_Q, 6MWT, forward and dual task walking velocities, and 9HPT in favor of the dance group.

Conclusions: Improvements in the Tango group were apparent off medication, suggesting that long-term participation in tango may modify progression of disability in PD.

Keywords: Parkinson disease, exercise, gait, balance, rehabilitation

Introduction

To combat mobility-related problems in PD, non-pharmacologic tactics such as exercise are needed. Traditional exercise is effective in improving balance, lower extremity strength, and gait speed.¹ Likewise, tango dancing, an alternative form of exercise, improves balance, gait, and quality of life.^{2,3} Previous studies examining the effects of dance on those with PD presented some limitations, however. First, participants were evaluated on anti-PD medications. While testing participants on medication may provide insight into how they perform daily activities, some of the deficits caused by PD may go unnoticed.⁴ For a more accurate picture of the underlying disease process, and to ultimately determine whether exercise may be disease-modifying, testing off medication is warranted. Second, although institution-based dance programs proved effective in prior studies, the Surgeon General's Call to Action encourages researchers and those with disabilities "to jointly develop community-based healthcare and wellness programs for people with disabilities and research their efficacy."⁵ Third, prior studies of dance utilized interventions of 3 months or less. There is a clear need for studies that determine the effects of long-term exercise in PD.⁶⁻⁸

The purpose of this study was to determine the effects of a long-term community-based dance program in people with PD, with all evaluations conducted off medication. As evidence emerges suggesting that exercise may positively impact multiple dimensions,⁹ and with recent animal studies suggesting exercise may have neuroprotective effects in PD,¹⁰ we were interested in the effects of dance on disease progression as assessed by our primary variable, the Movement Disorder Society-Unified Parkinson Disease Rating Scale 3 (MDS-UPDRS-3). We hypothesized that: 1) the Tango group would demonstrate improvements in disease severity and physical function not noted in the Control group and 2) Tango participants would show larger improvements in function at 6 and 12 months compared to 3 months.

Methods

Participants

The principal investigator (GME) recruited individuals with clinically defined “definite PD”¹¹⁻¹³ (Hoehn & Yahr Stages I-IV) from the Washington University Movement Disorders Center and through advertisements in a local PD newsletter. Phone interviews were conducted with potential participants to determine if they should be excluded based upon the following: 1) serious medical condition, 2) evidence of abnormality other than PD-related changes on brain imaging (previously done for clinical evaluations), 3) history or evidence of neurological deficit other than PD, or 4) history or evidence of musculoskeletal problem. This work was approved by the Human Research Protection Office (HRPO) at Washington University and each participant provided written informed consent. The trial was registered on ClinicalTrials.gov as “PD4PD: Partnered Dance for Parkinson Disease”, NCT01388556.

Study Design

This was a randomized controlled trial where participants were assigned to the Tango or Control group by the principal investigator using an online random number generator. Tango participants attended twice weekly, one hour community-based Argentine Tango classes for 12 months. Participants danced both leader and follower roles, changed partners frequently, and learned new steps and/or integrated previously learned steps in new ways at each class throughout the 12 months. The tango paradigm upon which classes were modeled has been described in detail¹⁴. Participants were encouraged to learn and perform the dance to the best of their abilities but dance performance was not evaluated. Control participants had no prescribed exercise and were instructed to go about their lives as usual.

Baseline evaluations were completed from October-December 2009 and included assessment of physical activity levels (Physical Activity Scale for the Elderly, PASE).¹⁵ Three-month evaluations were completed from January-March, 6-month evaluations from April-June, and 12-month evaluations from October-December 2010.

Participants were off all anti-PD medications for at least 12 hours prior to evaluation; time of day for all evaluations was kept the same for each participant. At each visit, participants were assessed by the same rater, a physical therapist (RPD), who was blinded to group.

Outcome Measures

Disease Severity

MDS-UPDRS sections 1-3 were used to measure disease severity.¹⁶ Section 1 examines non-motor experiences, section 2 covers ADLs, and section 3 assesses motor symptoms including tremor, rigidity, bradykinesia, gait, and postural instability. Higher scores indicate greater disease severity. The MDS-UPDRS-3 was our primary variable of interest. We conducted additional analyses on the specific motor components assessed within the MDS-UPDRS-3: tremor (sum of MDS-UPDRS-3 items 3.15-3.18), rigidity (sum of items 3.3a-3.3e), bradykinesia (sum of items 3.4-3.8 and 3.14), and postural instability/gait disorder (PIGD, sum of items 3.9-3.13).

Balance

Balance was assessed using the MiniBESTest, a 14-item tool measuring performance of dynamic balance tasks.¹⁷ This test has high inter-rater and test-retest reliability ($ICC \geq 0.92$ and $ICC \geq 0.88$, respectively) in PD.¹⁸ Lower scores indicate greater deficits in balance.

Gait

Freezing of gait was quantified using the Freezing of Gait Questionnaire (FOGQ).^{19,20} Higher scores indicate greater difficulty with walking and freezing. The Six Minute Walk Test (6MWT) was utilized to measure walking endurance.²¹ Individuals were instructed to cover as much ground as possible while walking at a safe, comfortable pace in a 30.48m hallway.

A 4.87m GAITRite (CIR Systems, Inc, Havertown, PA) was used to measure walking velocity during comfortable forward, fast as possible forward, dual task, and backward walking. Mean velocity for 3 trials of each condition was determined. For comfortable pace forward and backward walking, participants were instructed to walk at a self-determined “normal” speed. For fast as possible forward walking, participants were instructed to walk as quickly as possible without running. For dual task walking, participants were instructed to walk forward at a self-determined “normal” speed while naming as many words as possible that began with a certain letter. The same three letters were used for all participants at all time points. Subjects were instructed to begin walking as soon as the letter was provided. Number of correct and incorrect answers on each trial was recorded and the three trials averaged to provide a measure of overall task performance, which was not different between groups and did not change over the course of the study.

Upper Extremity Function

The Nine Hole Peg Test (9HPT) was utilized to evaluate upper extremity function^{22,23} to determine whether participation in the intervention might affect variables not directly related to the intervention. Participants performed 2 trials of the 9HPT with each hand. The average of all 4 trials yielded a composite mean value for 9HPT performance.

Procedures

The evening prior to testing, participants received a phone call and were reminded to not take any anti-PD medication for at least 12 hours prior to arrival. Upon arrival, the order of testing was: 1) MDS-UPDRS (1-3), 2) MiniBESTest, 3) gait, and 4) 9HPT. Several measures were included in the test battery to comprehensively evaluate the effects of the exercise intervention on disease progression, balance, gait, and upper extremity function. We deemed all of these to be important areas to include as few of these have been assessed in long-term exercise intervention studies.

Statistical Analyses

Power analyses based upon data from previous published tango studies^{2,3,24-26} and the minimal clinically important difference in the UPDRS-3²⁷ indicated a need for approximately 30 subjects per group to have 80% power at $p=0.05$. T-tests compared baseline demographic and disease severity characteristics between groups ($p = 0.05$). All other data were analyzed using two-way repeated measures ANOVAs with group (Tango or Control) and time (baseline, 3, 6, or 12 months) as factors ($p = 0.05$). When appropriate, Tukey-Kramer multiple comparison posthoc tests were used to determine specific differences between groups within a given time point and within a group across time points. An intent-to-treat analysis with the last observation carried forward was employed with any participants who completed the baseline visit and at least one other evaluation. Data analysis was completed in NCSS.²⁸

Results

123 participants were screened, with 62 randomized to the Tango or Control group. Reasons for exclusion included unwillingness to skip medications for the evaluations, failure to meet inclusion/exclusion criteria, and practical issues such as transportation difficulty (see Figure 1). Participants were lost at various time points (Figure 1). The final analysis included 52 individuals. Participants who dropped out at any point after 3 months were still included using the last observation carried forward. There were no significant differences at baseline between the Tango and Control groups for demographic measures (Table 1) and no differences at baseline in physical activity levels as assessed by the PASE (Tango = 124.2 ± 16.3 , Control = 115.4 ± 13.9).

Disease Severity

Motor symptoms (MDS-UPDRS-3) improved in the Tango group and did not change in the Control group, resulting in a significant group by time interaction ($F=9.82$, $p<0.001$, Figure 2A). There were also significant main effects of time ($F=9.40$, $p=0.004$) and group ($F=23.1$, $p<0.001$) for motor symptoms. Tango motor symptom scores at 3, 6, and 12 months were significantly better (i.e. lower) than Control scores at 3, 6, and

12 months, respectively. Within the Tango group, motor symptom scores were significantly better at 3, 6, and 12 months compared to baseline, and significantly better at 6 and 12 months compared to 3 months. Table 2 presents the average MDS-UPDRS-3 values and 95% confidence intervals for each group at each time point. There were no differences in ADLs (MDS-UPDRS-2) or non-motor symptoms (MDS-UPDRS-1) between groups, and no significant changes in ADLs or non-motor symptoms during the study (Figure 2B,C).

Tremor scores decreased very slightly over time in both groups (Figure 3A) resulting in a significant main effect of time for tremor ($F=6.70$, $p<0.001$). There were no differences between groups and no significant interaction for tremor. Rigidity did not change in the Tango group and increased in the Control group over the course of the study, resulting in a significant group by time interaction ($F=5.31$, $p=0.002$, Figure 3B). Rigidity was significantly lower in the Tango group compared to the Control group at 6 and at 12 months. There was also a significant main effect of time for rigidity ($F=11.72$, $p<0.001$). Bradykinesia decreased substantially in the Tango group and changed little in the Control group, resulting in a significant group by time interaction ($F=8.35$, $p<0.001$, Figure 3C). Bradykinesia was significantly lower in the Tango group compared to the Control group at 6 and at 12 months. Within the Tango group, bradykinesia was significantly lower at 3, 6, and 12 months compared to baseline and at 6 and 12 months compared to 3 months. There were also significant main effects of group ($F=5.65$, $p=0.02$) and time ($F=42.14$, $p<0.001$) for bradykinesia. PIGD scores decreased in the Tango group, resulting in a significant group by time interaction ($F=4.21$, $p=0.007$, Figure 3D). PIGD scores in the Tango group were significantly better than the Control group at 6 and at 12 months. Within the tango group, PIGD scores were significantly better at 12 months compared to 3 months.

Balance

Balance improved in the Tango group and worsened slightly in the Control group over the course of the study, resulting in a significant group by time interaction ($F=11.73$, $p<0.001$, Figure 4A). Balance scores in the Tango group at 3, 6, and 12 months were significantly better than Control scores at 3, 6, and 12 months,

respectively. Within the Tango group, balance scores at 3, 6 and 12 months were significantly better compared to baseline.

Gait

There were no significant differences between groups at any time point for freezing of gait as assessed by the FOG_Q. There was, however, a significant group by time interaction ($F=4.2$, $p=.006$) for freezing of gait (Figure 4B). Within the Control group, there was significantly more freezing reported at 12 months compared to baseline.

Six-minute walk distance held steady in the Tango group and decreased in the Control group over the course of the study, resulting in a significant group by time interaction ($F=3.33$, $p=0.02$, Figure 4C). Distance walked by the Tango group at 12 months was significantly longer than distance walked by the Control group at 12 months.

In general, walking velocity in all conditions increased over the course of the study for the Tango group and did not change in the Control group (Figure 5A-5D). For forward walking (Figure 5A) there was a significant group by time interaction ($F=2.74$, $p=0.04$). At 6 and 12 months, the Tango group had significantly higher preferred forward walking velocity than the Control group at 6 and 12 months, respectively. For fastest possible walking (Figure 5B), there were no significant differences between groups and no significant changes in velocity. For dual task walking (Figure 5C), there was an interaction between group and time ($F=3.57$, $p=0.02$) as well as a significant main effect of time ($F=3.31$, $p=0.02$). At 6 and 12 months, the Tango group had significantly faster dual task walking velocity than the Control group at 6 and 12 months, respectively. Within the Tango group, dual task walking velocity was significantly faster at 6 and 12 months compared to baseline. Finally, for backward walking velocity (Figure 5D), there was a main effect of time only ($F=3.04$, $p=0.03$).

Upper Extremity Function

Performance on the 9HPT improved in the Tango group and worsened slightly in the Control group over the course of the study, resulting in a significant group by time interaction ($F=3.83$, $p=0.01$, Figure 4D). Tango scores at 6 and 12 months were significantly better than Control scores at 6 and 12 months, respectively. Within the Tango group, scores were significantly better at 12 months compared to baseline.

Participant Adherence

Over the 12 month study, there was a 37% attrition rate in the Control group and a 50% attrition rate within the Tango group. Throughout the 12 months, the 16 Tango participants that attended consistently from start to end came to an average of $78.5 \pm 3\%$ of all classes. Following 12 months of participation, 11 of the 16 individuals in the Tango group chose to continue attending classes even though they had formally completed all study requirements and were no longer expected to attend.

Discussion

This study is the first to examine the effects of a community-based exercise program on individuals with PD tested off medication over a 12-month period. Following participation in Argentine Tango dance classes, participants demonstrated a significant reduction in disease severity, as well as significant improvements in gait, balance, and upper extremity function when compared to controls.

Disease Severity

This is, to our knowledge, the first study to report changes in motor symptom severity during a long-term exercise intervention with participants assessed off medication. Given this off medication testing, it is unlikely that improvements noted are due to improved response to pharmacologic interventions. Improvements in motor symptoms over the 12-month study, during which MDS-UPDRS-3 scores decreased by 12.8 points, were larger

than previously reported changes of 4.6 and 8 points in the UPDRS-3 in short-term studies examining the effects of tango in participants on medication.^{2,24}

Other exercise interventions have yielded improvements in PD motor symptom severity. Improvements of 25% (6-7 points) have been reported following treadmill training²⁹ and sensory attention focused exercise.³⁰ However, these studies examined subjects on medication. In the present study we noted MDS-UPDRS-3 score improvements of 10.3, 23.1, and 28.7 % at 3, 6, and 12 months, respectively. Given that these improvements in MDS-UPDRS-3 scores off medication in the Tango group, this suggests that participation in the Tango program may have a disease modifying effect. Examination of specific components of the MDS-UPDRS-3 indicates that tango may have a positive influence not only on balance and gait, as might be expected, but also bradykinesia and rigidity. This supports the idea that tango may have a broad impact on motor symptom progression rather than just targeting the gait and balance aspects specifically practiced in the context of dancing.

Although not significant, the Tango group demonstrated reduced scores on the MDS-UPDRS-1 and 2, indicating some improvement in non-motor symptoms and activities of daily living, respectively. It is disappointing that larger effects were not seen on these scales, but this was likely due in part to the much larger variability between subjects. This variability may reflect the fact that the MDS-UPDRS-1 and 2 are questionnaires answered by each participant, whereas the MDS-UPDRS-3 was rated by a single trained individual. Additional work is warranted, perhaps using measures in addition to the MDS-UPDRS-1 and 2, to explore the effects of long-term exercise on non-motor symptoms and ADLs.

Physical Function

Participants in Tango demonstrated significant improvements in balance at 3, 6, and 12 months. At 12 months, the Tango group had an average MiniBESTest score of 21.3 +/- 1.0 compared to the Control group which had an average score of 17.2 +/- 1.1 at the same time point. This difference of 4 points is substantial considering that the full MiniBESTest scale is only 32 points. Because the MiniBESTest is relatively new and MCID values

are not available, it is not clear if this 4 point difference is clinically meaningful. Previous literature shows that different forms of exercise can have a positive impact on balance for individuals with PD in the short-term,^{2,31-35} while reports on medium-term (6 months) interventions show mixed results. Direct comparison of the present study with previous work is difficult because of the different balance measures used and the fact that we tested participants off medication. However, our results are promising and suggest that balance can improve in the short-term (i.e. at 3 months), continue to improve out to 6 months, and can be maintained at 12 months with continued exercise participation. Additional work is warranted to determine if long-term participation in exercise modifies fall risk or fall rates.

With respect to gait, those in the Tango group demonstrated improved self-selected walking speed compared to Controls following 6 and 12 months of dance exercise. These findings are in concert with previous work demonstrating improved self-selected walking speed following a shorter dance intervention.²⁵ Treadmill training studies have reported larger increases in gait speed than those noted in the present study, with gains of approximately 0.15m/s^{36,29} Tango dancing may not be as intense as treadmill training, possibly accounting for the smaller improvements in gait speed in the present study compared to treadmill training. Nonetheless, the change of 0.09 m/s in gait speed from baseline to 12 months in the Tango group is close to the 0.1 m/s change generally accepted as a functionally meaningful difference.³⁷

We also noted a 0.12 m/s increase in dual task walking speed after 12 months of Tango. Improved dual task walking may relate to the task-specific practice of multitasking during dancing. As participants dance, they must execute one movement while planning the next, all while attending to other couples on the dance floor and to the music. This practice of multi-tasking, coupled with the presence of the music to serve as an auditory cue, may facilitate dual task walking. Improvements in dual task walking have also been demonstrated with cueing,³⁸ and through specific practice of multiple-task walking.³⁹ With respect to FOG, within the Control group there was a significant difference from baseline to 12 months. This is not surprising as it is well known that FOG occurs more frequently as PD progresses.^{40,41} It is interesting, though, that the Tango group presented no differences in

FOG from baseline to 12 months, suggesting that exercise may delay the progression of FOG. This seems feasible, as tango dancing requires many starts, stops, turns, and movement in confined spaces, thereby incorporating direct practice of walking in the situations that commonly provoke freezing.⁴²

The Tango group covered significantly greater distance in the 6MWT compared to Controls at the end of the study. Improvements in 6MWT with shorter tango interventions have been reported previously.^{26,43} Improvements in 6MWT have also been demonstrated with other exercise interventions including boxing⁴⁴ and treadmill training.⁴⁵ As with gait velocity, changes in 6MWT in the present study were much smaller than those reported following treadmill training,^{46,47} but it is possible that the low intensity aerobic exercise provided through dance may have helped Tango participants to maintain 6MWT performance while Control participants declined. Alternatively, or in addition, maintenance of 6MWT distance in the Tango group may be related to improvements in balance, as balance has been suggested to play a significant role in 6MWT performance.⁴⁸

Somewhat surprisingly, we noted significant improvements in 9HPT performance in the Tango group compared to Controls. While a previous exercise study also demonstrated improvements in 9HPT performance, that program included movement exercises for the upper extremities and hands.⁴⁹ As there are no specific hand movements practiced with tango dancing, we propose that improved 9HPT performance may be reflective of a global impact of exercise on bradykinesia. This is supported by the improvements in MDS-UPDRS-3 bradykinesia scores and is in keeping with the suggestion that participation in exercise may have disease modifying effects. This further suggests that the effects of tango dancing may extend beyond the specific types of tasks practiced in the context of the dance.

Participant Adherence

The attrition rate within the Tango group was relatively high; however, it is important to note that this study was much longer than previous tango studies.^{2,26} The 16 participants completing the full 12 months of the study attended 78% of dance classes. Studies of long-term exercise in older adults without PD have reported

attendance rates of 65% over a 6 month period⁵⁰ and 58% over a 12 month period.⁵¹ While attendance rates should be cautiously interpreted when comparing this study to the aforementioned studies due to differences in sample sizes, we think the higher attendance rate among those who completed the full 12 months of tango may speak to the high level of satisfaction with the dance classes. However, there was an overall 50% attrition rate in the tango group over 12 months. More than half of the individuals who stopped participating did so due to unrelated medical conditions that developed over the course of the 12 months. Our attrition in the Tango group at 3 months was 18%, which is in line with reported attrition rates of 14% and 23% for traditional 12-week exercise programs³⁰. At 6 months, attrition in the Tango group was 37%, which is high compared to a 6-month study of traditional exercise that reported 13% attrition.³⁵ One possible explanation for our higher attrition rate is the fact that the 6-month study with low attrition required only monthly visits to an exercise class with the other exercise sessions done at home. In addition, our study required participants to be off medication for all evaluations. This may have deterred some people from continuing if the experience of withdrawing from medications was unpleasant or stressful. Finally, the present study was longer than most prior work and as such one would expect higher attrition.

Why Dance May Be Effective

We suggest that tango may be especially helpful compared to other dances because of the specific movements it incorporates, such as backward walking.²⁶ Tango offers both physical and cognitive challenges, as it incorporates low level aerobic activity and movements that challenge gait and balance while also requiring high level multi-tasking and progressive motor skill learning in the presence of external cues provided by the music and the partner.

Study Limitations

This study has several limitations. We only evaluated subjects off medication in order to eliminate the potentially confounding effects of medication. However, testing both off and on medication is recommended for

future studies to increase relevance to everyday functioning. Another limitation is that the Control group received no intervention and no control for attention or socialization. As such we cannot say whether and how the social and attentional aspects of participation contributed to the outcomes. The Control group did enable us to examine the natural history in the absence of a specific intervention. Future work should control for attention and socialization, and compare other exercise approaches and intensities.

Conclusion

Long-term participation in community-based dance exercise benefits people with PD. Socially engaging and enjoyable skill-based exercise may help to promote long-term participation.

Acknowledgements

This study was funded by the Parkinson's Disease Foundation. General support came from the Greater St. Louis Chapter of the American Parkinson Disease Association (APDA) and the APDA Center for Advanced PD Research at Washington University. Thanks to Ruth Porter, DPT and John Michael Rotello for Tango class instruction and to the student volunteers who assisted.

References

1. Goodwin VA, Richards SH, Taylor RS, Taylor AH, Campbell JL. The effectiveness of exercise interventions for people with Parkinson's disease: a systematic review and meta-analysis. *Mov Disord* 2008; 23: 631-40.
2. Hackney ME, Earhart GM. Short duration, intensive tango dancing for people with Parkinson disease: an uncontrolled pilot study. *Complement Ther Med* 2009; 17: 203-7.
3. Hackney ME, Earhart GM. Health-related quality of life and alternative forms of exercise in Parkinson disease. *Parkinsonism Relat Disord* 2009; 15: 644-8.
4. Gordon AM, Reilmann R. Getting a grasp on research: does treatment taint testing of parkinsonian patients? *Brain* 1999; 122: 1597-8.
5. U.S. Department of Health and Human Services. The Surgeon General's Call to Action to Improve the Health and Wellness of Persons with Disabilities. U.S. Department of Health and Human Services, Office of the Surgeon General, 2005.
6. Earhart GM. Dance as therapy for individuals with Parkinson disease. *Eur J Phys Rehabil Med* 2009; 45: 231-8.
7. Crizzle AM, Newhouse IJ. Is physical exercise beneficial for persons with Parkinson's disease? *Clin J Sport Med* 2006; 16: 422-5.
8. Morris ME, Iansek R, Kirkwood B. A randomized controlled trial of movement strategies compared with exercise for people with Parkinson's disease. *Mov Disord* 2009; 24: 64-71.
9. Zesiewicz TA, Evatt ML. Potential influences of complementary therapy on motor and non-motor complications in Parkinson's disease. *CNS Drugs* 2009; 23: 817-35.

10. Hirsch MA, Farley BG. Exercise and neuroplasticity in persons living with Parkinson's disease. *Eur J Phys Rehabil Med* 2009; 45: 215-29.
11. Racette BA, Rundle M, Parsian A, Perlmutter JS. Evaluation of a screening questionnaire for genetic studies of Parkinson's disease. *Am J Med Genet* 1999; 88: 539-43.
12. Calne DB, Snow BJ, Lee C. Criteria for diagnosing Parkinson's disease. *Ann Neurol* 1992; 32: S125-7.
13. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992; 55: 181-4.
14. Hackney ME, Earhart GM. Recommendations for implementing partnered tango classes for persons with Parkinson disease. *Amer J Dance Ther* 2010; 32: 41.
15. Washburn R, Smith K, Jette A, Janney C. The Physical Activity Scale for the Elderly (PASE): development and evaluation. *J Clin Epidemiol.* 1993 Feb;46(2):153-62.
16. Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, Martinez-Martin P, et al. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPRDS): scale presentation and clinimetric testing results. *Mov Disord* 2008; 23: 2129-70.
17. Franchignoni F, Horak F, Godi M, Nardone A, Giordano A. Using psychometric techniques to improve the Balance Evaluation Systems Test: the mini-BESTest. *J Rehabil Med* 2010; 42: 323-31.
18. Leddy AL, Crowner B, Earhart GM. Utility of the Mini-BESTest, BESTest, and BESTest sections for balance assessments in Parkinson disease. *J Neurol Phys Ther.* In press.

19. Giladi N, Shabtai H, Simon ES, Biran S, Tal J, Korczyn AD. Construction of freezing of gait questionnaire for patients with Parkinsonism. *Parkinsonism Relat Disord* 2000; 6: 165-170.
20. Giladi N, Tal J, Azulay T, Rascol O, Brooks DJ, Melamed E, et al. Validation of the freezing of gait questionnaire in patients with Parkinson's disease. *Mov Disord* 2009; 24: 655-61.
21. Steffen T, Seney M. Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-item short-form health survey, and the unified Parkinson disease rating scale in people with parkinsonism. *Phys Ther* 2008; 88: 733-46.
22. Oxford Grice K, Vogel KA, Le V, Mitchell A, Muniz S, Vollmer MA. Adult norms for a commercially available Nine Hole Peg Test for finger dexterity. *Am J Occup Ther* 2003; 57: 570-3.
23. Earhart, G.M. Cavanaugh, J.T., Ellis, T., Ford, M.P., Foreman, K.B., Dibble, L.E. The 9 Hole Peg Test of Upper Extremity Function: Normative Values, Test-retest Reliability, and Factors Contributing to Performance in People with Parkinson Disease. *J Neurol Phys Ther*, in press.
24. Hackney ME, Kantorovich S, Levin R, Earhart GM. Effects of tango on functional mobility in Parkinson's disease: a preliminary study. *J Neurol Phys Ther* 2007; 31: 173-9.
25. Hackney ME, Earhart GM. Effects of dance on gait and balance in Parkinson's disease: a comparison of partnered and nonpartnered dance movement. *Neurorehabil Neural Repair* 2010; 24: 384-92.
26. Hackney ME, Earhart GM. Effects of dance on movement control in Parkinson's disease: a comparison of Argentine tango and American ballroom. *J Rehabil Med* 2009; 41: 475-81.
27. Schrag A, Sampaio C, Counsell N, Poewe W. Minimal clinically important change on the unified Parkinson's disease rating scale. *Mov Disord* 2006; 21: 1200-7.

28. Hintze J. 2009 NCSS. NCSS, LLC. Kaysville, Utah. www.ncss.com
29. Herman T, Giladi N, Gruendlinger L, Hausdorff JM. Six weeks of intensive treadmill training improves gait and quality of life in patients with Parkinson's disease: a pilot study. *Arch Phys Med Rehabil* 2007; 88: 1154-8.
30. Sage MD, Almeida QJ. Symptom and gait changes after sensory attention focused exercise vs aerobic training in Parkinson's disease. *Mov Disord* 2009; 24: 1132-8.
31. Nocera J, Horvat M, Ray CT. Effects of home-based exercise on postural control and sensory organization in individuals with Parkinson disease. *Parkinsonism Relat Disord* 2009; 15: 742-5.
32. Hackney ME, Earhart GM. Tai Chi improves balance and mobility in people with Parkinson disease. *Gait Posture* 2008; 28: 456-60.
33. Hirsch MA, Toole T, Maitland CG, Rider RA. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Arch Phys Med Rehabil* 2003; 84: 1109-17.
34. Gobbi LT, Oliveira-Ferreira MD, Caetano MJ, Lirani-Silva E, Barbieri FA, Stella F, et al. Exercise programs to improve mobility and balance in people with Parkinson's disease. *Parkinsonism Relat Disord* 2010; 15 Suppl 3: S49-52.
35. Allen NE, Canning CG, Sherrington C, Lord SR, Latt MD, Close JC, et al. The effects of an exercise program on fall risk factors in people with Parkinson's disease: a randomized controlled trial. *Mov Disord* 2010; 25: 1217-25.
36. Protas EJ, Mitchell K, Williams A, Qureshy H, Caroline K, Lai EC. Gait and step training to reduce falls in Parkinson's disease. *NeuroRehabilitation* 2005; 20: 183-90.
37. Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc* 2006; 54: 743-9.

38. Rochester L, Rafferty D, Dotchin C, Msuya O, Minde V, Walker RW. The effect of cueing therapy on single and dual-task gait in drug naïve population of people with Parkinson's disease in northern Tanzania. *Mov Disord* 2010; 25: 906-11.
39. Canning CG, Ada L, Woodhouse E. Multiple-task walking training in people with mild to moderate Parkinson's disease: a pilot study. *Clin Rehabil* 2008; 22: 226-33.
40. Lamberti P, Armenise S, Castaldo V, de Mari M, Iliceto G, Tronci P, et al. Freezing gait in Parkinson's disease. *Eur Neurol* 1997; 38: 297-301.
41. Giladi N, Treves TA, Simon ES, Shabtai H, Orlov Y, Kandinov B, et al. Freezing of gait in patients with advanced Parkinson's disease. *J Neural Transm* 2001; 108: 53-61.
42. Schaafsma JD, Balash Y, Gurevich T, Bartels AL, Hausdorff JM, Giladi N. Characterization of freezing of gait subtypes and the response to each to levodopa in Parkinson's disease. *Eur J Neurol* 2003; 10: 391-8.
43. Hackney ME, Earhart GM. Effects of dance on balance and gait in severe Parkinson disease: a case study. *Disabil Rehabil* 2010; 32: 679-84.
44. Combs SA, Diehl MD, Staples WH, Conn L, Davis K, Lewis N, et al. Boxing training for patients with Parkinson disease: a case series. *Phys Ther* 2011; 91:132-42.
45. Pelosin E, Faelli E, Lofrano F, Avanzino L, Marinelli L, Bove M, Ruggeri P, Abbruzzese G. Effects of treadmill training on walking economy in Parkinson's disease: a pilot study. *Neurol Sci.* 2009;30:499-504.
46. Burini D, Farabollini B, Iacucci S, Rimatori C, Riccardi C, Riccardi G, Capecci M, Provinciali L, Ceravolo MG. A randomised controlled cross-over trial of aerobic training versus Qigong in advanced Parkinson's disease. *Eura Medicophys* 2006; 42: 231-8.

47. Mirelman A, Maidan I, Herman T, Deutsch JE, Giladi N, Hausdorff JM. Virtual reality for gait training: can it induce motor learning to enhance complex walking and reduce fall risk in patients with Parkinson's disease? *J Gerontol A Biol Med Sci* 2011; 66: 234-40.
48. Falvo MJ, Earhart GM. Six-minute walk distance in persons with Parkinson disease: a hierarchical regression model. *Arch Phys Med Rehabil* 2009; 90: 1004-8.
49. Caglar AT, Gurses HN, Mutluay FK, Kiziltan G. Effects of home exercises on motor performance in patients with Parkinson's disease. *Clin Rehabil* 2005; 19: 870-7.
50. Shumway-Cook A, Silver IF, LeMier M, York S, Cummings P, Koepsell TD. Effectiveness of a community-based multifactorial intervention on falls and fall risk factors in community-living older adults: a randomized, controlled trial. *J Gerontol A Biol Sci Med Sci* 2007; 62: 1420-7.
51. Fielding RA, Katula J, Miller ME, Abbott-Pillola K, Jordan A, Glynn NW, et al. Activity adherence and physical function in older adults with functional limitations. *Med Sci Sports Exerc* 1997; 39: 1997-2004.

Table 1. Participant Demographics

	Tango (n=26)	Control (n=26)	Tango Dropped (n=6)	Control Dropped (n=4)
Age (yrs)	69.3 ± 1.9 (48-89)	69.0 ± 1.5 (48-81)	76.2 ± 3.6 (60-86)	70.8 ± 6.2 (59-85)
Gender	15 male/11 female	15 male/11 female	4 male/2 female	1 male/3 female
Years with PD	5.8 ± 1.1 (1-20)	7.0 ± 1.0 (1-21)	3.5 ± 1.3 (0.5-9)	6.0 ± 1.5 (2.5-10)
Hoehn & Yahr Stage	2.6 ± 0.1 (1-4)	2.5 ± 0.1 (2-4)	2.4 ± 0.2 (2-3)	2.8 ± 0.4 (2-4)

* Values are means ± SEs (ranges). Right two columns provide demographic information for those participants who were lost to follow-up after the baseline visit and prior to the 3 month visit.

Table 2. MDS-UPDRS-III Means \pm SEs (95% CI)

MDS-UPDRS-III	Tango (n=26)	Control (n=26)
Baseline	44.5 \pm 2.3 (37-53)	48.0 \pm 1.8 (45-56)
3 Months	39.9 \pm 2.3 (28-45)	45.6 \pm 1.8 (38-49)
6 Months	34.2 \pm 2.2 (28-38)	45.2 \pm 1.9 (41-50)
12 Months	31.7 \pm 2.4 (24-36)	45.0 \pm 1.9 (39-48)




Figure 1. Consort flow diagram illustrating participant recruitment, randomization, and tracking over the course of the study. Note that the final analyzed sample included all participants retained through 3 months, with those who dropped out at 6 or 12 months remaining in the sample through use of a last observation carried forward intent to treat analysis.




Figure 2. Scores on MDS-UPDRS-1 non-motor symptoms scale (A), MDS-UPDRS-2 ADLs scale (B), and MDS-UPDRS-3 motor symptoms scale (C) at baseline, 3, 6, and 12 month evaluations for the Tango (filled circles) and Control (open circles) groups. Values are means \pm SEs. Evaluations were conducted with participants off medication. Asterisks denote significant differences between Tango and Control within that time point. Each horizontal line indicates a significant difference within Tango between the two time points spanned by the line. For complete statistical comparison results please see text.




Figure 3. Scores on MDS-UPDRS-3 individual motor components of tremor (A), rigidity (B), bradykinesia (C), and PIGD (D) at baseline, 3, 6, and 12 month evaluations for the Tango (filled circles) and Control (open circles) groups. Values are means \pm SEs. Evaluations were conducted with participants off medication. Asterisks denote significant differences between Tango and Control within that time point. Each horizontal line indicates a significant difference within Tango between the two time points spanned by the line. For complete statistical comparison results please see text.




Figure 4. Scores on the nine hole peg test (A), MiniBESTest balance test (B), freezing of gait questionnaire (C), and six minute walk test (D) at baseline, 3, 6, and 12 month evaluations for the Tango (filled circles) and Control (open circles) groups. Values are means \pm SEs. Asterisks denote significant differences between Tango and Control within that time point. Each horizontal line indicates a significant difference within Tango between the two time points spanned by the line. For complete statistical comparison results please see text.




Figure 5. Walking velocities for forward (A), fast as possible (B), dual task (C), and backward (D) walking at baseline, 3, 6, and 12 month evaluations for the Tango (filled circles) and Control (open circles) groups. Values are means \pm SEs. Asterisks denote significant differences between Tango and Control within that time point. Each horizontal line indicates a significant difference within Tango between the two time points spanned by the line. For complete statistical comparison results please see text.