

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2017

Comparisons of ELISA and Western blot assays for detection of autophagy flux

Sung-hee Oh
University of Suwon

Yong-bok Choi
University of Suwon

June-hyun Kim
University of Suwon

Conrad C. Wehl
Washington University School of Medicine in St. Louis

Jeong-sun Ju
University of Suwon

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Oh, Sung-hee; Choi, Yong-bok; Kim, June-hyun; Wehl, Conrad C.; and Ju, Jeong-sun, "Comparisons of ELISA and Western blot assays for detection of autophagy flux." *Data in Brief*. 13, 696-699. (2017).
https://digitalcommons.wustl.edu/open_access_pubs/6039

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Comparisons of ELISA and Western blot assays for detection of autophagy flux

Sung-hee Oh ^a, Yong-bok Choi ^b, June-hyun Kim ^b,
Conrad C. Wehl ^c, Jeong-sun Ju ^{a,*}^a Department of Exercise Science, Research Institute of Sports Science, the University of Suwon, 17 Wauan-gil, Bongdam-eup, Hwaseong-si, Gyeonggi-do 18322, South Korea^b Department of Biotechnology and Bioscience, School of Bioindustry, the University of Suwon, 17 Wauan-gil, Bongdam-eup, Hwaseong-si, Gyeonggi-do 18322, South Korea^c Department of Neurology, Washington University School of Medicine, PO Box 8111, 660 South Euclid Avenue, St Louis, MO 63110, USA

ARTICLE INFO

Article history:

Received 3 May 2017

Received in revised form

14 June 2017

Accepted 28 June 2017

Available online 4 July 2017

Keywords:

Autophagy

Mitophagy

ELISA

Western blot

Skeletal muscle

ABSTRACT

We analyzed autophagy/mitophagy flux *in vitro* (C2C12 myotubes) and *in vivo* (mouse skeletal muscle) following the treatments of autophagy inducers (starvation, rapamycin) and a mitophagy inducer (carbonyl cyanide m-chlorophenylhydrazone, CCCP) using two immunodetection methods, ELISA and Western blotting, and compared their working range, accuracy, and reliability. The ELISAs showed a broader working range than that of the LC3 Western blots (Table 1). Table 2 showed that data value distribution was tighter and the average standard error from the ELISA was much smaller than those of the Western blot, directly relating to the accuracy of the assay. Test-retest reliability analysis showed good reliability for three individual ELISAs (interclass correlation, ≥ 0.7), but poor reliability for three individual Western blots (interclass correlation, ≤ 0.4) (Table 3).

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.ab.2017.05.003>

* Corresponding author.

E-mail address: jju625@hotmail.com (J.-s. Ju).<http://dx.doi.org/10.1016/j.dib.2017.06.045>2352-3409/© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

| | |
|----------------------------|--|
| Subject area | Biochemistry |
| More specific subject area | Assay development and method |
| Type of data | Tables |
| How data was acquired | enzyme-linked immunosorbent assay (ELISA), Western blotting |
| Data format | Descriptive data: mean \pm SD, analyzed |
| Experimental factors | C2C12 myotubes and male wild-type C57BL/6 mouse skeletal muscle |
| Experimental features | <i>in vitro</i> and <i>in vivo</i> autophagy/mitophagy flux were measured using two immunodetection techniques |
| Data source location | Hwaseong, South Korea |
| Data accessibility | Data are available with this article |

Value of the data

- The presented data indicated that the ELISA had smaller data distribution and was more repeatable to measure autophagy/mitophagy flux, compared with Western blot data.
- These data could be helpful for many autophagy researchers to obtain more accurate and reproducible data using this ELISA technique.
- These data could also be beneficial for researchers in other areas to adapt the ELISA-based assay strategy from the Western blot.

1. Data

We compared the ELISA with Western blot data from C2C12 myotubes and male wild-type C57BL/6 mouse skeletal muscle tissue in terms of the working range, accuracy, and reliability for autophagy/mitophagy flux measurements. The ELISAs could surpass the Western blot in all criteria for quantification of autophagy flux. The ELISAs showed a broader dynamic range than the Western blot (5.3 *versus* 1.4, the ratio of the highest O.D. value to the lowest) (Table 1). The values of standard error from the ELISA were much smaller than those of the Western blot ($P < 0.05$, Table 2), which implies the accuracy of the assays. When each of the three individual assays was compared, the ELISA was more reliable than the Western blot (interclass correlation, ≥ 0.7 *versus* ≤ 0.4) (Table 3).

Table 1

The working range of the ELISA and the LC3 Western blot.

| ELISA | Blanks | Lowest O.D. | Highest O.D. |
|---------------------|-------------------|-------------------|-------------------|
| Average values | 0.03 \pm 0.002 | 0.183 \pm 0.028 | 0.961 \pm 0.031 |
| Ratio to blanks | | 6 | 32 |
| Dynamic range | | | 5.3 ¹ |
| Western blot | Background | Lowest density | Highest density |
| Average values | 0.071 \pm 0.005 | 0.301 \pm 0.093 | 0.407 \pm 0.052 |
| Ratio to background | | 4.2 | 5.7 |
| Dynamic range | | | 1.4 ¹ |

Values are means \pm SD; $n = 18$ /group.

¹ The ratio of the highest O.D. value to the lowest.

Table 2

Comparisons of average standard errors of data from C2C12 cells and TA muscles treated with starvation, rapamycin, and CCCP.

| Average Standard Error | | | |
|------------------------|-------------------|------------------|-------------------|
| C2C12 Cells | Starvation | Rapamycin | CCCP |
| Western blot | 0.180 ± 0.082 | 0.240 ± 0.088 | 0.172 ± 0.037 |
| ELISA | 0.07 ± 0.009 | 0.161 ± 0.013 | 0.018 ± 0.005 |
| Fold difference | 2.6 [*] | 1.5 [*] | 9.6 [*] |
| TA muscles | Starvation | Rapamycin | CCCP |
| Western blot | 0.778 ± 0.105 | 0.301 ± 0.093 | 0.407 ± 0.052 |
| ELISA | 0.041 ± 0.014 | 0.042 ± 0.006 | 0.032 ± 0.003 |
| Fold difference | 19.0 [*] | 7.2 [*] | 12.7 [*] |

Values are means ± SD; $n = 18$ /group. Difference was evaluated using a paired t -test at $P < 0.05$.

*Fold change compared to control groups.

Table 3

Test-retest reliability for three inter-assays.

| 3 ELISAs | ICC | 95% CI |
|------------------------|------------|---------------|
| Rapamycin | 0.889 | 0.53, 0.98 |
| CCCP | 0.778 | 0.06, 0.966 |
| 3 Western blots | ICC | 95% CI |
| Starvation | 0.3804 | −1.63, 0.91 |
| Rapamycin | 0.234 | −2.25, 0.88 |
| CCCP | 0.077 | −2.911, 0.86 |

ICC: intraclass correlation, CI: confidence interval.

2. Experimental design, materials and methods

We performed the autophagic flux assays using *in vitro* and *in vivo* LC3-II turnover measured by Western blot in the presence and absence of an autophagy inhibitor as previously described [1,2] and the ELISA using LC3 antibodies following subcellular fractionation of mouse skeletal muscle cells and tissue as described in a recent paper [3].

2.1. Measurement of autophagic flux in cultured cells

C2C12 myotubes grown on 10-cm dishes were incubated in amino acid and serum-free starvation buffer or treated with 10 µg/mL rapamycin or 25 µg/mL CCCP with and without 200 nM BafilomycinA1 for 8 h.

2.2. Measurement of autophagy flux in animals

Ten week old male wild-type C57BL/6 mice were divided into four groups: fed and starvation, or vehicle, rapamycin, and CCCP. Starvation was performed by removing food for 48 h. 10 mg/kg/day rapamycin, 4 mg/kg/day CCCP or vehicle was *i.p.* injected to mice daily for two days. Mice were also treated with and without 0.4 mg/kg/day colchicine administration for 48 h. Control mice received an equal volume of *i.p.* saline.

2.3. Membrane/cytosol fractionation

Subcellular fractionations from cultured cells or tissue were performed as previously described [4].

2.4. Mitochondrial/cytosol fractionation

Mitochondrial fractionations from cultured cells or tissue were performed as previously described [5].

2.5. Sandwich ELISA assay

Following subcellular fractionation of mouse skeletal muscle cells and tissue, cytosolic, membrane, and mitochondrial fractions were analyzed through a sandwich ELISA using two LC3 antibodies, LC3 capture and HRP-conjugated LC3 detection antibodies as previously described [3].

2.6. Immunoblot analysis

LC3 protein turnover and other protein levels were measured using immunoblot techniques as previously described [3].

2.7. Statistical analysis

Data are presented as means \pm SD and were evaluated by a paired t-test or reliability analysis at $p < 0.05$.

Funding

This work was supported by the National Research Foundation of Korea Grant funded by the Korean Government (NRF-2013-332-2013S1A5A8021905).

Acknowledgements

The authors thank Yolanda Mathews for critical reading of the manuscript.

Transparency document. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2017.06.045>.

References

- [1] J.S. Ju, A.S. Varadhacharty, S.E. Miller, C.C. Wehl, Quantification of “autophagic flux” in mature skeletal muscle, *Autophagy* 6 (2010) 929–935.
- [2] J.S. Ju, S.I. Jeon, J.Y. Park, J.Y. Lee, S.C. Lee, K.J. Cho, J.M. Jeong, Autophagy plays a role in skeletal muscle mitochondrial biogenesis in an endurance exercise-trained condition, *J. Physiol. Sci.* 66 (2016) 417–430.
- [3] S.H. Oh, Y.B. Choi, J.H. Kim, C.C. Wehl, J.S. Ju, Quantification of autophagy flux using LC3 ELISA, *Anal. Biochem.: Methods Biol. Sci.* 530 (2017) 57–67.
- [4] S. Baghirova, B.G. Hughes, M.J. Hendzel, R. Schulz, Sequential fractionation and isolation of subcellular proteins from tissue or cultured cells, *MethodsX* 2 (2015) 440–445.
- [5] I. Dim Mauro, T. Pearson, D. Caporossi, M.J. Jackson, A simple protocol for the subcellular fractionation of skeletal muscle cells and tissue, *BMC Res. Notes* 5 (2010) 1–5.