

Washington University School of Medicine

Digital Commons@Becker

---

Conference Abstracts and Posters

Division of Emergency Medicine/Emergency  
Care Research Section

---

2010

## Comparing throughput of physician extenders in various volume EDs

Lawrence Lewis

*Washington University School of Medicine in St. Louis*

Melanie Sutter

*Washington University School of Medicine in St. Louis*

Follow this and additional works at: [https://digitalcommons.wustl.edu/em\\_conf](https://digitalcommons.wustl.edu/em_conf)

---

### Recommended Citation

Lewis, Lawrence and Sutter, Melanie, "Comparing throughput of physician extenders in various volume EDs" (2010). *Conference Abstracts and Posters*. Paper 5.  
[https://digitalcommons.wustl.edu/em\\_conf/5](https://digitalcommons.wustl.edu/em_conf/5)

This Presentation Poster is brought to you for free and open access by the Division of Emergency Medicine/ Emergency Care Research Section at Digital Commons@Becker. It has been accepted for inclusion in Conference Abstracts and Posters by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

## Comparing Throughput of Physician Extenders in Various Volume EDs

Larry Lewis MD, Melanie Sutter MD, MSCI

Division of Emergency Medicine, Washington University School of Medicine, St. Louis, Missouri



### BACKGROUND

Nurse practitioners (NP) and physician assistants (PA) are commonly utilized in busy emergency departments (ED). Each has a separate training program but typically see similar types of patients within the ED.

### OBJECTIVES

To evaluate the efficiency of NPs and PAs in the ED setting based on ED volume.

Hypothesis: We believe that PAs based on the similarities in their training program to that of MD/DOs will have increased levels of efficiency regardless of ED volume.

### METHODS

We did a 3 year retrospective review of data from 41 EDs, varying in size from 15,000 to > 45,000 visits/yr. Data from NP's (N=588) and PA's (N=1314) were stratified by ED volume: 15-30,000 (low); 30-45,000 (mid); or > 45,000 visits/yr (high).


### STATISTICAL METHODS

Median time-to-decision (TTD) (triage time to disposition time) in minutes with interquartile ranges was compared between PAs and NPs for each ED volume using a t-test. Mean number of patients/hr (with 95% CI) were compared between PAs and NPs for each category of ED volume using ANOVA. Statistical significance was set at  $p < 0.05$ .


### RESULTS

Median TTD for NPs was greater than PAs: 2.30, 2.20, and 2.17 hrs for low, mid, and high volume EDs respectively vs. 1.95, 1.88, and 2.05 hrs ( $p < 0.001$ ,  $p = 0.002$ ,  $p = 0.009$  for each volume). TTD differed significantly with ED volume for NPs ( $p = 0.030$ ), but not for PAs ( $p = 0.349$ ). Mean number of pts/hr for NPs was 1.32, 1.73, and 1.62 for low, mid, and high volume EDs vs. 1.52, 1.71, and 1.90 pts/hr for PAs ( $p < 0.001$ ,  $p = 0.589$ ,  $p < 0.001$  respectively). There was a significant difference by ED volume for both NPs and PAs ( $p < 0.001$  for both).

Time-to-decision (TTD) for NPs vs. PAs



Mean number of patients/hour for NPs vs. PAs



### CONCLUSIONS

ED volume is related to throughput by TTD and patients/hr; however, PAs had shorter TTD and saw more patients/hr than NPs in most settings.

### LIMITATIONS

Patient diagnosis and consultant use were not included in the study, which can both affect the overall TTD. Years in practice of the practitioners was not included. This could have an effect on both TTD and patients/hr.

### DISCLOSURES

We have no disclosures and did not receive any funding for this study.

### ACKNOWLEDGEMENTS

We would like to thank Dr. Christopher Richter for compiling the study information into a database and for all of his work on the statistical aspects of the study.