

Washington University School of Medicine

Digital Commons@Becker

Posters

2010: Disentangling the Genetics of Alcoholism:
Understanding Pathophysiology and Improving
Treatment

2010

The association between Generalized Anxiety Disorder and alcohol abuse and dependence

Kevin Y. Chung

Washington University School of Medicine in St. Louis

Alexis E. Duncan

Washington University in St Louis

Kathleen Keenan Bucholz

Washington University School of Medicine in St. Louis

Follow this and additional works at: <https://digitalcommons.wustl.edu/guzeposter2010>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Chung, Kevin Y.; Duncan, Alexis E.; and Bucholz, Kathleen Keenan, "The association between Generalized Anxiety Disorder and alcohol abuse and dependence" (2010). *Posters*. Paper 4 Samuel B. Guze Symposium on Alcoholism.

<https://digitalcommons.wustl.edu/guzeposter2010/4>

This Poster is brought to you for free and open access by the 2010: Disentangling the Genetics of Alcoholism: Understanding Pathophysiology and Improving Treatment at Digital Commons@Becker. It has been accepted for inclusion in Posters by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

THE ASSOCIATION BETWEEN GENERALIZED ANXIETY DISORDER AND ALCOHOL ABUSE AND DEPENDENCE

Kevin Y. Chung, Alexis E. Duncan,
Kathleen Keenan Bucholz

Washington University Department of Psychiatry, Midwest
Alcoholism Research Center

George Warren Brown School of Social Work, Washington
University



Supported by NIH Grant AA12640



BACKGROUND

- Generalized Anxiety Disorder (GAD) is a chronic disorder that is often comorbid with other psychiatric disorders (Ballenger et al., 2001; Grant, Hasin, Stinson, et al. 2005)
 - The association between GAD and Alcohol Use Disorders (AUD) has received attention due to the possibility that individuals with GAD may resort to alcohol for self-medication (Grant et al., 2005)
 - In a population based study, 65% of males and 32.8% of females with a lifetime GAD diagnosis also had a lifetime AUD diagnosis (Vesga-Lopez et al. 2008)
-

OBJECTIVE

To characterize the association between GAD and alcohol use disorders, and to determine whether depression is a moderator of this association.

METHODS - SAMPLE

- Family study of Missouri residents (MOFAM)
 - Over-sampled for African American (AfAm) race (>50% AA)
 - Three risk groups based on paternal excessive alcohol use:
 - Recurrent drunk driving (RDD; ascertained from driving records), Maj. n=267, AfAm n=151
 - High risk (mother of children reported father drank excessively), Maj. n=147, AfAm n=210
 - Control (drawn from general population, irrespective of paternal alcohol status), Maj n=190, AfAm n=319
-

METHODS – Data Analysis

- Dependent Variable: Alcohol Abuse and Dependence
 - Independent Variable: GAD
 - Covariates: Race, age (<18y), Sex, Income (<\$45,000), Regular Marijuana Use, Marijuana Abuse and Dependence, Regular Smoker, Nicotine Dependence, Social Phobia, Panic Attacks (>3), Panic Disorder
 - Potential Effect Modifier: Major Depressive Disorder (MDD)
-

METHODS - Data Analysis

- Bivariable relationships were assessed using the chi-square statistic
 - The Breslow-Day test for homogeneity was used to test for effect modification by MDD
 - Logistic Regression was employed to construct a multivariable model
 - All covariates were assessed as confounders: A variable was considered a confounder if its addition to the model resulted in a >10% change in the OR for GAD
-

RESULTS: Table 1. Characteristics of MOFAM sample by GAD diagnosis.

	GAD Diagnosis (n=78)	No GAD Diagnosis (n=1199)	P-value
African American	61.54	52.21	0.110
Family type			0.064
Repeat drunk driving	32.05	32.53	
High risk	38.46	27.19	
Control	29.49	40.28	
Age <18y	42.31	57.38	0.009
Female	58.97	50.04	0.126
Income (<\$45,000)	60.0	55.34	0.431

Regular Marijuana User	20.51	16.01	0.297
Marijuana Abuse and Dependence	21.79	5.50	<0.001
Regular Smoking	32.05	18.85	0.005
Nicotine Dependence	28.21	13.76	0.001
Major Depressive Disorder	37.18	8.59	<0.001
Social Phobia	57.69	20.85	<0.001
Panic Attack*	38.46	6.92	<0.001
Panic Disorder	10.26	1.08	<0.001

*At least 4 panic attack episodes that peak within 10 minutes

RESULTS – Interaction between GAD and Major Depression

- Using the Breslow-Day test, we identified a significant interaction between GAD and major depression ($p=.028$)
 - Therefore, the interaction was modeled as a set of dummy variables:
 - GAD with MDD (GAD+MDD+) $n=29$
 - GAD without MDD (GAD+MDD-) $n=49$
 - MDD without GAD (GAD-MDD+) $n=103$
-

RESULTS: Table 2. Logistic Regression Model predicting AUD by GAD and MDD status

	OR (95% CI)	
	Crude	Adjusted*
GAD+MDD+	2.075 (.904-4.762)	0.547 (0.194-1.543)
GAD+MDD-	2.403 (1.281-4.508)	1.774 (0.816-3.854)
GAD-MDD+	2.927 (1.891-4.530)	1.966 (1.151-3.357)

*Adjusted for race, family status, age, sex, marijuana abuse and dependence, nicotine dependence and social phobia

RESULTS: SUMMARY

- In unadjusted analyses, GAD without depression and depression without GAD are significantly positively associated with AUD
 - After controlling for relevant confounders (race, family status, age, sex, marijuana abuse and dependence, nicotine dependence, and social phobia) the strength of the association is attenuated for GAD+MDD- and GAD-MDD+, and that for GAD+MDD+ changes direction, although only the OR for GAD-MDD+ is statistically significant
-

CONCLUSIONS

- The relationship between GAD and AUD may be moderated by MDD such that having GAD+MDD+ is negatively associated with AUD and having GAD+MDD- is positively associated with AUD
 - Due to a relatively small number of individuals with GAD in our sample, we lacked statistical power to detect significant effects of this magnitude
 - Attempts should be made to replicate these results in larger samples
-

REFERENCES

Ballenger JC, et al. (2001). Consensus statement on generalized anxiety disorder from the International Consensus Group on Depression and Anxiety. *Journal of Clinical Psychiatry*; 62:53-58

Grant BF, Hasin DS, Stinson FS, et al. (2005). Prevalence, correlates, co-morbidity, and comparative disability of DSM-IV generalized anxiety disorder in the USA: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychological Medicine*; 35:1747-1759.

Vesga-Lopez, O, et al. (2008). Gender differences in generalized anxiety disorder: results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Journal of Clinical Psychiatry*; 69:1606-1616
