

2003

Effects of childhood exposure to paternal alcoholism on substance use disorders in adolescents and young adults

A. E. Duncan

Q. Fu

Kathleen K. Bucholz

J. F. Scherrer

T. Jacob

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.wustl.edu/guzeposter2003>

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Duncan, A. E.; Fu, Q.; Bucholz, Kathleen K.; Scherrer, J. F.; Jacob, T.; and True, W. R., "Effects of childhood exposure to paternal alcoholism on substance use disorders in adolescents and young adults" (2003). *Posters*. Paper 30 Samuel B. Guze Symposium on Alcoholism.

<http://digitalcommons.wustl.edu/guzeposter2003/30>

This Poster is brought to you for free and open access by the 2003: Drinking and the High School Student at Digital Commons@Becker. It has been accepted for inclusion in Posters by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

Authors

A. E. Duncan, Q. Fu, Kathleen K. Bucholz, J. F. Scherrer, T. Jacob, and W. R. True

EFFECTS OF CHILDHOOD EXPOSURE TO PATERNAL ALCOHOLISM ON SUBSTANCE USE DISORDERS IN ADOLESCENTS AND YOUNG ADULTS

A.E. Duncan, Q. Fu, K.K. Bucholz, J.F. Scherrer, T.
Jacob, and W.R. True



OBJECTIVE

To determine whether exposure to paternal alcoholism in the first 12 years of life is related to the development of substance use disorders

METHODS

COA Study

- Fathers
 - 1464 twin fathers sampled from the Vietnam Era Twin Registry
 - All had at least 1 child 12-26 yrs old in 2000
 - Twin pairs either concordant or discordant for alcohol dependence. Controls were non-alcoholic twin pairs
- Mothers
 - 1,000 biological and/or rearing (i.e. lived regularly with child between ages 6 and 13) mother identified, 864 interviewed
 - Mothers asked to give permission to interview offspring
- Offspring
 - Mothers identified and gave permission for 1487 offspring to be interviewed. 1,270 offspring of 730 fathers were interviewed.

Assessment Domains

- Twin fathers' alcohol lifetime Dx and zygosity obtained from '92 data (Tsuang and Lyons).
- 1,464 twin fathers sampled with 84% response rate (n=1176). Updated lifetime drinking history (LDH) collected from twin fathers by telephone interview.
- Mothers and offspring were interviewed by telephone and completed a questionnaire by mail to provide:
 - Telephone interview: Alcohol abuse, dependence, lifetime drinking history (offspring only), substance use/abuse, psychopathology (depression, anxiety, conduct disorder, oppositional defiant disorder, attention deficit/hyperactivity disorder)
 - Mail questionnaire: psychosocial variables (family & relationship quality, peers, school attendance, work environment, personality, drinking expectancies, etc.)

Data Analysis

- Latent class analysis was done using LCAP (<http://hardy.wustl.edu>) to group children of alcoholics by pattern of exposure to paternal alcoholism
- STATA was used to develop Cox Proportional Hazards models for risk of developing alcohol abuse/dependence, marijuana abuse/dependence and nicotine dependence and adjust for confounding variables and familial clustering

Latent Class Analysis

- Statistical method for uncovering patterns or structure in multivariate categorical data (like factor analysis for categorical data)
- Defines “clusters” or “classes” of cases that share similar characteristics
- Cases placed in clusters based upon membership probabilities


Cox Proportional Hazards Models

- Survival analysis – accounts for the possibility that some subjects have not yet had time to develop the outcome
- Assumption that risk remains constant over time
- Hazard ratios can be thought of like odds ratios

LCA for child exposure to paternal alcoholism

- COA's only (n=618)
- Father's lifetime drinking history (LDH) matched to child's lifetime
- Binary variables created indicating whether or not father met criteria for alcohol abuse or dependence during each of the first 12 years of the child's life
- 4 class solution was selected

Child exposure to paternal alcoholism LCA – 4 class solution



Characteristics of the sample

	<u>Latent class for exposure to paternal alcoholism</u>				
	Non-alcoholic	None	Chronic	Early	Late
N (%)	652 (51.3)	264 (20.8)	195 (15.4)	114 (9.0)	45 (3.5)
% Male	47.70	52.65	46.15	42.11	60.00
Age:					
12-16 yrs	28.22	38.65	18.46	29.82	20.00
17-21 yrs	37.88	35.98	34.87	38.60	31.11
22-28 yrs	33.90	25.38	46.67	31.58	48.89

Proportion of offspring with fathers with lifetime psychiatric and drug use disorders

DSM-III-R Disorder	<u>Latent class for exposure to paternal alcoholism</u>				
	Non-alcoholic	None	Chronic	Early	Late
N (%)	652 (51.3)	264 (20.8)	195 (15.4)	114 (9.0)	45 (3.5)
Drug abuse or dependence	6.6	22.4	18.0	16.7	13.3
ASPD	6.2	11.0	13.3	16.7	15.6
Depression	4.8	11.7	13.9	10.5	22.2

Offspring substance use disorders and other psychopathology

	<u>Latent class for exposure to paternal alcoholism</u>				
	Father not alcoholic	None	Chronic	Early	Late
N (%)	652 (51.3)	264 (20.8)	195 (15.4)	114 (9.0)	45 (3.5)
Alcohol abuse/ dependence	25.27	22.35	38.86	35.96	40.00
Marijuana abuse/ dependence	12.27	13.64	13.33	17.54	11.11
Nicotine Dependence	9.23	10.61	11.92	15.04	8.89
Conduct Disorder	10.68	14.50	12.95	12.73	16.67
ODD	15.36	18.18	15.46	18.42	28.29
Depression	17.64	21.97	16.92	24.56	26.67
Social Phobia	8.44	8.71	7.18	12.28	15.56

Cox proportional hazards models for alcohol abuse/dependence – adjusting for gender, age cohort, and clustering

	Hazard Ratios (95% CI) *			
	None	Chronic	Early	Late
All	1.10 (0.81, 1.51)	1.44 (1.09, 1.90)	1.93 (1.31, 2.84)	1.57 (1.00, 2.46)
Male	1.20 (0.81, 1.79)	1.46 (1.01, 2.09)	1.45 (0.84, 2.51)	1.80 (1.06, 3.05)
Female	1.03 (0.64, 1.66)	1.46 (0.95, 2.24)	2.35 (1.39, 3.98)	1.19 (0.46, 3.09)

*Comparison group: children of non-alcoholics

Cox proportional hazards models for marijuana abuse/dependence – adjusting for gender, age cohort, and clustering

	Hazard Ratios (95% CI)*			
	None	Chronic	Early	Late
All	1.21 (0.79, 1.84)	1.06 (0.65, 1.72)	1.62 (0.94, 2.82)	0.65 (0.24, 1.75)
Males	1.63 (0.99, 2.66)	0.99 (0.54, 1.80)	1.17 (0.50, 2.71)	0.70 (0.22, 2.22)
Females	0.69 (0.31, 1.52)	1.19 (0.59, 2.40)	2.03 (0.96, 4.31)	0.54 (0.07, 4.00)

*Comparison group: children of non-alcoholics

Cox proportional hazards models for nicotine dependence – adjusting for gender, age cohort, and clustering

	Hazard Ratios (95% CI)*			
	None	Chronic	Early	Late
All	1.34 (0.68, 1.21)	1.15 (0.82, 2.17)	1.90 (1.07, 3.38)	0.86 (0.30, 2.49)
Male	1.22 (0.62, 2.39)	1.20 (0.61, 2.36)	2.07 (0.42, 2.72)	0.34 (0.04, 5.79)
Female	1.49 (0.75, 2.96)	1.12 (0.53, 2.37)	2.79 (1.31, 5.94)	1.66 (0.48, 5.79)

*Comparison group: children of non-alcoholics

Cox proportional hazards models for alcohol abuse/dependence before and after adjusting for father psychopathology*

		Hazard Ratios (95% CI)**			
		None	Chronic	Early	Late
Males					
Before		1.20 (0.81, 1.79)	1.46 (1.01, 2.09)	1.45 (0.84, 2.51)	1.80 (1.06, 3.05)
After		1.24 (0.82, 1.87)	1.45 (0.99, 2.11)	1.46 (0.84, 2.55)	1.71 (0.95, 3.07)
Females					
Before		1.03 (0.64, 1.66)	1.46 (0.95, 2.24)	2.35 (1.39, 3.98)	1.19 (0.46, 3.09)
After		0.96 (0.59, 1.57)	1.39 (0.90, 2.14)	2.26 (1.32, 3.87)	1.14 (0.44, 2.96)

*drug abuse/dependence, major depression and ASPD

** Comparison group: children of non-alcoholics

Cox proportional hazards models for marijuana abuse/dependence before and after adjusting for father psychopathology*

		Hazard Ratios (95% CI)			
		None	Chronic	Early	Late
Males					
Before		1.63 (0.99, 2.66)	0.99 (0.54, 1.80)	1.17 (0.50, 2.71)	0.70 (0.22, 2.22)
After		1.39 (0.82, 2.36)	0.87 (0.48, 1.60)	0.92 (0.39, 2.16)	0.59 (0.19, 1.37)
Females					
Before		0.69 (0.31, 1.52)	1.19 (0.59, 2.40)	2.03 (0.96, 4.31)	0.54 (0.07, 4.00)
After		0.62 (0.28, 1.38)	1.15 (0.56, 2.34)	2.01 (0.92, 4.36)	0.45 (0.06, 3.28)

*drug abuse/dependence, major depression and ASPD

** Comparison group: children of non-alcoholics

Cox proportional hazards models for nicotine dependence before and after adjusting for father psychopathology*

		Hazard Ratios (95% CI)			
		None	Chronic	Early	Late
Males					
Before	1.22 (0.62, 2.39)	1.20 (0.61, 2.36)	2.07 (0.42, 2.72)	0.34 (0.04, 5.79)	
After	1.15 (0.60, 2.22)	1.02 (0.53, 1.98)	0.95 (0.35, 2.52)	0.26 (0.03, 2.19)	
Females					
Before	1.49 (0.75, 2.96)	1.12 (0.53, 2.37)	2.79 (1.31, 5.94)	1.66 (0.48, 5.79)	
After	1.30 (0.65, 2.62)	0.99 (0.47, 2.07)	2.55 (1.15, 5.66)	1.68 (0.49, 5.78)	

*drug abuse/dependence, major depression and ASPD

** Comparison group: children of non-alcoholics

CONCLUSIONS

- After adjusting for age cohort and father drug abuse/dependence, depression, and ASPD, exposure to paternal alcoholism in the first 12 years of life is associated with over a 2-fold risk of alcohol abuse/dependence and nicotine dependence in females, when exposure to paternal alcoholism is early in the first 12 years of life.

Future analyses

- Explore the role of maternal alcoholism, depression, and ASPD symptoms
- Consider the role of paternal alcoholism severity
- Devise a variable for child contact with father and use as a covariate
- Examine the role of exposure during the teen years

We would like to acknowledge:

...the **Support of NIAAA** for

- Grant P50-AA11998 to Andrew Heath for the Missouri Alcoholism Research Center
- Grant R01-AA11667 to Theodore Jacob for Adult Children of Alcoholism Discordant Twins, and
- Grant R01-AA11822 to William True for Adolescent Children of Alcoholics: A Twin Family Design.