2014

Do comorbid ulnar symptoms or ulnar neuropathy affect the prognosis of workers with carpal tunnel syndrome?

Alexis Descatha

Univ Versailles St-Quentin

Ann Marie Dale

Washington University School of Medicine in St. Louis

Angelique Zeringue

Washington University School of Medicine in St. Louis

Bradley Evanoff

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/ohs_facpubs

Recommended Citation

This is a non-final version of an article published in final form in Descatha A, Dale AM, Zeringue A, Evanoff B. Do Co-Morbid Ulnar Symptoms Or Ulnar Neuropathy Affect The Prognosis Of Workers With Carpal Tunnel Syndrome? *Journal of occupational and environmental medicine / American College of Occupational and Environmental Medicine*. 2014;56(3):e2-e3. doi:10.1097/JOM.000000000000118.

This Article is brought to you for free and open access by the Occupational Health and Safety at Digital Commons@Becker. It has been accepted for inclusion in OHS Faculty Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Readers are invited to submit letters for publication in this department. Submit letters online at <http://joem.edmgr.com>. Choose "Submit New Manuscript." A signed copy-right assignment and financial disclosure form must be submitted with the letter. Form available at www.joem.org under Author and Reviewer information.

Do Comorbid Ulnar Symptoms or Ulnar Neuropathy Affect the Prognosis of Workers With Carpal Tunnel Syndrome?

To the Editor:

Carpal tunnel syndrome (CTS) is a common and costly disease among working-aged adults, and a leading cause of work-related disability¹ and has affected almost five million US workers with prevalence estimated between 3.1% and 7.8%.² Although prognosis after surgery has been described in previous studies,^{3,4} many questions exist about predictors of CTS prognosis in working populations.⁵ The presence of ulnar symptoms or neuropathy have been considered predictive of a poor outcome in CTS cases. We aimed to describe a 3-year evolution of CTS with and without ulnar symptoms in a large cohort of workers.

This study presents data collected prospectively on 1107 newly hired workers in the United States recruited between July 2004 and October 2006 from eight employers and three trade unions, representing manufacturing, construction, biotechnology, and health care. Subjects were followed for 3 years ($n = 888$) and came from three main occupational groups: construction apprentices, office and laboratory workers, and service workers/housekeepers.⁶ At baseline, subjects answered questions about the presence of hand pain in the past year (occurring three times or lasting at least 1 week) and about the nature and location of symptoms.

This study was supported by CDC/NIOSH (grant # R01OH008017-01) and by the National Center for Research Resources (NCRR) (grant # UL1 RR024992), a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of NIOSH, NCRR or NIH. NEUROMetrix provided electrodes for nerve conduction testing.

The authors declare no other conflicts of interest. Address correspondence to: Alexis Descatha, MD, AP-HP, Occupational Health Unit, University hospital of West suburb of Paris, F92380 Garches, France (alexis.descatha@uvsq.fr). Copyright © 2014 by American College of Occupational and Environmental Medicine DOI: 10.1097/JOM.0000000000000118

Bilateral nerve conduction studies were performed for median and ulnar nerves at the wrist, using the NC-Stat automated nerve conduction testing device (NEUROMetrix, Inc, Waltham, MA). The NC-Stat device followed an automated testing protocol to measure median and ulnar distal motor latencies (wrist-thenar eminence and wrist-hypothenar eminence) and distal sensory latencies (wrist-third finger and wrist-fifth finger).⁷ Abnormal median nerve conduction was defined as sensory latency more than 3.5 ms (14 cm) or motor latency more than 4.5 ms or median-ulnar sensory latency difference of more than 0.5 ms (14 cm). For the ulnar nerve, we used sensory or motor latency more than the 95th percentile. Subjects were categorized according to baseline data into mutually exclusive groups (see Table 1). Workers with symptoms of CTS with and without fifth finger involvement were also described, including those with confirmed CTS and confirmed ulnar syndrome (symptoms and abnormal nerve studies).⁴

Three outcomes were used at follow-up: "severe hand pain," defined as hand pain within the past 30 days with a rating of 5 or higher on a scale of 0 (no discomfort) to 10 (worst discomfort imaginable); "functional status limitations," assessed via the Levine Functional Status Scale; and "job limitation," a dichotomous composite outcome that included all workers who reported a limitation attributed to hand symptoms in one or more of the following areas: (1) limited ability to work, (2) decreased productivity, (3) lost time from work, (4) placed on job restrictions, and (5) changed job or employer.

Among the 888 workers followed, baseline mean age was 30.3 years (range, 18 to 66 years), and 63.9% were men. Carpal tunnel syndrome prevalence and incidence were similar to that reported in other worker populations.^{8–10} From Table 1, ulnar neuropathy at the wrist was associated with CTS in only 4 of the 21 CTS cases at follow-up. These workers had a slightly higher proportion of severe hand pain but similar limitations. More than half of the subjects with symptoms of CTS had symptoms in the fifth finger ($n = 45$ of 79), without differences on the other outcomes, and only one worker had confirmed CTS and ulnar syndrome.

This study was limited by the small number of cases and by the use of the NC-Stat automated nerve testing device,⁷ which limited our testing of ulnar neuropathy to the wrist. Nonetheless, our findings provide an interesting perspective. In clinical settings,

some case reports suggest that associated median and ulnar neuropathies affect prognosis after nerve release.^{3,11–13} Nevertheless, our results are consistent with a recent study of ulnar neuropathy at the elbow; while 60.5% of subjects had persistent symptoms at follow-up, there were no differences in disability and symptom severity seen among the 5% of patients with ulnar neuropathy who had comorbid CTS.¹⁴

Ulnar neuropathy is relatively understudied, and future research is needed from large prospective studies to determine the prognosis, management, and prevention of ulnar neuropathies from entrapment at the wrist or elbow.

Alexis Descatha, MD

Univ Versailles St-Quentin, Versailles France; Inserm, Centre for Research in Epidemiology and Population Health "Population-Based Epidemiological Cohorts" Research Platform, Villejuif France; and AP-HP, Occupational Health Unit/EMS (Samu92), University hospital of West suburb of Paris, Garches, France

Ann-Marie Dale, PhD

Division of General Medical Sciences Washington University School of Medicine, St Louis, Mo

Angelique Zeringue, MSc

Division of General Medical Sciences Washington University School of Medicine, St Louis, Mo

Bradley Evanoff, MD, MPH

Division of General Medical Sciences Washington University School of Medicine, St Louis, Mo

REFERENCES

1. Bland JDP. Carpal tunnel syndrome. *BMJ*. 2007;335:343–346.
2. Luckhaupt SE, Sweeney MH, Sestito JP, Calvert GM. What is the true prevalence of carpal tunnel syndrome among US workers [published online ahead of print October 11, 2013]? *Scand J Work Environ Health*. doi:10.5271/sjweh.3391.
3. Turner A, Kimble F, Gulyás K, Ball J. Can the outcome of open carpal tunnel release be predicted?: a review of the literature. *ANZ J Surg*. 2010;80:50–54.
4. Shi Q, MacDermid JC. Is surgical intervention more effective than non-surgical treatment for carpal tunnel syndrome? A systematic review. *J Orthop Surg*. 2011;6:17.
5. Descatha A, Dale AM, Franzblau A, Evanoff B. Natural history and predictors of long-term pain and function among workers with hand symptoms. *Arch Phys Med Rehabil*. 2013;94:1293–1299.

TABLE 1. Description of Outcomes at Follow-Up (Severe Hand Pain, Functional Status Limitation on Levine Scale, and Job Limitation) Among Subjects Categorized at Baseline on Clinical Status and Nerve Conduction Status

Outcomes (at Follow-Up)	Variables (Baseline)	Total (N)	Cases (n)	n/N, %	Crude Odds Ratios (95% Confidence Interval)
Severe hand pain	No hand pain	732	111	15.16	1.00 (—)
	Other hand symptoms	82	22	26.83	2.05 (1.21–3.48)
	Symptoms of CTS, no nerve conduction abnormalities	53	16	30.19	2.42 (1.30–4.50)
	CTS confirmed without ulnar neuropathy	17	10	58.82	7.99 (2.98–21.44)
	CTS confirmed with ulnar neuropathy	4	3	75	16.76 (1.73–162.49)
Functional status limitation on Levine scale	No hand pain	732	59	8.06	1.00 (—)
	Other hand symptoms	82	6	7.32	0.90 (0.38–2.16)
	Symptoms of CTS, no nerve conduction abnormalities	53	7	13.21	1.74 (0.75–4.01)
	CTS confirmed without ulnar neuropathy	17	4	23.53	3.51 (1.11–11.10)
	CTS confirmed with ulnar neuropathy	4	1	25	3.80 (0.39–37.13)
Job limitation	No hand pain	732	72	9.84	1.00 (—)
	Other hand symptoms	82	19	23.17	2.76 (1.57–4.88)
	Symptoms of CTS, no nerve conduction abnormalities	53	9	16.98	1.88 (0.88–4.00)
	CTS confirmed without ulnar neuropathy	17	5	29.41	3.82 (1.31–11.15)
	CTS confirmed with ulnar neuropathy	4	1	25	3.06 (0.31–29.76)

CTS, carpal tunnel syndrome.

- Armstrong TN, Dale AM, Franzblau A, Evanoff BA. Risk factors for carpal tunnel syndrome and median neuropathy in a working population. *J Occup Environ Med.* 2008;50:1355–1364.
- Armstrong TN, Dale AM, Al-Lozi MT, Franzblau A, Evanoff BA. Median and ulnar nerve conduction studies at the wrist: criterion validity of the NC-stat automated device. *J Occup Environ Med.* 2008;50:758–764.
- Atroshi I, Gummesson C, Johnsson R, Ornstein E, Ranstam J, Rosén I. Prevalence of carpal tunnel syndrome in a general population. *JAMA.* 1999;282:153–158.
- Dale AM, Harris-Adamson C, Rempel D, et al. Prevalence and incidence of carpal tunnel syndrome in US working populations: pooled analysis of six prospective studies. *Scand J Work Environ Health.* 2013;39:495–505.
- Roquelaure Y, Ha C, Pelier-Cady MC, et al. Work increases the incidence of carpal tunnel syndrome in the general population. *Muscle Nerve.* 2008;37:477–482.
- Vimercati L, Lorusso A, L'abbate N, Assennato G. Bilateral carpal tunnel syndrome and ulnar neuropathy at the elbow in a pizza chef. *BMJ Case Rep.* 2009;2009. doi:10.1136/bcr.11.2008.1293:1–3.
- Desmoineaux P, Nezry N, Prové S, Boisrenoult P. Libération endoscopique du nerf ulnaire au coude technique—résultats. *Chir Main.* 2004;23(suppl 1):S128–S133.
- Ozdemir O, Calisaneller T, Gulsen S, Caner H. Ulnar nerve entrapment in Guyon's canal due to recurrent carpal tunnel syndrome: case report. *Turk Neurosurg.* 2011;21:435–437.
- Svensen SW, Johnsen B, Fuglsang-Frederiksen A, et al. Prognosis of ulnar neuropathy and ulnar neuropathy-like symptoms in relation to occupational biomechanical exposures and lifestyle. *Scand J Work Environ Health.* 2013;39:506–514.