

Washington University School of Medicine

Digital Commons@Becker

---

Open Access Publications

---

6-25-2020

## Exploiting the circuit breaker cancer evolution model in human clear cell renal cell carcinoma

James J Hsieh

*Washington University School of Medicine in St. Louis*

Emily H Cheng

*Weill Cornell Medical College*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

Please let us know how this document benefits you.

---

### Recommended Citation

Hsieh, James J and Cheng, Emily H, "Exploiting the circuit breaker cancer evolution model in human clear cell renal cell carcinoma." *Cell Stress*. 4, 8. 191 - 198. (2020).

[https://digitalcommons.wustl.edu/open\\_access\\_pubs/9617](https://digitalcommons.wustl.edu/open_access_pubs/9617)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

# Exploiting the circuit breaker cancer evolution model in human clear cell renal cell carcinoma

James J. Hsieh<sup>1,\*</sup> and Emily H. Cheng<sup>2,3,4</sup>

<sup>1</sup> Molecular Oncology, Department of Medicine, Washington University, St. Louis, MO 63110, USA.

<sup>2</sup> Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA.

<sup>3</sup> Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA.

<sup>4</sup> Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, Cornell University, New York, NY 10065, USA.

\* Corresponding Authors:

James J. Hsieh, M.D., Ph.D., 660 S. Euclid Avenue, Box 8069, St. Louis, MO 63110, USA; Tel: 314-273-1688; E-mail: Jhsieh@wustl.edu

**ABSTRACT** The incessant interactions between susceptible humans and their respective macro/microenvironments registered throughout their lifetime result in the ultimate manifestation of individual cancers. With the average lifespan exceeding 50 years of age in humans since the beginning of 20<sup>th</sup> century, aging – the “time” factor – has played an ever-increasing role alongside host and environmental factors in cancer incidences. Cancer is a genetic/epigenetic disease due to gain-of-function mutations in cancer-causing genes (oncogene; OG) and/or loss-of-function mutations in tumor-suppressing genes (tumor suppressor genes; TSG). In addition to their integral relationship with cancer, a timely deployment of specific OG and/or TSG is in fact needed for higher organisms like human to cope with respective physiological and pathological conditions. Over the past decade, extensive human kidney cancer genomics have been performed and novel mouse models recapitulating human kidney cancer pathobiology have been generated. With new genomic, genetic, mechanistic, clinical and therapeutic insights accumulated from studying clear cell renal cell carcinoma (ccRCC) – the most common type of kidney cancer, we conceived a cancer evolution model built upon the OG-TSG signaling pair analogous to the electrical circuit breaker (CB) that permits necessary signaling output and at the same time prevent detrimental signaling overdrive. Hence, this viewpoint aims at providing a step-by-step mechanistic explanation/illustration concerning how inherent OG-TSG CBs intricately operate in concert for the organism’s wellbeing; and how somatic mutations, the essential component for genetic adaptability, inadvertently triggers a sequential outage of specific sets of CBs that normally function to maintain and protect and individual tissue homeostasis.

doi: 10.15698/cst2020.08.227

Received originally: 19.04.2020;

in revised form: 16.06.2020

Accepted 17.06.2020.

Published 25.06.2020.

**Keywords:** kidney cancer evolution, circuit breaker, VHL, HIF, PBRM1, MTOR.

## Abbreviations:

CB – circuit breaker;  
ccRCC – clear cell RCC;  
HIF – hypoxia inducible factor;  
OG – oncogene;  
mccRCC – metastatic ccRCC;  
RCC – renal cell carcinoma;  
TSG – tumor suppressor gene;  
VHL – Von-Hippel Lindau.

## INTRODUCTION

On average a human adult body encompasses fifty trillion cells ( $\sim 5 \times 10^{13}$ ) with a daily turnover of hundred billion cells ( $\sim 10^{11}$ ) within which each contains two copies of  $\sim$ three billion base-pair DNA haploid genome. Approximately, the routine maintenance of a healthy human body necessitates  $\sim 10^{21}$  DNA base pairs transaction every day. Remarkably, despite this astronomical chemical challenge genome integrity is inherently maintained by sophisticated DNA replication, proof-reading, and repair mechanisms evolved over billion years. However, the precision of DNA transaction in

multi-cellular complex organisms can only be near-perfect to allow finite replication errors needed for genetic adaptation and thereby evolution. Nevertheless, individual cells are equipped with intrinsic cell death machineries to assure continuous genome integrity, which works alongside with immune system’s surveillance, killing, and removal of detected pre-cancer/cancer cells [1-3]. The human genome encodes  $\sim 20,000$  protein coding sequences of which 2-10% are cancer-related genes such as oncogenes (OGs) and tumor suppressor genes (TSGs) [2, 4]. Notably, these cancer-related genes normally participate in key biological

processes such as embryonic development and tissue homeostasis such as MLL (a Trx-G gene) and BMI-1 (a Pc-G gene)[5]. In response to imminent or chronic tissue stresses such as ischemia tissue injury these “specialty genes” are called upon to act appropriately till resolution of respective insults [2, 5-6]. Hence, we envisioned that specific oncogene and tumor suppressor gene pairs (OG-TSG) could function as regulatory modules similar to electrical circuit breakers (CBs) that mitigate power/signal overload [5]. Additionally, these OG-TSG CBs can interconnect to provide additional layers of regulation and safety that operate in a tissue context-dependent manner, which helps explain the tissue-type specific propensity of dysfunction in certain OG and TSG [7]. With these basic principles, hereafter we will begin to explore this hypothetical OG-TSG CB cancer evolution model inspired when we studied clear cell renal cell carcinoma (ccRCC) in human, mouse, and cell line models.

### Kidney cancer

Kidney cancer accounts for ~2% of all cancer diagnoses (~74,000; ~300,000) and deaths (~15,000; ~134,000) annually at the United States and the World, respectively[8, 9]. Renal cell carcinoma (RCC) encompasses a large heterogeneous group of cancers derived from renal tubular epithelial cells, which encompasses >10 molecular and histopathological subtypes [10]. RCC major subtypes (≥5%) include clear cell RCC (ccRCC) at ~75%, papillary RCC (pRCC) at ~15%, and chromophobe RCC (chRCC) at ~5% [11]. Accordingly, metastatic ccRCC (mccRCC) accounts for most of the kidney cancer mortality [12]. Although mccRCC is refractory to conventional chemotherapy, marked therapeutic advances have been made over the past 15 years, culminating in 15 FDA-approved agents for mccRCC [9]. These agents exploit seven molecular mechanisms: (1) IL-2 and α-Interferon are cytokines; (2) Sorafenib, Sunitinib, Pazopanib, Axitinib and Bevacizumab inhibit VEGF pathway; (3) Cabozantinib inhibits VEGFR2, cMET and AXL; (4) Lenvatinib inhibits VEGFR2 and FGFR; (5) Everolimus and Temsirolimus are inhibitors of mTORC1; (6) Nivolumab, Avelumab, and Pembrolizumab are PD-1/L1 blocking antibodies; and (7) Ipilimumab is an anti-CTLA-4 antibody [13]. However, mccRCC remains lethal and treatment response is highly heterogeneous among patients upon individual treatments [14], likely due to tumor genomic and host genetic heterogeneities.

### The genomics of ccRCC

Modern multi-omics of human ccRCC consisting of genomics, transcriptomics, proteomics and metabolomics not only highlights its profound inter-/intra-tumor heterogeneity but also showcases underline cancer evolution constraints that could be exploited as therapeutic biomarkers [14-25]. The most conspicuous genomic event in ccRCC is the near universal (>90%) one copy loss of the short arm of Chromosome 3 [26]. The most striking genomic discovery in ccRCC is the extreme chromosomal proximity of the four most prevalently mutated kidney cancer TSGs—*VHL*

(~80%), *PBRM1* (29-46%), *SETD2* (8-30%), and *BAP1* (6-19%)—spanning chromosome 3p21-3p25 [6, 27-28]. Hence, one genetic event incurred the 3p loss in renal epithelial cells simultaneously creates a haplo-insufficient state of four critical ccRCC TSGs. Remarkably, 3p loss represents the first somatic driver event in sporadic ccRCC, which takes place during adolescence that predates the most common second somatic event by ~20 years, i.e., the genetic/epigenetic inactivation of the *VHL* gene, giving rise to the most recent common ancestor (MRCA) within a given ccRCC patient whose ccRCC is eventually diagnosed at 60 years of age [29]. Notably, unlike most cancers that are initiated by gain-of-function mutations in OGs ccRCC results from sequential losses of TSGs [6].

### The shared genetic events of hereditary and sporadic in human ccRCC

The Von-Hippel Lindau (VHL) disease, an autosomal dominant hereditary cancer syndrome caused by the loss-of-function germline mutation in the *VHL* gene [30], is characterized by the development of hemangioblastoma of the central nervous system and retina, ccRCC, and pheochromocytoma [31-32]. It was later demonstrated that *VHL* is inactivated in ~90% of sporadic ccRCC through either genetic mutation or promoter methylation [9, 33]. Biallelic inactivation of the TSG *VHL* is therefore established as an early event in both germline mutant VHL-associated and sporadic ccRCC [6].

## DISCUSSION

### The VHL-HIF-hypoxia-metabolism

Studies on oxygen sensing led to the discovery of Hypoxia-Inducible Factors (HIFs) [34]. VHL is a multipurpose adaptor protein and chiefly effects as the substrate recognition module of the VCB (*VHL*-Elongin *C*-Elongin *B*)-Cul2 E3 ligase which ubiquitinates HIF-1α and HIF-2α [35]. Under normal oxygen conditions, HIF1/2α is prolyl hydroxylated by EGLN, ubiquitinated by VCB-Cul2-VHL, and rapidly degraded by the 26S Proteasome [36]; whereas under low oxygen conditions such as high altitude or ischemia, HIFα is stabilized to initiate a myriad of hypoxia-specific transcriptional programs [34, 36-38]. The pathologic loss of VHL in ccRCC results in persistently elevated HIFs accounting for the observed clear cell morphology and highly vascularity [11, 12, 38-40]. However, the long latency (>30 years) for VHL syndrome patients to develop ccRCC [31] and the insufficiency of VHL loss alone to induce ccRCC in mice [41] argue for the necessity of cooperative events [42].

### ccRCC signifies prevalent loss-of-function mutations in TSGs at the renal epithelium

Unlike many cancers that originate from gain-of-function mutations in OGs such as *EGFR* and *RAS*, ccRCC manifests with prevalent loss-of-function mutations in TSGs, making the development of predictive biomarkers for individual targeted therapies and/or immunotherapies extremely challenging. Nevertheless, new therapeutic modalities, novel genetically engineered mouse models, clinically rele-

vant patient-derived cell line/xenograft models, and outcome-based biomarker studies altogether have begun to shed light on how these seemingly distinct research areas are in fact exquisitely interconnected [43]. Here, we wish to update essential findings in ccRCC, and present a novel concept of “Interconnected OG-TSG Circuit Breaker Cancer Evolution Model in ccRCC”.

### The first oncogenic driver event in ccRCC

In hereditary VHL-loss ccRCC, the first genetic event is the inheritance of a loss-of-function copy of the VHL gene; whereas in sporadic ccRCC, the first genetic event is the loss of chromosome 3p. The ensuing epi/genetic event in developing both kinds of ccRCC converges on the complete inactivation of VHL [6]. Hence, generally speaking ccRCC is a VHL-loss kidney cancer, and complete VHL inactivation is the quintessential first functional/genetic truncal event [7].

### The exemplary interconnected VHL/HIF/PBRM1/TSC/MTORC1 CBs in ccRCC

Among the three newly identified 3p21 TSGs in ccRCC, *PBRM1* is best studied in molecular mechanisms, mouse models, and human clinical outcomes [22, 28, 32, 42, 44–48]. Accordingly, we will further elaborate recent key research findings on *PBRM1* and attempt to reconcile how and why *PBRM1* loss in ccRCC impact the efficacy of select targeted therapies and immune checkpoint inhibitors.

#### *PBRM1*

The SWI/SNF are macromolecular protein complexes that utilize ATP to mobilize nucleosome, modulate chromatin structure, and thereby regulate central cellular, developmental and oncogenic processes [49–50]. They come with many flavors due to their interchangeable, dynamic compositions in nature [51]. Notably, mutations of individual SWI/SNF subunits are detected and exhibit preferential enrichment in ~20% human cancer of various types [52]. *PBRM1* is the defining component of the PBAF complex and is most commonly mutated in ccRCC [53]. Remarkably, the *in vivo* tumor suppressor role of *PBRM1* in ccRCC was confirmed and reported in 2017 by three independent laboratories using three different genetically engineered mouse models where combined losses of VHL and *PBRM1* lead to multifocal ccRCC in mouse kidney, whereas individual losses did not [42, 54–55]. How *PBRM1* loss might have contributed to the ccRCC pathogenesis will be discussed hereafter.

#### *The hypoxia scenario: the normal physiology*

Upon tissue injury, insufficient blood supply incurs, thereby resulting in low oxygen tension, which in turn stabilizes HIF-1 to act accordingly (Fig. 1A), i.e. inhibition of mitochondria oxidative phosphorylation and promotion of re-vascularization. Once the tissue repair is complete, normal oxygen tension is re-established, HIF-1 is prolyl hydroxylated which is recognized by VHL, ubiquitinated by VCB-Cullin 2 E3 ligase, and degraded by 26S proteasome, and tissue returns to a normal homeostatic state [34, 36].

#### *The VHL loss scenario: inactivation of one CB*


The complete pathologic loss of VHL due to chromosome 3p loss, mutations, and/or promoter methylation results in the aberrant abundance of HIF-1 protein, resulting in a hypoxia-like molecular response despite normal oxygen tension, i.e. pseudohypoxia. Under this scenario, the VHL loss inactivates the first ccRCC CB “VHL-HIF” (Fig. 1B) present in the renal cortex. *In vivo* mouse model studies demonstrated that the activation of pseudohypoxia program is evident when comparing the twelve week old normal appearing mouse renal cortex with kidney specific deletion of *Vhl* to that of wild-type control [42].

#### *The VHL/PBRM1 loss scenario: inactivation of two CBs*

The singular loss of VHL is insufficient in initiating ccRCC, which is recognized in both mouse and human VHL loss models [56]. The ensuing complete loss of *PBRM1* inactivates the second CB “HIF-PBRM1-STAT” in the renal cortex (Fig. 1C), leading to the increased transcription output of HIF and STAT targets [42]. Of note, the singular loss of *PBRM1* did not activate HIF or STAT targets [42]. It is known that HIF-1 and STAT3 cooperate to activate the expression of HIF-1 targets [57–58]. Consequently, the dysregulated interplay between HIF and STAT upon combined losses of VHL and *PBRM1* creates a feed-forward amplification loop that maximizes downstream gene expression [42]. The role of *PBRM1* in restricting HIF signaling output was independently reported using cell-based assays [59]. Under this scenario, the subsequent *PBRM1* loss inactivates the second ccRCC CB “HIF-PBRM1-STAT”.

#### *The VHL/PBRM1/TSC loss scenario: inactivation of three CBs*

Despite the fact that HIF/STAT signaling overdrive was evident in the twelve week-old *Vhl*<sup>-/-</sup>;*Pbrm1*<sup>-/-</sup> mouse kidney, the long latency (ten months) and the incomplete penetrance (~50%) to the ultimate development of multifocal ccRCC in this model suggested that additional OG-TSG CBs could be at play [42]. To this end, gene expression and immunohistochemical analyses comparing twelve month-old ccRCC tumors to twelve week-old renal cortices of *Vhl*<sup>-/-</sup>;*Pbrm1*<sup>-/-</sup> mice detected hyperactive mTORC1 signaling in tumors in addition to the demonstrated pre-existing activation of HIF/STAT and suppression of mitochondrial pathways [42]. Hence, the prevention of aberrant activation of mTORC1 pathway probably constitutes the third ccRCC CB. Of note, mTORC1 serves as the central nutrition state integrator of the cell and its main control is conferred by the TSG TSC1/2 complex [60]. Remarkably, loss-of-function mutations in TSC1 or TSC2 (~10%), and gain-of-function mutations in MTOR (~6%) are common in ccRCC and correlated with rapalog response in therapeutic outlier studies; and *Tsc1* and *Tsc2* expression levels are down-regulated in *Vhl*<sup>-/-</sup>;*Pbrm1*<sup>-/-</sup> mouse ccRCC tumors [17, 22, 42, 61–63]. Accordingly, the third CB in place to prevent ccRCC pathogenesis after losses of VHL and *PBRM1* is “TSC-mTORC1” (Fig. 1D). Of note, the inactivation of the “PTEN/PI3K” CB which functions upstream of the “TSC-mTORC1” CB is observed in 7% of ccRCC [22]. Mechanistic



**FIGURE 1: The interconnected OG-TSG CBs operate in renal epithelium to control signaling output and prevent tumorigenesis. (A)** Depicts the physiological employment of the inherent VHL-HIF CB to gauge tissue response to low oxygen tension, and **(B-D)** examine how the sequential losses of a pre-determined set of “OG-TSG CBs” once the first CB is inactivated result in ccRCC.

cally, the inactivation of the first “VHL-HIF” and the second “HIF-PBRM1-STAT” CBs and the resulting HIF output override apparently activates *Redd1*, a known HIF1 target and an activator of TSC2 [42, 64], which positions “TSC-mTORC1” as the preferred third CB after the losses of VHL and PBRM1 [56].

#### *A hypothetical immunosurveillance CB in ccRCC*

As STATs are key transcription factors in cancer inflammation and immunity [65], the activation of STAT pathway due to the combined loss of VHL and PBRM1 could render

the resulting tumors prone to immune regulation [42]. Recent approvals of single agent Nivolumab (anti-PD-1 antibody) as second line and the combination of Ipilimumab (anti-CTLA-4 antibody) and Nivolumab as first line treatment options for ccRCC have dramatically altered the therapeutic landscape of metastatic kidney cancer [66-67]. Intriguingly, a recent paper identified PBRM1 loss as a potential genomic biomarker for the treatment response to these immune checkpoint inhibitors [68-69] and others suggested otherwise [70-71], which needs further validation facing the daunting intratumor heterogeneity of ccRCC.

Nevertheless, these mechanistic, mouse, and human ccRCC studies support a working hypothesis in which the disarmed “STAT-PD-1” immunosurveillance CB can be reactivated through biological means such as anti-PD-1/L1 antibodies for therapeutic exploitation (Fig. 1D). This might be one of the important rationales of why immunotherapy has activity in ccRCC, a tumor generally associated with low tumor mutation burden and a lack of microsatellite instability.

### FUTURE DIRECTION


#### Integrated applied pathology to exploit functional pathology in advancing precision cancer therapeutics

The holy grail of contemporary cancer research is to be able to predict not only how an individual patient may benefit from currently available front-line therapies, but also how an individual tumor’s molecular identity could potentially inform resistance mechanisms and thereby help implement a novel, tailored combination therapeutic strategy to greatly improve clinical outcome. One of the most challenging issues concerning metastatic ccRCC care is the known conspicuous intra-tumor and inter-tumor heterogeneity, which probably contributes to clinical outcomes. Nevertheless, like all individual human subjects develop from the same two copies of genome, all cancer cells evolve from the same set of genetic materials carried in the host cell. Hence, we hypothesize that intrinsic programming principles are in place to guide embryonic development, maintain tissue homeostasis, and restrict tumorigenesis [7, 14]. To visualize this third-generation hypothesis, we propose a “OG-TSG CB Constrained Braided Cancer River” model by integrating our first-generation “Braided River Model” and second-generation “CB Model” to further expound on this carcinogenic principle inspired from studying ccRCC, which might be applicable to additional cancer types.

#### The OG-TSG CB constrained braided cancer river model

Despite conspicuous tumor heterogeneity, long-term clinical

benefits on single agent targeted therapy are not uncommonly observed with metastatic ccRCC patients, implicating underlying cancer evolutionary constraints that force nonrandom sequences of parallel gene/pathway/function/phenotype convergences (Fig. 2). Indeed, our multi-region genomics study performed on ccRCC therapeutic outliers treated with single agent mTORC1 inhibitors rendered invaluable insights concerning this hypothesis [62]. We first proposed a braided cancer river model to help depict individual cancer evolution and advise therapeutic options.[7] The “Braided River” model highlights parallel and convergent events occurring throughout individual ccRCC tumorigenesis. It illustrates the stepwise, ordered accumulation of different driver mutations in kidney cancer development to acquire cancer hallmarks. A late chaotic evolutionary time point was incorporated to explain the limited effectiveness of combined targeted therapies in highly aggressive cancers, when “speedy” driver mutations quickly accumulate to enable extreme subclonal evolution. With the new concept of interconnected OG-TSG CBs, we propose a “CB-Constrained Braided Cancer River” model in which inherent context-dependent OG-TSG CBs are positioned at the gene/pathway/function/phenotype convergent points (Fig. 2). In other words, CBs function like dams to limit flow and prevent flooding. This model stipulates that each individual cancer river initiates with a truncal driver event, and once that occurs the ensuing branching events could be predicted. Accordingly, restoring a pre-determined, preferred set of CBs at once could offer effective measures and guide trial designs.



**FIGURE 2.** The diagram depicts the “OG-TSG CB Constrained Braided Cancer River” model to explain the non-randomness of cancer evolution and devise personalized cancer treatment strategy.



## ACKNOWLEDGMENTS

JJH and EHC are supported by NIH R01 CA223231.

## CONFLICT OF INTEREST

JJH has received consulting fees from Eisai Inc., Japan; clinical trial funding from Merck & Co., Inc., USA, and Astra-Zeneca Pharmaceuticals, UK; and research funding from BostonGene, USA.

## REFERENCES

- Nowell PC (1976). The clonal evolution of tumor cell populations. *Science* 194(4260): 23-8. doi: 10.1126/science.959840
- Hanahan D, Weinberg RA (2011). Hallmarks of cancer: the next generation. *Cell* 144(5):646-74. doi: 10.1016/j.cell.2011.02.013
- Dunn GP, Old LJ, Schreiber RD (2004). The immunobiology of cancer immunosurveillance and immunoediting. *Immunity* 21(2):137-48. doi: 10.1016/j.immuni.2004.07.017
- Stratton MR, Campbell PJ, Futreal PA (2009). The cancer genome. *Nature* 458(7239):719-24. doi: 10.1038/nature07943
- Liu H, Takeda S, Cheng EH, Hsieh JJ (2008). Biphasic MLL takes helm at cell cycle control: implications in human mixed lineage leukemia. *Cell Cycle* 7(4):428-35. doi: 10.4161/cc.7.4.5426
- Hsieh JJ, Le VH, Oyama T, Ricketts CJ, Ho TH, Cheng EH (2018). Chromosome 3p Loss-Orchestrated VHL, HIF, and Epigenetic Deregulation in Clear Cell Renal Cell Carcinoma. *J Clin Oncol* 2018:JCO2018792549. doi: 10.1200/JCO.2018.79.2549
- Wei EY, Hsieh JJ (2015). A river model to map convergent cancer evolution and guide therapy in RCC. *Nat Rev Urol* 12(12):706-12. doi: 10.1038/nrurol.2015.260
- Siegel RL, Miller KD, Jemal A (2020). Cancer statistics, 2020. *CA Cancer J Clin* 70(1):7-30. doi: 10.3322/caac.21590
- Huang JJ, Hsieh JJ (2020). The Therapeutic Landscape of Renal Cell Carcinoma: From the Dark Age to the Golden Age. *Semin Nephrol* 40(1):28-41. doi: 10.1016/j.semnephrol.2019.12.004
- Moch H, Humphrey PA, Ulbright TM, Reuter VE (2016). Who classification of tumours of the urinary system and male genital organs. 4th Edition, Volume 8. **WHO/IARC Classification of Tumours, Lyon, France.**
- Hsieh JJ, Le V, Cao D, Cheng EH, Creighton CJ (2018). Genomic classifications of renal cell carcinoma: a critical step towards the future application of personalized kidney cancer care with pan-omics precision. *J Pathol* 244(5):525-37. doi: 10.1002/path.5022
- Hsieh JJ, Purdue MP, Signoretti S, Swanton C, Albiges L, Schmidinger M, Heng DY, Larkin J, Ficarra V (2017). Renal cell carcinoma. *Nat Rev Dis Primers* 3:17009. doi: 10.1038/nrdp.2017.9
- McKay RR, Bosse D, Choueiri TK (2018). Evolving Systemic Treatment Landscape for Patients With Advanced Renal Cell Carcinoma. *J Clin Oncol* 2018:JCO2018790253. doi: 10.1200/JCO.2018.79.0253
- Hsieh JJ, Manley BJ, Khan N, Gao J, Carlo MI, Cheng EH (2017). Overcome tumor heterogeneity-imposed therapeutic barriers through convergent genomic biomarker discovery: A braided cancer river model of kidney cancer. *Semin Cell Dev Biol* 64:98-106. doi: 10.1016/j.semcdb.2016.09.002

## COPYRIGHT

© 2020 Hsieh and Cheng. This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: James J. Hsieh and Emily H. Cheng (2020). Exploiting the circuit breaker cancer evolution model in human clear cell renal cell carcinoma. *Cell Stress* 4(8): 191-198. doi: 10.15698/cst2020.08.227

- Gerlinger M, Horswell S, Larkin J, Rowan AJ, Salm MP, Varela I, Fisher R, McGranahan N, Matthews N, Santos CR, Martinez P, Phillimore B, Begum S, Rabinowitz A, Spencer-Dene B, Gulati S, Bates PA, Stamp G, Pickering L, Gore M, Nicol DL, Hazell S, Futreal PA, Stewart A, Swanton C (2014). Genomic architecture and evolution of clear cell renal cell carcinomas defined by multiregion sequencing. *Nat Genet* 46(3):225-33. doi: 10.1038/ng.2891
- Gerlinger M, Rowan AJ, Horswell S, Larkin J, Endesfelder D, Gronroos E, Martinez P, Matthews N, Stewart A, Tarpey P, Varela I, Phillimore B, Begum S, McDonald NQ, Butler A, Jones D, Raine K, Latimer C, Santos CR, Nohadani M, Eklund AC, Spencer-Dene B, Clark G, Pickering L, Stamp G, Gore M, Szallasi Z, Downward J, Futreal PA, Swanton C (2012). Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med* 366(10):883-92. doi: 10.1056/NEJMoa1113205
- Hsieh JJ, Chen D, Wang PI, Marker M, Redzematovic A, Chen YB, Selcuklu SD, Weinhold N, Bouvier N, Huberman KH, Bhanot U, Chevinsky MS, Patel P, Pinciroli P, Won HH, You D, Viale A, Lee W, Hakimi AA, Berger MF, Socci ND, Cheng EH, Knox J, Voss MH, Voi M, Motzer RJ (2017). Genomic Biomarkers of a Randomized Trial Comparing First-line Everolimus and Sunitinib in Patients with Metastatic Renal Cell Carcinoma. *Eur Urol* 71(3):405-14. doi: 10.1016/j.eururo.2016.10.007
- Pena-Llopis S, Vega-Rubin-de-Celis S, Liao A, Leng N, Pavia-Jimenez A, Wang S, Yamasaki T, Zhrebker L, Sivanand S, Spence P, Kinch L, Hambuch T, Jain S, Lotan Y, Margulis V, Sagalowsky AI, Summerour PB, Kabbani W, Wong SW, Grishin N, Laurent M, Xie XJ, Haudenschild CD, Ross MT, Bentley DR, Kapur P, Brugarolas J (2012). BAP1 loss defines a new class of renal cell carcinoma. *Nat Genet* 44(7):751-9. doi: 10.1038/ng.2323
- Durinck S, Stawiski EW, Pavia-Jimenez A, Modrusan Z, Kapur P, Jaiswal BS, Zhang N, Toffessi-Tcheuyap V, Nguyen TT, Pahuja KB, Chen YJ, Saleem S, Chaudhuri S, Heldens S, Jackson M, Pena-Llopis S, Guillory J, Toy K, Ha C, Harris CJ, Holloman E, Hill HM, Stinson J, Rivers CS, Janakiraman V, Wang W, Kinch LN, Grishin NV, Haverty PM, Chow B, Gehring JS, Reeder J, Pau G, Wu TD, Margulis V, Lotan Y, Sagalowsky A, Pedrosa I, de Sauvage FJ, Brugarolas J, Seshagiri S (2015). Spectrum of diverse genomic alterations define non-clear cell renal carcinoma subtypes. *Nat Genet* 47(1):13-21. doi: 10.1038/ng.3146
- Sato Y, Yoshizato T, Shiraishi Y, Maekawa S, Okuno Y, Kamura T, Shimamura T, Sato-Otsubo A, Nagae G, Suzuki H, Nagata Y, Yoshida K, Kon A, Suzuki Y, Chiba K, Tanaka H, Niida A, Fujimoto A, Tsunoda T, Morikawa T, Maeda D, Kume H, Sugano S, Fukayama M, Aburatani H, Sanada M, Miyano S, Homma Y, Ogawa S (2013). Integrated molecular analysis of clear-cell renal cell carcinoma. *Nat Genet* 45(8):860-7. doi: 10.1038/ng.2699
- Casuscelli J, Weinhold N, Gundem G, Wang L, Zabor EC, Drill E, Wang PI, Nanjangud GJ, Redzematovic A, Nargund AM, Manley BJ,

- Arcila ME, Donin NM, Cheville JC, Thompson RH, Pantuck AJ, Russo P, Cheng EH, Lee W, Tickoo SK, Ostrovnaya I, Creighton CJ, Papaemmanuil E, Seshan VE, Hakimi AA, Hsieh JJ (2017). Genomic landscape and evolution of metastatic chromophobe renal cell carcinoma. *JCI insight* 2(12): e92688. doi: 10.1172/jci.insight.92688
22. Cancer Genome Atlas Research Network (2013). Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature* 499(7456):43-9. doi: 10.1038/nature12222
23. Cancer Genome Atlas Research Network, *et al.* (2016). Comprehensive Molecular Characterization of Papillary Renal-Cell Carcinoma. *N Engl J Med* 374(2):135-45. doi: 10.1056/NEJMoa1505917
24. Davis CF, Ricketts CJ, Wang M, Yang L, Cherniack AD, Shen H, Buhay C, Kang H, Kim SC, Fahey CC, Hacker KE, Bhanot G, Gordenin DA, Chu A, Gunaratne PH, Biehl M, Seth S, Kaiparettu BA, Bristow CA, Donehower LA, Wallen EM, Smith AB, Tickoo SK, Tamboli P, Reuter V, Schmidt LS, Hsieh JJ, Choueiri TK, Hakimi AA, Cancer Genome Atlas Research N, Chin L, Meyerson M, Kucherlapati R, Park WY, Robertson AG, Laird PW, Henske EP, Kwiatkowski DJ, Park PJ, Morgan M, Shuch B, Muzny D, Wheeler DA, Linehan WM, Gibbs RA, Rathmell WK, Creighton CJ (2014). The somatic genomic landscape of chromophobe renal cell carcinoma. *Cancer Cell* 26(3):319-30. doi: 10.1016/j.ccr.2014.07.014
25. Sankin A, Hakimi AA, Mikkilineni N, Ostrovnaya I, Silk MT, Liang YP, Mano R, Chevinsky M, Motzer RJ, Solomon SB, Cheng EH, Durack JC, Coleman JA, Russo P, Hsieh JJ (2014). The impact of genetic heterogeneity on biomarker development in kidney cancer assessed by multiregional sampling. *Cancer Med-Us* 3(6):1485-92. doi: 10.1002/cam4.293
26. Zbar B, Brauch H, Talmadge C, Linehan M (1987). Loss of alleles of loci on the short arm of chromosome 3 in renal cell carcinoma. *Nature* 327(6124):721-4. doi: 10.1038/327721a0
27. Hakimi AA, Pham CG, Hsieh JJ (2013). A clear picture of renal cell carcinoma. *Nat Genet* 45(8):849-50. doi: 10.1038/ng.2708
28. Casuscelli J, Vano Y-A, Fridman WH, Hsieh JJ (2017). Molecular Classification of Renal Cell Carcinoma and Its Implication in Future Clinical Practice. *Kidney Cancer* 1(1):3-13. doi: 10.3233/KCA-170008
29. Mitchell TJ, *et al.* (2018). Timing the Landmark Events in the Evolution of Clear Cell Renal Cell Cancer: TRACERx Renal. *Cell* 173(3):611-23 e17. doi: 10.1016/j.cell.2018.02.020
30. Linehan WM, Lerman MI, Zbar B (1995). Identification of the von Hippel-Lindau (VHL) gene. Its role in renal cancer. *JAMA* 273(7):564-70. doi: 10.1001/jama.1995.03520310062031
31. Lonser RR, Glenn GM, Walther M, Chew EY, Libutti SK, Linehan WM, Oldfield EH (2003). von Hippel-Lindau disease. *Lancet*. 361(9374):2059-67. doi: 10.1016/S0140-6736(03)13643-4
32. Fisher R, Horswell S, Rowan A, Salm MP, de Bruin EC, Gulati S, McGranahan N, Stares M, Gerlinger M, Varela I, Crockford A, Favero F, Quidville V, Andre F, Navas C, Gronroos E, Nicol D, Hazell S, Hrouda D, O'Brien T, Matthews N, Phillimore B, Begum S, Rabinowitz A, Biggs J, Bates PA, McDonald NQ, Stamp G, Spencer-Dene B, Hsieh JJ, Xu J, Pickering L, Gore M, Larkin J, Swanton C (2014). Development of synchronous VHL syndrome tumors reveals contingencies and constraints to tumor evolution. *Genome Biol* 15(8):433. doi: 10.1186/s13059-014-0433-z
33. Linehan WM, Srinivasan R, Schmidt LS (2010). The genetic basis of kidney cancer: a metabolic disease. *Nat Rev Urol* 7(5):277-85. doi: 10.1038/nrur.2010.47
34. Semenza GL (2013). HIF-1 mediates metabolic responses to intratumoral hypoxia and oncogenic mutations. *J Clin Invest* 123(9):3664-71. doi: 10.1172/JCI67230
35. Frew IJ, Krek W (2008). pVHL: a multipurpose adaptor protein. *Sci Signal* 1(24):pe30. doi: 10.1126/scisignal.124pe30
36. Ivan M, Kaelin WG Jr (2017). The EGLN-HIF O<sub>2</sub>-Sensing System: Multiple Inputs and Feedbacks. *Mol Cell* 66(6):772-9. doi: 10.1016/j.molcel.2017.06.002
37. Masson N, Ratcliffe PJ (2014). Hypoxia signaling pathways in cancer metabolism: the importance of co-selecting interconnected physiological pathways. *Cancer Metab* 2(1):3. doi: 10.1186/2049-3002-2-3
38. Nakazawa MS, Keith B, Simon MC (2016). Oxygen availability and metabolic adaptations. *Nat Rev Cancer* 16(10):663-73. doi: 10.1038/nrc.2016.84
39. Schito L, Semenza GL (2016). Hypoxia-Inducible Factors: Master Regulators of Cancer Progression. *Trends Cancer* 2(12):758-70. doi: 10.1016/j.trecan.2016.10.016
40. Hakimi AA, Reznik E, Lee CH, Creighton CJ, Brannon AR, Luna A, Aksoy BA, Liu EM, Shen RL, Lee W, Chen Y, Stirdivant SM, Russo P, Chen YB, Tickoo SK, Reuter VE, Cheng EH, Sander C, Hsieh JJ (2019). An Integrated Metabolic Atlas of Clear Cell Renal Cell Carcinoma. *Cancer Cell* 29(1):104-16. doi: 10.1016/j.ccell.2015.12.004
41. Kapitsinou PP, Haase VH (2008). The VHL tumor suppressor and HIF: insights from genetic studies in mice. *Cell Death Differ* 15(4):650-9. doi: 10.1038/sj.cdd.4402313
42. Nargund AM, Pham CG, Dong Y, Wang PI, Osmangoyoglu HU, Xie Y, Aras O, Han S, Oyama T, Takeda S, Ray CE, Dong Z, Berge M, Hakimi AA, Monette S, Lekaye CL, Koutcher JA, Leslie CS, Creighton CJ, Weinhold N, Lee W, Tickoo SK, Wang Z, Cheng EH, Hsieh JJ (2017). The SWI/SNF Protein PBRM1 Restrains VHL-Loss-Driven Clear Cell Renal Cell Carcinoma. *Cell Rep* 18(12):2893-906. doi: 10.1016/j.celrep.2017.02.074
43. Wolf MM, Kimryn Rathmell W, Beckermann KE (2020). Modeling clear cell renal cell carcinoma and therapeutic implications. *Oncogene* 39(17):3413-3426. doi: 10.1038/s41388-020-1234-3
44. Lee CH, Pham CG, Hsieh JJ (2015). PBRM1: A Critical Subunit of the SWI/SNF Chromatin Remodeling Complex. Renal Cell Carcinoma. In: Bukowski RM, Motzer RJ, Figlin RA, editors. *Molecular Targets and Clinical Applications*. Springer, New York; pp 111-51. doi: 10.1007/978-1-4939-1622-1\_5
45. Becerra MF, Manley BJ, Casuscelli J, Redzematovic A, Tennenbaum DM, Motzer RJ, Feldman DR, Voss MH, Arcila M, Russo P, Coleman JA, Hsieh JJ, Hakimi AA (2016). Genomic Landscape of Clear Cell Renal Cell Carcinoma Metastasis. *J Urology* 195(4):E919-E20.
46. Chen FJ, *et al.* (2016). Multilevel Genomics-Based Taxonomy of Renal Cell Carcinoma. *Cell Rep* 14(10):2476-89. doi: 10.1016/j.celrep.2016.02.024
47. Hakimi AA, Ostrovnaya I, Reva B, Schultz N, Chen YB, Gonen M, Liu H, Takeda S, Voss MH, Tickoo SK, Reuter VE, Russo P, Cheng EH, Sander C, Motzer RJ, Hsieh JJ (2013). Adverse Outcomes in Clear Cell Renal Cell Carcinoma with Mutations of 3p21 Epigenetic Regulators BAP1 and SETD2: A Report by MSKCC and the KIRC TCGA Research Network. *Clin Cancer Res* 19(12):3259-67. doi: 10.1158/1078-0432.CCR-12-3886
48. Hakimi AA, Chen YB, Wren J, Gonen M, Abdel-Wahab O, Heguy A, Liu H, Takeda S, Tickoo SK, Reuter VE, Voss MH, Motzer RJ, Coleman JA, Cheng EH, Russo P, Hsieh JJ (2013). Clinical and pathologic impact of select chromatin-modulating tumor suppressors in clear cell renal cell carcinoma. *Eur Urol* 63(5):848-54. doi: 10.1016/j.eururo.2012.09.005
49. Clapier CR, Cairns BR (2009). The biology of chromatin remodeling complexes. *Annu Rev Biochem* 78:273-304. doi: 10.1146/annurev.biochem.77.062706.153223



50. Biegel JA, Busse TM, Weissman BE (2014). SWI/SNF chromatin remodeling complexes and cancer. **Am J Med Genet C Semin Med Genet** 166C(3):350-66. doi: 10.1002/ajmg.c.31410
51. Hodges C, Kirkland JG, Crabtree GR (2016). The Many Roles of BAF (mSWI/SNF) and PBAF Complexes in Cancer. **Cold Spring Harb Perspect Med** 6(8): a026930. doi: 10.1101/cshperspect.a026930
52. Kadoch C, Hargreaves DC, Hodges C, Elias L, Ho L, Ranish J, Crabtree GR (2013). Proteomic and bioinformatic analysis of mammalian SWI/SNF complexes identifies extensive roles in human malignancy. **Nat Genet** 45(6):592-601. doi: 10.1038/ng.2628
53. Varela I, *et al.* (2011). Exome sequencing identifies frequent mutation of the SWI/SNF complex gene PBRM1 in renal carcinoma. **Nature** 469(7331):539-42. doi: 10.1038/nature09639
54. Gu YF, Cohn S, Christie A, McKenzie T, Wolff NC, Do QN, Madhuranthakam A, Pedrosa I, Wang T, Dey A, Busslinger M, Xie XJ, Hammer RE, McKay RM, Kapur P, Brugarolas J (2017). Modeling Renal Cell Carcinoma in Mice: Bap1 and Pbrm1 Inactivation Drive Tumor Grade. **Cancer Discov** 7(8):900-917. doi: 10.1158/2159-8290.CD-17-0292
55. Espana-Agusti J, Warren A, Chew SK, Adams DJ, Matakidou A (2017). Loss of PBRM1 rescues VHL dependent replication stress to promote renal carcinogenesis. **Nat Commun** 8(1):2026. doi: 10.1038/s41467-017-02245-1
56. Nargund AM, Osmanbeyoglu HU, Cheng EH, Hsieh JJ (2017). SWI/SNF tumor suppressor gene PBRM1/BAF180 in human clear cell kidney cancer. **Mol Cell Oncol** 4(4):e1342747. doi: 10.1080/23723556.2017.1342747
57. Luo W, Semenza GL (2012). Emerging roles of PKM2 in cell metabolism and cancer progression. **Trends Endocrinol Metab** 23(11):560-6. doi: 10.1016/j.tem.2012.06.010
58. Jung JE, Lee HG, Cho IH, Chung DH, Yoon SH, Yang YM, Lee JW, Choi S, Park JW, Ye SK, Chung MH (2005). STAT3 is a potential modulator of HIF-1-mediated VEGF expression in human renal carcinoma cells. **FASEB J** 19(10):1296-8. doi: 10.1096/fj.04-3099fje
59. Gao W, Li W, Xiao T, Liu XS, Kaelin WG Jr (2017). Inactivation of the PBRM1 tumor suppressor gene amplifies the HIF-response in VHL/-clear cell renal carcinoma. **Proc Natl Acad Sci U S A** 114(5):1027-32. doi: 10.1073/pnas.1619726114
60. Saxton RA, Sabatini DM (2017). mTOR Signaling in Growth, Metabolism, and Disease. **Cell** 169(2):361-71. doi: 10.1016/j.cell.2017.03.035
61. Xu J, Pham CG, Albanese SK, Dong Y, Oyama T, Lee CH, Rodrik-Outmezguine V, Yao Z, Han S, Chen D, Parton DL, Chodera JD, Rosen N, Cheng EH, Hsieh JJ (2016). Mechanistically distinct cancer-associated mTOR activation clusters predict sensitivity to rapamycin. **J Clin Invest** 126(9):3526-40. doi: 10.1172/JCI86120
62. Voss MH, Hakimi AA, Pham CG, Brannon AR, Chen YB, Cunha LF, Akin O, Liu H, Takeda S, Scott SN, Succi ND, Viale A, Schultz N, Sander C, Reuter VE, Russo P, Cheng EH, Motzer RJ, Berger MF, Hsieh JJ (2014). Tumor genetic analyses of patients with metastatic renal cell carcinoma and extended benefit from mTOR inhibitor therapy. **Clin Cancer Res** 20(7):1955-64. doi: 10.1158/1078-0432.CCR-13-2345
63. Kwiatkowski DJ, Choueiri TK, Fay AP, Rini BI, Thorner AR, De Velasco G, Tyburczy M, Hamieh L, Albiges L, Agarwal N, Ho TH, Song J, Pignon JC, Barrios PM, Michaelson MD, Van Allen EM, Krajewski KM, Porta C, Pal SK, Bellmunt J, McDermott DF, Heng DY, Gray KP, Signoretti S (2016). Mutations in TSC1, TSC2, and MTOR are associated with response to rapalogs in patients with metastatic Renal Cell Carcinoma. **Clin Cancer Res** 22(10):2445-52. doi: 10.1158/1078-0432.CCR-15-2631
64. Brugarolas J, Lei K, Hurley RL, Manning BD, Reiling JH, Hafen E, Witters LA, Ellisen LW, Kaelin WG Jr (2004). Regulation of mTOR function in response to hypoxia by REDD1 and the TSC1/TSC2 tumor suppressor complex. **Genes Dev** 18(23):2893-904. doi: 10.1101/gad.1256804
65. Yu H, Pardoll D, Jove R (2009). STATs in cancer inflammation and immunity: a leading role for STAT3. **Nat Rev Cancer** 9(11):798-809. doi: 10.1038/nrc2734
66. Motzer RJ, Escudier B, McDermott DF, George S, Hammers HJ, Srinivas S, Tykodi SS, Sosman JA, Procopio G, Plimack ER, Castellano D, Choueiri TK, Gurney H, Donskov F, Bono P, Wagstaff J, Gauder TC, Ueda T, Tomita Y, Schutz FA, Kollmannsberger C, Larkin J, Ravaud A, Simon JS, Xu LA, Waxman IM, Sharma P, CheckMate I (2015). Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. **N Engl J Med** 373(19):1803-13. doi: 10.1056/NEJMoa1510665
67. Motzer RJ, Tannir NM, McDermott DF, Aren Frontera O, Melichar B, Choueiri TK, Plimack ER, Barthelemy P, Porta C, George S, Powles T, Donskov F, Neiman V, Kollmannsberger CK, Salman P, Gurney H, Hawkins R, Ravaud A, Grimm MO, Bracarda S, Barrios CH, Tomita Y, Castellano D, Rini BI, Chen AC, Mekan S, McHenry MB, Wind-Rotolo M, Doan J, Sharma P, Hammers HJ, Escudier B, CheckMate I (2018). Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. **N Engl J Med** 378(14):1277-1290. doi: 10.1056/NEJMoa1712126
68. Miao D, Margolis CA, Gao W, Voss MH, Li W, Martini DJ, Norton C, Bosse D, Wankowicz SM, Cullen D, Horak C, Wind-Rotolo M, Tracy A, Giannakis M, Hodi FS, Drake CG, Ball MW, Allaf ME, Snyder A, Hellmann MD, Ho T, Motzer RJ, Signoretti S, Kaelin WG, Jr., Choueiri TK, Van Allen EM (2018). Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma. **Science** 359(6377):801-806. doi: 10.1126/science.aan5951
69. Braun DA, Ishii Y, Walsh AM, Van Allen EM, Wu CJ, Shukla SA, Choueiri TK (2019). Clinical Validation of PBRM1 Alterations as a Marker of Immune Checkpoint Inhibitor Response in Renal Cell Carcinoma. **JAMA Oncol** 5(11):1631-1633. doi: 10.1001/jamaoncol.2019.3158
70. Liu XD, Kong W, Peterson CB, McGrail DJ, Hoang A, Zhang X, Lam T, Pilie PG, Zhu H, Beckermann KE, Haake SM, Isgandrova S, Martinez-Moczygemba M, Sahni N, Tannir NM, Lin SY, Rathmell WK, Jonasch E (2020). PBRM1 loss defines a nonimmunogenic tumor phenotype associated with checkpoint inhibitor resistance in renal carcinoma. **Nat Commun** 11(1):2135. doi: 10.1038/s41467-020-15959-6
71. McDermott DF, Huseni MA, Atkins MB, Motzer RJ, Rini BI, Escudier B, Fong L, Joseph RW, Pal SK, Reeves JA, Sznol M, Hainsworth J, Rathmell WK, Stadler WM, Hutson T, Gore ME, Ravaud A, Bracarda S, Suarez C, Danielli R, Gruenwald V, Choueiri TK, Nickles D, Jhunjhunwala S, Piault-Louis E, Thobhani A, Qiu J, Chen DS, Hegde PS, Schiff C, Fine GD, Powles T (2018). Clinical activity and molecular correlates of response to atezolizumab alone or in combination with bevacizumab versus sunitinib in renal cell carcinoma. **Nat Med** 24(6):749-57. doi: 10.1038/s41591-018-0053-3