

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2020

Surgical treatment of femoroacetabular impingement: Hip arthroscopy versus surgical hip dislocation: A propensity-matched analysis

Jeffrey J Nepple

Washington University School of Medicine in St. Louis

Ira Zaltz

Beaumont Health System

Christopher M Larson

Twin Cities Orthopedics

Paul E Beaulé

The Ottawa Hospital

Young-Jo Kim

Harvard University

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Nepple, Jeffrey J; Zaltz, Ira; Larson, Christopher M; Beaulé, Paul E; Kim, Young-Jo; Millis, Michael B; Sierra, Rafael J; Clohisey, John C; and ANCHOR Group, "Surgical treatment of femoroacetabular impingement: Hip arthroscopy versus surgical hip dislocation: A propensity-matched analysis." *The Journal of Bone and Joint Surgery. American Volume*. 102, Suppl 2. 51 - 58. (2020).
https://digitalcommons.wustl.edu/open_access_pubs/9821

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Jeffrey J Nepple, Ira Zaltz, Christopher M Larson, Paul E Beaulé, Young-Jo Kim, Michael B Millis, Rafael J Sierra, John C Clohisy, and ANCHOR Group

Surgical Treatment of Femoroacetabular Impingement: Hip Arthroscopy Versus Surgical Hip Dislocation

A Propensity-Matched Analysis

Jeffrey J. Nepple, MD, Ira Zaltz, MD, Christopher M. Larson, MD, Paul E. Beaulé, MD, FRCSC, Young-Jo Kim, MD, Michael B. Millis, MD, Rafael J. Sierra, MD, the ANCHOR Group*, and John C. Clohisy, MD

Investigation performed at the Department of Orthopaedic Surgery, Washington University School of Medicine, St. Louis, Missouri

Background: Surgical treatment of femoroacetabular impingement (FAI) continues to evolve and is most commonly approached with either hip arthroscopy (HA) or surgical dislocation (SD) of the hip. The purpose of this study was to compare the outcomes of similar patients undergoing surgical treatment of FAI with either HA or SD.

Methods: A prospective multicenter cohort study of patients undergoing primary surgical treatment of FAI was performed. Follow-up at a minimum of 1 year (mean, 4.3 years) was available for 621 hips (81.7%), including 399 procedures with HA and 222 procedures with SD. Propensity scores were calculated and reflect the likelihood of surgical treatment with HA versus SD for a given set of covariates. Propensity scores allowed 1:1 matching to identify similar patients at baseline. After propensity matching, 128 matched pairs of patients who underwent HA and 128 matched pairs of those who underwent SD were included in the study. The primary outcome was the postoperative modified Harris hip score (mHHS); secondary outcomes included the Hip disability and Osteoarthritis Outcome Score (HOOS), the University of California Los Angeles (UCLA) activity score, and the Short Form-12 (SF-12) physical and mental subscores, as well as the rate of persistent symptoms, revision surgery, and total hip arthroplasty (THA).

Results: After propensity matching, the 2 groups exhibited similar distributions of all of the covariates that were included in the model. Both groups demonstrated significant improvements in all patient-reported outcomes (PROs). The final mHHS was not significantly different between the 2 matched groups (81.3 for the HA group versus 80.2 for the SD group, $p = 0.67$). Likewise, the HOOS pain subscale was similar at the time of final follow-up (77.6 versus 80.5, respectively, $p = 0.32$). No difference between the HA group and the SD group was identified in the rate of THA (0% and 3.1%, respectively, $p = 0.41$) and revision surgery (7.8% and 10.9%, respectively, $p = 0.35$); overall rates of persistent symptoms were 21.9% for the HA group and 24.4% for the SD group ($p = 0.55$).

Conclusions: In a propensity-matched analysis of patients who were treated with either approach, patients undergoing HA or SD demonstrated similar outcomes at a mean of 4 years postoperatively.

Level of Evidence: Therapeutic Level II. See Instructions for Authors for a complete description of levels of evidence.

Surgical treatment of femoroacetabular impingement (FAI) continues to evolve¹. Surgical treatment of FAI is most commonly approached with either hip arthroscopy (HA) or surgical dislocation (SD) of the hip. HA offers advantages

with its minimally invasive nature and quicker early rehabilitation², but inadequate deformity correction appears to be a common reason for surgical failure^{3,4}. On the other hand, SD offers global access to hip deformities that may be

*A list of the ANCHOR Group members is included as a note at the end of the article.

Disclosure: This study was supported in part by a clinical research grant from Zimmer-Biomet, Inc. to some of the authors (J.C.C., J.J.N.), The Curing Hip Disease Fund (J.C.C.), and the ANCHOR Fund (J.C.C.). On the **Disclosure of Potential Conflicts of Interest** forms, which are provided with the online version of the article, one or more of the authors checked "yes" to indicate that the author had a relevant financial relationship in the biomedical arena outside the submitted work (<http://links.lww.com/JBJS/G107>).

challenging or inaccessible by HA^{5,6}, but it requires a more invasive procedure, including trochanteric osteotomy. Decision-making between these 2 approaches has evolved substantially over the past decade¹. However, the effect of surgical approach on outcomes is not well established.

The literature is limited with regard to rigorous, direct comparisons of the results of these 2 techniques. Several studies have demonstrated a relatively similar ability to achieve typical cam correction with HA and SD^{6,7}, although SD may have some advantages in more severe proximal femoral and acetabular deformities^{5,6}. Several studies and systematic reviews have investigated the outcomes of HA and SD, with differing conclusions^{2,8-10}. Comparisons are challenging given the potentially different patient characteristics (age, sex, preoperative patient-reported outcomes [PROs], deformity severity), which often are not controlled for, of patients undergoing these 2 distinct procedures. Additionally, a randomized controlled trial on this topic may not be feasible given the potential for lack of surgeon equipoise.

The purpose of the current study was to compare the outcomes of surgical treatment of FAI with HA and SD utilizing a propensity analysis to identify hips with similar characteristics at baseline to allow a valid comparison¹¹.

Materials and Methods

This study utilized data from a prospective multicenter cohort study of patients undergoing primary surgical treatment of FAI. The entire cohort included 1,130 patients (1,076 hips); all patients had a primary clinical diagnosis of FAI and had failure of nonoperative treatment lasting at least 3 months¹². Inclusion criteria for the present study were primary surgery and a diagnosis of FAI. Exclusion criteria included previous surgery or residual pediatric hip disease (including residual slipped capital femoral epiphysis and deformities from Legg-Calvé-Perthes disease). A total of 760 hips undergoing surgical treatment of FAI met the inclusion criteria for this study. Eleven experienced surgeons at 9 sites contributed to the enrollment of the prospective cohort. Contributing surgeons had varying skill sets and surgical approach preferences, and included those performing only HA ($n = 1$), those performing only SD ($n = 4$), or surgeons performing both procedures ($n = 6$). SD was performed with trochanteric osteotomy, as described by Ganz et al.¹³. HA was performed with the patient in the supine position in all of the cases. The presence of cam and/or pincer FAI was determined by the clinical diagnosis by the treating surgeon. The maximum alpha angle was determined on anteroposterior, 45° Dunn, or frog-leg/cross-table lateral radiographs.




Fig. 1
Study flow diagram. The initial enrolled cohort included 760 hips; 621 hips with appropriate follow-up were eligible for inclusion. Propensity matching was performed to identify 256 hips (128 matched pairs) for inclusion in the study cohort.

TABLE I Baseline Characteristics of Matched and Unmatched Groups for Variables Utilized in Propensity Matching*

Variable	Propensity Matched				Unmatched			
	Surgical Dislocation (N = 128)	Hip Arthroscopy (N = 128)	P Value	Absolute Standardized Difference	Surgical Dislocation (N = 222)	Hip Arthroscopy (N = 399)	P Value	Absolute Standardized Difference
Female†	50.8% (65)	49.2% (63)	0.80	0.03	51.4% (114)	59.6% (238)	0.046	0.19
Involved in competitive sports†	22.7% (29)	22.7% (29)	1.00	0.00	20.7% (46)	11.8% (47)	0.003	0.23
Alpha angle >70°†	32.8% (42)	33.6% (43)	0.894	0.02	34.2% (76)	28.3% (113)	0.12	0.11
Lateral center-edge angle >40°†	7.8% (10)	7.0% (9)	0.812	0.03	14% (31)	6% (24)	0.001	0.34
FAI subtype†			0.876				<0.001	
Cam	35.9% (46)	37.5% (48)		0.03	23.9% (53)	57.4% (229)		0.66
Combined	56.3% (72)	56.3% (72)		0	64.0% (142)	38.3% (153)		0.46
Pincer	7.8% (10)	6.3% (8)		0.06	12.2% (27)	4.3% (17)		0.3
Tönnis osteoarthritis grade†			0.804				<0.001	
0	48.4% (62)	50.0% (64)		0.03	54.1% (120)	36.3% (145)		0.43
1	50.8% (65)	48.4% (62)		0.05	42.8% (95)	61.9% (247)		0.45
2	0.8% (1)	1.6% (2)		0.05	3.2% (7)	1.8% (7)		0.05
Age† (yr)	24.9 ± 9.0	24.9 ± 9.2	0.98	0.003	22.9 ± 8.3	33.7 ± 11.8	<0.001	1.01
Preop. mHHS†	61.4 ± 14.7	62.5 ± 15.9	0.54	0.08	62.4 ± 14.3	60.1 ± 15.6	0.08	0.16

*Significant p values are bolded. FAI = femoroacetabular impingement, and mHHS = modified Harris hip score. †The values for the 2 groups are given as the percentage, with the number in parentheses. ‡The values for the 2 groups are given as the mean and standard deviation.

In general, during the study period, pincer FAI was felt to be present in the setting of global acetabular overcoverage (lateral center-edge angle [LCEA] of >40°) or a positive crossover sign, while cam FAI was defined by an alpha angle of >50°. Surgical procedure type (HA or SD) was determined by the treating surgeon. Labral and cartilage pathology were classified using the modified Beck classification¹⁴⁻¹⁶. Follow-up at a minimum of 1 year was available with 621 hips (81.7%) (mean, 4.3 ± 2.4 years), including assessment of PROs (the modified Harris hip score [mHHS], the University of California Los Angeles (UCLA) activity score, 5 Hip disability and Osteoarthritis Outcome Score [HOOS] subscales, and the Short Form-12 [SF-12] physical and mental subscores) as well as reoperations (including total hip arthroplasty [THA] and revision surgery) (Fig. 1).

Propensity score matching is a powerful method to adjust for confounding variables and reduce, although not completely eliminate, selection bias in nonrandomized studies. Matching on propensity scores identifies similar patients undergoing HA and SD based on the set of included variables (e.g., preoperative

clinical characteristics) in order to compare outcomes between the surgical treatment types¹¹. Propensity scores were calculated (possible range, 0.00 to 1.00) using a logistic regression model with surgical treatment type (HA versus SD) as the outcome, and independent variables included factors that were felt to play a role in surgical approach decision-making based on clinical experience or univariate statistical analysis (Table I), as well as to play a role in the pathophysiology and outcome of treatment of FAI. Covariates that were included were sex, age, participation in competitive sports, preoperative mHHS, FAI subtype, moderate cam morphology (defined as a maximum alpha angle of >70°), global acetabular overcoverage (defined as an LCEA of >40°), and Tönnis osteoarthritis classification. HA and SD patients were randomly matched 1:1 based on propensity scores (within an absolute distance of 0.01) to allow for a comparison of similar patients who were commonly treated with either procedure. Standardized differences were used to assess the quality of the matching results, where an after-matching standardized difference of 0.1 implied that there was a

TABLE II Baseline Characteristics of Matched and Unmatched Groups for Variables Not Utilized in Propensity Matching*

Variable	Propensity Matched			Unmatched		
	Surgical Dislocation (N = 128)	Hip Arthroscopy (N = 128)	P Value	Surgical Dislocation (N = 222)	Hip Arthroscopy (N = 399)	P Value
Alpha angle	63.2 ± 12.5	63.1 ± 12.8	0.93	63.2 ± 13.3	62.5 ± 11.6	0.50
LCEA	30.2 ± 6.9	29.0 ± 6.3	0.13	31.9 ± 7.6	28.7 ± 6.6	<0.001
Acetabular inclination	4.1 ± 5.6	5.5 ± 5.7	0.047	3.8 ± 5.8	6.4 ± 6.9	<0.001
Preop. PROs						
HOOS domain						
Pain	55.1 ± 20.0	58.1 ± 21.0	0.25	57.0 ± 19.9	55.6 ± 21.1	0.42
ADL	65.9 ± 20.2	67.7 ± 21.3	0.49	66.8 ± 19.8	64.9 ± 22.2	0.28
QOL	29.2 ± 19.5	33.7 ± 20.8	0.08	30.3 ± 19.1	30.7 ± 20.0	0.83
Symptoms	53.8 ± 19.5	57.9 ± 21.8	0.12	55.5 ± 19.0	56.0 ± 21.4	0.75
Sport	46.5 ± 23.6	45.5 ± 22.8	0.73	47.3 ± 23.8	44.0 ± 25.0	0.11
UCLA activity score	7.2 ± 2.8	7.6 ± 2.7	0.22	7.1 ± 2.7	6.9 ± 2.8	<0.001
SF-12 Physical	38.4 ± 10.7	38.8 ± 10.8	0.79	38.6 ± 10.0	37.5 ± 10.7	0.22
SF-12 Mental	54.2 ± 10.2	52.7 ± 9.0	0.23	54.2 ± 10.1	51.8 ± 11.0	0.008

*LCEA = lateral center-edge angle, PROs = patient-reported outcomes, HOOS = Hip disability and Osteoarthritis Outcome Score, ADL = activities of daily living, QOL = quality of life, UCLA = University of California Los Angeles, and SF-12 = Short Form-12. The values are given as the mean. Significant p values are bolded.

negligible correlation and good balance between the group and each covariate¹⁷. The cohort initially included 621 patients (399 who underwent HA and 222 who underwent SD). After propensity matching, a total of 256 patients (128 matched pairs of patients undergoing HA and SD) were included in the current study and subsequent analysis (Fig. 1).

The primary outcome was the final mHHS, while secondary outcomes included the HOOS, the UCLA activity score, and the SF-12 score. Secondary outcomes also included the rates of revision surgery or THA and persistent symptoms. Persistent symptoms were defined as reoperation (for continued symptoms) or a symptomatic hip. A “symptomatic” hip

was defined as improvement in the mHHS of <8 (minimal clinically important difference [MCID])¹⁸ and an mHHS of <74 (patient acceptable symptom state [PASS])¹⁹; symptomatic hips reached neither the MCID nor the PASS. Reoperation was defined as revision surgery or THA and did not include hardware removal. Outcomes were compared in propensity-matched groups. Categorical variables were analyzed with the McNemar test (or Fisher exact test). Continuous variables were analyzed with paired t tests (a positive difference indicated a larger value for SD, and a negative value indicated a larger value for HA). Patients with a missing post-operative mHHS prior to undergoing a reoperation (revision

TABLE III Characteristics of HA and SD Patients Who Were Included and Excluded from Propensity-Matched Analysis*

	Hip Arthroscopy		P Value	Surgical Dislocation		P Value
	Included (N = 128)	Excluded (N = 271)		Included (N = 128)	Excluded (N = 94)	
Female†	49.2% (63)	64.6% (175)	0.004	50.8% (65)	52.1% (49)	0.84
Competitive sports†	22.7% (29)	6.6% (18)	<0.001	22.7% (29)	18.1% (17)	0.41
Alpha angle‡ (°)	63.1	62.2	0.52	63.2	63.2	1.00
Alpha angle >70°†	33.6% (43)	25.8% (70)	0.11	32.8% (42)	36.2% (34)	0.56
LCEA‡ (°)	28.9	28.5	0.55	30.2	34.2	<0.001
LCEA >40°†	7.0% (9)	5.5% (15)	0.56	7.8% (10)	22.3% (21)	0.002
Age‡ (yr)	24.8	37.8	<0.001	24.9	20.1	<0.001
Preoperative mHHS‡	62.5	59	0.034	61.4	63.8	0.23

*LCEA = lateral center-edge angle, and mHHS = modified Harris hip score. †The values for the 2 groups are given as the percentage, with the number in parentheses. ‡The values are given as the mean.

TABLE IV Intraoperative Disease Characteristics for the Hip Arthroscopy and Surgical Dislocation Groups*

	Hip Arthroscopy	Surgical Dislocation		Hip Arthroscopy	Surgical Dislocation	P Value
Modified Beck labral grade			Labral treatment			0.001
Normal	3.9% (5)	8.6% (11)	None	6.3% (8)	21.9% (28)	
Degeneration	12.5% (16)	25.0% (32)	Debridement	23.4% (30)	24.2% (31)	
Full-thickness tear	3.1% (4)	6.3% (8)	Repair	70.3% (90)	53.9% (69)	
Detachment	76.6% (98)	47.7% (61)				
Ossification	3.9% (5)	12.5% (16)				
Modified Beck acetabular cartilage grade			Acetabular cartilage treatment			<0.001
Normal	7.8% (10)	27.3% (35)	None	32.0% (41)	57.0% (73)	
Malacia	22.7% (29)	15.6% (20)	Chondroplasty	63.3% (81)	33.6% (43)	
Debonding	32.8% (42)	19.5% (25)	Microfracture	4.7% (6)	9.4% (12)	
Cleavage	29.7% (38)	27.3% (35)				
Defect	7.0% (9)	10.2% (13)				
Modified Beck femoral cartilage grade			Femoral cartilage treatment			0.046
Normal	89.1% (114)	71.9% (92)	None	95.3% (122)	90.6% (116)	
Malacia	7.8% (10)	23.4% (30)	Chondroplasty	4.7% (6)	4.7% (6)	
Debonding	0% (0)	0.8% (1)	Microfracture	0% (0)	4.7% (6)	
Cleavage	1.6% (2)	0.8% (1)				
Defect	1.6% (2)	3.1% (4)				

*N = 128 for each group. The values for the 2 groups are given as the percentage, with the number in parentheses.

or THA) were assigned an imputed mHHS value of 55 (n = 15 of 256, 5.9%) based on existing data in similar patients rather than excluding such patients from the anal-

ysis. Regarding cases of incomplete secondary PRO data, only matched pairs with complete data for each metric were included (n = 108 to 121).

Results

The cohort included 256 hips (128 with HA, 128 with SD) with a patient mean age of 24.9 years (range, 13 to 52 years) and a mean follow-up (and standard deviation) of 4.3 ± 2.4 years. FAI was classified as cam in 36.7%, combined cam/pincer in 56.3%, and isolated pincer in 7.0% of cases, including moderate cam morphology in 33.2% and global acetabular overcoverage in 7.4% of cases.

Prior to propensity matching, the standardized differences in baseline characteristics were generally high between hips undergoing HA versus SD (Table I). After propensity matching, the standardized differences were <0.1 for all covariates that were included in the match (Tables I and II). Hips included in the propensity-matched analysis had significantly different patient characteristics than those that were not included in the match (those without an appropriate match) (Table III). The patients who were treated with HA in the propensity analysis included greater proportions of men, those involved in competitive sports, those with combined FAI, those with a Tönnis osteoarthritis grade of 0, younger patients, and those with a higher preoperative mHHS score than the non-matched HA cases (all $p < 0.05$). The hips that were treated with SD in the propensity analysis included those with less global acetabular overcoverage,

TABLE V Differences in Postoperative PROs Between the Hip Arthroscopy and Surgical Dislocation Groups*

Postop. PRO	Difference†	P Value‡
mHHS	-1.0 (26.8)	0.67
HOOS domain		
ADL	0.9 (28.1)	0.75
Pain	2.9 (30.6)	0.32
QOL	3.7 (40.6)	0.34
Sport	5.8 (38.7)	0.12
Symptoms	2.8 (28.3)	0.30
SF-12 Mental	1.2 (11.8)	0.30
SF-12 Physical	1.5 (17.2)	0.36
UCLA activity score	0.1 (3.2)	0.63

*PRO = patient-reported outcome, mHHS = modified Harris hip score, HOOS = Hip disability and Osteoarthritis Outcome Score, ADL = activities of daily living, QOL = quality of life, SF-12 = Short Form-12, and UCLA = University of California Los Angeles. †The pairwise difference was calculated as surgical dislocation - hip arthroscopy, and is given as the mean and standard deviation. A positive difference indicates a larger value for surgical dislocation and a negative value indicates a larger value for hip arthroscopy. ‡Paired-sample t test.

TABLE VI Patient-Reported Outcomes at Preoperative and Postoperative Time Points*

	N	Surgical Dislocation		Change	Hip Arthroscopy		Change
		Preop.†	Postop.†		Preop.†	Postop.†	
mHHS	128	61.4 (14.7)	80.2 (19.5)	18.8	62.5 (15.9)	81.3 (20.4)	18.8
HOOS domain							
Pain	111	55.8 (20.1)	80.5 (22.3)	24.7	57.1 (21.1)	77.6 (22.3)	20.5
ADL	111	66.0 (20.7)	85.8 (21.1)	19.8	66.9 (21.7)	84.9 (20.8)	18.0
QOL	110	29.1 (19.8)	65.9 (28.1)	36.8	32.2 (19.8)	62.2 (30.6)	30.0
Symptoms	111	53.9 (18.8)	76.3 (21.5)	22.4	57.3 (22.3)	73.5 (21.8)	16.2
Sport	108	45.3 (22.8)	73.2 (27.7)	27.9	43.8 (21.6)	67.3 (30.1)	23.5
UCLA activity score	121	7.5 (2.7)	7.8 (2.2)	0.3	7.5 (2.7)	7.6 (2.6)	0.1
SF-12 Physical	111	38.7 (10.7)	47.6 (12.0)	8.9	38.2 (10.5)	46.1 (12.7)	7.9
SF-12 Mental	111	54.1 (10.1)	54.8 (7.9)	0.7	53.3 (8.7)	53.7 (8.4)	0.4

*mHHS = modified Harris hip score, HOOS = Hip disability and Osteoarthritis Outcome Score, ADL = activities of daily living, QOL = quality of life, UCLA = University of California Los Angeles, and SF-12 = Short Form-12. †The values are given as the mean and standard deviation.

more severe cam FAI, and older patient age than non-matched hips that were treated with SD (all $p < 0.05$). Intraoperative classification of labral, acetabular cartilage, and femoral cartilage lesions by the modified Beck classification demonstrated some differences between HA and SD in the matched groups (Table IV). Hips undergoing HA (when compared with those undergoing SD) were more likely to be classified with labral and acetabular cartilage pathology and undergo labral repair or acetabular chondroplasty, and less likely to report femoral head cartilage treatment (all $p < 0.05$) (Table IV).

The mean mHHS was similar at baseline between the 2 matched groups (62.5 with HA versus 61.4 with SD). Both groups demonstrated significant improvements in all PRO measures. The final mHHS was not significantly different between the 2 groups (81.3 versus 80.2, $p = 0.67$). Similarly, the HOOS pain subscale was not significantly different at the time of final follow-up (77.6 versus 80.5, $p = 0.32$). Also, the HOOS subscales for sport and recreation, activities of daily living, quality of life, and symptoms demonstrated no significant differences at the time of the final follow-up (Tables V and VI). No differences in the SF-12 physical and mental subscores were noted at the time of final follow-up.

No significant difference between groups was present for the rates of the 3 defined outcome states. The rate of THA

conversion was 0% for the HA group versus 3.1% for the SD group ($p = 0.12$) (Table VII). The rate of revision surgery was 7.8% for the HA group and 10.9% for the SD group ($p = 0.35$). The overall rate of persistent symptoms (reoperation or symptomatic hip) was 21.9% for the HA group and 24.4% for the SD group ($p = 0.55$).

Discussion

FAI is a common cause of hip pain and a recognized precursor to hip osteoarthritis²⁰⁻²². HA is increasingly being utilized for the treatment of FAI. Montgomery et al. found a 365% increase in HA from 2004 to 2009²³, while Maradit Kremers et al. saw an increase of 3.6 to 16.7 per 100,000 individuals per year from 2005 to 2013²⁴. Bozic et al. reported a fivefold increase in the use of HA to treat FAI from 2006 to 2010²⁵. While utilization of HA for the treatment of FAI continues to rise, SD continues to be used for the more severe or complex cases. There remains a lack of consensus regarding the indications for each procedure, and rigorous, direct comparisons between the outcomes of HA and SD are limited. Direct comparison studies are challenging and may be biased by differences in the patients in whom these procedures are performed. Thus, large patient populations with similar patients being treated with the 2 different procedures are required for meaningful outcome comparisons.

TABLE VII Differences Between Postoperative Outcome States for the Hip Arthroscopy and Surgical Dislocation Groups*

Outcome Variable	Surgical Hip Dislocation	Hip Arthroscopy	P Value
THA	3.1%	0%	0.12†
THA or revision	10.9%	7.8%	0.35
Persistent symptoms	24.4%	21.9%	0.55

*THA = total hip arthroplasty. †Analyzed with the Fisher exact test.

In our large multicenter study, after propensity matching to allow comparison of similar patients who were treated with HA and SD, no significant differences in PROs or rates of failure were seen. Our failure rate, as defined by reoperations or conversion to THA, was 7.8% for HA and 10.9% for SD and was not significantly different ($p = 0.35$) between the groups. Our findings were comparable with those reported in a systematic review that reported 90.5% survival for HA and 93.0% survival for SD⁸.

The literature directly comparing the outcome of HA versus SD remains sparse and often fails to control for differing patient populations undergoing the 2 procedures. Domb et al. compared the outcomes of the 2 treatment approaches in a single-surgeon cohort, with 10 patients undergoing SD and 20 age and sex-matched patients undergoing HA, but they failed to control for deformity severity and other differences in the groups⁸. They found no difference in the mHHS, but there were differences in the Non-Arthritic Hip Score (NAHS) ($p = 0.01$) and the Hip Outcome Score sport-specific subscale ($p = 0.047$), with HA patients scoring higher. The senior author performed the 10 SDs over a nearly 4-year study period during which 785 HAs were performed. Zingg et al. compared outcomes of 15 patients undergoing SD with 23 patients undergoing HA in a 2-surgeon study². They found a higher mHHS in the HA group compared with the SD group at 1 year postoperatively (93.4 versus 84.9, $p = 0.027$). Rego et al. compared the outcomes of 102 patients undergoing HA with those of 96 undergoing SD for FAI and found no difference in clinical outcomes between the groups as assessed by the NAHS⁹. Nwachukwu et al. performed a systematic review comparing 600 hips treated with SD at 57.0 months of follow-up with 1,484 hips treated with HA at 50.8 months of follow-up¹⁰. With conversion to THA as an end point, 93% survival was seen in the SD group and 90.5% ($p = 0.06$) survival was seen in the HA group. This large systematic review data set is very consistent with the findings of our large multicenter study.

In our study, the similar ultimate outcomes of patients undergoing treatment of FAI with either HA or SD support the continued use of both of these surgical procedures. It is important to recognize that this conclusion was applicable to the subgroup of patients commonly treated with either procedure during the study period. Additional research is needed to better define if other subsets of FAI patients are better treated with a specific technique. HA allows advantages in postoperative patient recovery and is the preferred method of treatment for most mild and moderate deformities. However, rates of failure with arthroscopic treatment remain at approximately 10% to 20%. This underscores the need for better understanding of the patient subgroups that are at risk for poor outcomes. Previous studies also have shown that inadequate deformity correction is a common reason for HA failure^{3,4} and emphasize the continued need for improved surgeon education and skills training. Severe FAI deformities, both cam and pincer, can be difficult to fully correct with HA even though the technical aspects of HA continue to improve^{5,6}. Inadequate osseous correction remains a common reason for failure of hip preservation surgery. Severe deformities may be more consistently corrected with SD.

Decision-making regarding HA versus SD is likely best undertaken with the expertise of the individual surgeon. The ability to perform adequate deformity correction is perhaps more important than the actual surgical approach. In cases where access for arthroscopic deformity correction is questionable, SD provides a very effective alternative with the potential for more comprehensive access to the FAI deformity. On the other hand, in hips with deformities that are adequately assessed by HA, HA may have advantages with respect to perioperative recovery and avoidance of a potential need for later hardware removal. With improved arthroscopic techniques, the limits of HA have improved and will continue to improve, although this remains surgeon-dependent.

The current study has several limitations. A randomized trial would be the optimal research design to answer the proposed questions but is likely not feasible given limited surgeon equipoise in this area. Through propensity matching, the current study provides a relevant comparison across a large number of patients at early postoperative time points. Propensity matching allows for limiting the bias between groups at baseline by controlling for known differences that cause bias. However, it is unable to control for unknown sources of bias that may still exist to some degree. The current multicenter study included a variety of hip preservation surgeons, including surgeons who perform HA, surgeons who perform open SD of the hip, and surgeons who perform both arthroscopic and open procedures. Although these different surgeon groups may influence the treatment and decision-making procedure, we saw no major effects in the current study. All surgeons treating patients in the current study were very experienced in the utilized surgical approach, which is important to understand when applying the results of the current study. In this study, we lacked radiographic confirmation of deformity correction, which may be influenced by surgical approach and affect outcomes. Thus, we were unable to assess the role of osseous correction in the outcomes. The study had a mean 4-year follow-up, and longer-term results may be needed to adequately demonstrate the differences between the 2 groups over time. Despite propensity matching, the study demonstrated significant differences in labral and cartilage pathology between the 2 groups. Given the highly matched baseline characteristics, we feel that this finding is likely due to arthroscopic visualization being more sensitive to early labral and cartilage disease in the central compartment than visualization during SD, while visualization during SD may be more sensitive to femoral head cartilage abnormalities. To our knowledge, the direct comparability of disease classification with HA or SD has not been previously established. However, the need to treat these earlier findings is unclear given the similar outcomes in the 2 groups in the current study. Additionally, the results of this study apply to hips that were able to be matched in the propensity analysis, whereas more extreme or very mild FAI cases may not have had adequate matching. In a propensity-matched study, the study design precludes the ability to perform subgroup analyses of all individual factors (as not all hips are matched on each factor

used in the matching, such as moderate cam or global overcoverage deformities) without losing the benefit of the propensity matching.

Conclusions

In a propensity-matched analysis, similar patients undergoing HA or SD demonstrated similar outcomes at a mean of 4 years postoperatively. Results of this multicenter study reinforce that both HA and SD of the hip are acceptable methods of treatment for FAI. ■

NOTE: The ANCHOR Group investigators on this project include Ernest L. Sink, MD; Daniel J. Sucato, MD, MS; David A. Podeszwa, MD; Perry L. Schoenecker, MD; Wudbhav N. Sankar, MD; Asheesh Bedi, MD; Eduardo N. Novais, MD; Travis H. Matheney, MD; and Etienne L. Belzile, MD.

Jeffrey J. Nepple, MD¹
Ira Zaltz, MD²
Christopher M. Larson, MD³
Paul E. Beaulé, MD, FRCSC⁴
Young-Jo Kim, MD⁵
Michael B. Millis, MD⁵
Rafael J. Sierra, MD⁶
John C. Clohisy, MD¹

¹Department of Orthopaedic Surgery, Washington University, St. Louis, Missouri

²Department of Orthopaedic Surgery, Beaumont Health System, Royal Oak, Michigan

³Twin Cities Orthopedics, Edina, Minnesota

⁴Department of Orthopaedic Surgery, The Ottawa Hospital, Ottawa, Ontario, Canada

⁵Department of Orthopaedic Surgery, Boston Children's Hospital and Harvard Medical School, Boston, Massachusetts

⁶Department of Orthopedic Surgery, Mayo Clinic, Rochester, Minnesota

Email address for J.C. Clohisy: jclohisy@wustl.edu

ORCID iD for J.J. Nepple: [0000-0002-7582-1415](https://orcid.org/0000-0002-7582-1415)
ORCID iD for I. Zaltz: [0000-0003-4036-6149](https://orcid.org/0000-0003-4036-6149)
ORCID iD for C.M. Larson: [0000-0002-9910-0145](https://orcid.org/0000-0002-9910-0145)
ORCID iD for P.E. Beaulé: [0000-0001-7667-9994](https://orcid.org/0000-0001-7667-9994)
ORCID iD for Y.-J. Kim: [0000-0002-0855-0168](https://orcid.org/0000-0002-0855-0168)
ORCID iD for M.B. Millis: [0000-0002-1380-5495](https://orcid.org/0000-0002-1380-5495)
ORCID iD for R.J. Sierra: [0000-0002-8513-1477](https://orcid.org/0000-0002-8513-1477)
ORCID iD for J.C. Clohisy: [0000-0001-7040-616X](https://orcid.org/0000-0001-7040-616X)

References

- Nepple JJ, Clohisy JC; ANCHOR Study Group Members. Evolution of femoroacetabular impingement treatment: the ANCHOR experience. *Am J Orthop* (Belle Mead NJ). 2017 Jan/Feb;46(1):28-34.
- Zingg PO, Ulbrich EJ, Buehler TC, Kalberer F, Poutawera VR, Dora C. Surgical hip dislocation versus hip arthroscopy for femoroacetabular impingement: clinical and morphological short-term results. *Arch Orthop Trauma Surg*. 2013 Jan;133(1):69-79. Epub 2012 Oct 14.
- Heyworth BE, Shindle MK, Voos JE, Rudzki JR, Kelly BT. Radiologic and intra-operative findings in revision hip arthroscopy. *Arthroscopy*. 2007 Dec;23(12):1295-302.
- Bogunovic L, Gottlieb M, Pashos G, Baca G, Clohisy JC. Why do hip arthroscopy procedures fail? *Clin Orthop Relat Res*. 2013 Aug;471(8):2523-9.
- Ahmad SS, Heilgmeier M, Anwender H, Beck M. Surgical hip dislocation is more powerful than arthroscopy for achieving high degrees of acetabular correction in pincer type impingement. *Orthop Traumatol Surg Res*. 2019 Nov;105(7):1339-44. Epub 2019 Sep 26.
- Bedi A, Zaltz I, De La Torre K, Kelly BT. Radiographic comparison of surgical hip dislocation and hip arthroscopy for treatment of cam deformity in femoroacetabular impingement. *Am J Sports Med*. 2011 Jul;39(Suppl):20S-8S.
- Büchler L, Neumann M, Schwab JM, Iselin L, Tannast M, Beck M. Arthroscopic versus open cam resection in the treatment of femoroacetabular impingement. *Arthroscopy*. 2013 Apr;29(4):653-60. Epub 2013 Feb 6.
- Domb BG, Stake CE, Botser IB, Jackson TJ. Surgical dislocation of the hip versus arthroscopic treatment of femoroacetabular impingement: a prospective matched-pair study with average 2-year follow-up. *Arthroscopy*. 2013 Sep;29(9):1506-13.
- Rego PA, Mascarenhas V, Oliveira FS, Pinto PC, Sampaio E, Monteiro J. Arthroscopic versus open treatment of cam-type femoroacetabular impingement: retrospective cohort clinical study. *Int Orthop*. 2018 Apr;42(4):791-7. Epub 2018 Jan 3.
- Nwachukwu BU, Rebolledo BJ, McCormick F, Rosas S, Harris JD, Kelly BT. Arthroscopic versus open treatment of femoroacetabular impingement: a systematic review of medium- to long-term outcomes. *Am J Sports Med*. 2016 Apr;44(4):1062-8. Epub 2015 Jun 9.
- Baser O. Too much ado about propensity score models? Comparing methods of propensity score matching. *Value Health*. 2006 Nov-Dec;9(6):377-85.
- Clohisy JC, Baca G, Beaulé PE, Kim YJ, Larson CM, Millis MB, Podeszwa DA, Schoenecker PL, Sierra RJ, Sink EL, Sucato DJ, Trousdale RT, Zaltz I; ANCHOR Study Group. Descriptive epidemiology of femoroacetabular impingement: a North American cohort of patients undergoing surgery. *Am J Sports Med*. 2013 Jun;41(6):1348-56. Epub 2013 May 13.
- Ganz R, Gill TJ, Gautier E, Ganz K, Krügel N, Berlemann U. Surgical dislocation of the adult hip a technique with full access to the femoral head and acetabulum without the risk of avascular necrosis. *J Bone Joint Surg Br*. 2001 Nov;83(8):1119-24.
- Beck M, Kalhor M, Leunig M, Ganz R. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br*. 2005 Jul;87(7):1012-8.
- Beck M, Leunig M, Parvizi J, Boutier V, Wyss D, Ganz R. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res*. 2004 Jan;418:67-73.
- Nepple JJ, Larson CM, Smith MV, Kim YJ, Zaltz I, Sierra RJ, Clohisy JC. The reliability of arthroscopic classification of acetabular rim labrochondral disease. *Am J Sports Med*. 2012 Oct;40(10):2224-9. Epub 2012 Aug 27.
- Austin PC. Using the standardized difference to compare the prevalence of a binary variable between two groups in observational research. *Commun Stat Simul Comput*. 2009;38(6):1228-34.
- Kemp JL, Collins NJ, Roos EM, Crossley KM. Psychometric properties of patient-reported outcome measures for hip arthroscopic surgery. *Am J Sports Med*. 2013 Sep;41(9):2065-73. Epub 2013 Jul 8.
- Chahal J, Van Thiel GS, Mather RC 3rd, Lee S, Song SH, Davis AM, Salata M, Nho SJ. The patient acceptable symptomatic state for the modified Harris hip score and hip outcome score among patients undergoing surgical treatment for femoroacetabular impingement. *Am J Sports Med*. 2015 Aug;43(8):1844-9. Epub 2015 Jun 15.
- Ganz R, Parvizi J, Beck M, Leunig M, Nötzli H, Siebenrock KA. Femoroacetabular impingement: a cause for osteoarthritis of the hip. *Clin Orthop Relat Res*. 2003 Dec;417:112-20.
- Beaulé PE, Speirs AD, Anwender H, Melkus G, Rakhra K, Frei H, Lamontagne M. Surgical correction of cam deformity in association with femoroacetabular impingement and its impact on the degenerative process within the hip joint. *J Bone Joint Surg Am*. 2017 Aug 16;99(16):1373-81.
- Wyles CC, Norambuena GA, Howe BM, Larson DR, Levy BA, Yuan BJ, Trousdale RT, Sierra RJ. Cam deformities and limited hip range of motion are associated with early osteoarthritic changes in adolescent athletes: a prospective matched cohort study. *Am J Sports Med*. 2017 Nov;45(13):3036-43. Epub 2017 Aug 18.
- Montgomery SR, Ngo SS, Hobson T, Nguyen S, Alluri R, Wang JC, Hame SL. Trends and demographics in hip arthroscopy in the United States. *Arthroscopy*. 2013 Apr;29(4):661-5. Epub 2013 Feb 1.
- Maradit Kremers H, Schilz SR, Van Houten HK, Herrin J, Koenig KM, Bozic KJ, Berry DJ. Trends in utilization and outcomes of hip arthroscopy in the United States between 2005 and 2013. *J Arthroplasty*. 2017 Mar;32(3):750-5. Epub 2016 Sep 28.
- Bozic KJ, Chan V, Valone FH 3rd, Feeley BT, Vail TP. Trends in hip arthroscopy utilization in the United States. *J Arthroplasty*. 2013 Sep;28(8)(Suppl):140-3. Epub 2013 Aug 1.