Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

1-1-2021

Taking account of asymptomatic infections: A modeling study of the COVID-19 outbreak on the Diamond Princess cruise ship

Li-Shan Huang

National Tsing Hua University, Hsinchu

Li Li

AT&T, Bedminster, New Jersey

Lucia Dunn

The Ohio State University

Mai He

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Huang, Li-Shan; Li, Li; Dunn, Lucia; and He, Mai, "Taking account of asymptomatic infections: A modeling study of the COVID-19 outbreak on the Diamond Princess cruise ship." PLoS One. 16, 3. e0248273 (2021). https://digitalcommons.wustl.edu/open_access_pubs/10239

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

RESEARCH ARTICLE

Taking account of asymptomatic infections: A modeling study of the COVID-19 outbreak on the Diamond Princess cruise ship

Li-Shan Huang^{1*}, Li Li², Lucia Dunn³, Mai He⁴

1 Institute of Statistics, National Tsing Hua University, Hsinchu, Taiwan, **2** AT&T, Bedminster, New Jersey, United States of America, **3** Department of Economics, The Ohio State University, Columbus, Ohio, United States of America, **4** Department of Pathology and Immunology, Washington University School of Medicine in St. Louis, St. Louis, Missouri, United States of America

* lhuan@stat.nthu.edu.tw

Abstract

The COVID-19 outbreak on the Diamond Princess (DP) cruise ship has provided empirical data to study the transmission potential of COVID-19 with the presence of pre/asymptomatic cases. We studied the changes in R_0 on DP from January 21 to February 19, 2020 based on chain binomial models under two scenarios: no quarantine assuming a random mixing condition, and quarantine of passengers in cabins—passengers may get infected either by an infectious case in a shared cabin or by pre/asymptomatic crew who continued to work. Estimates of R_0 at the beginning of the epidemic were 3.27 (95% CI, 3.02–3.54) and 3.78 (95% CI, 3.49–4.09) respectively for serial intervals of 5 and 6 days; and when quarantine started, with the reported asymptomatic ratio 0.505, R_0 rose to 4.18 (95%CI, 3.86–4.52) and 4.73 (95%CI, 4.37–5.12) respectively for passengers who might be exposed to the virus due to pre/asymptomatic crew. Results confirm that the higher the asymptomatic ratio is, the more infectious contacts would happen. We find evidence to support a US CDC report that “a high proportion of asymptomatic infections could partially explain the high attack rate among cruise ship passengers and crew.” Our study suggests that if the asymptomatic ratio is high, the conventional quarantine procedure may not be effective to stop the spread of virus.

OPEN ACCESS

Citation: Huang L-S, Li L, Dunn L, He M (2021) Taking account of asymptomatic infections: A modeling study of the COVID-19 outbreak on the Diamond Princess cruise ship. PLoS ONE 16(3): e0248273. <https://doi.org/10.1371/journal.pone.0248273>

Editor: Qiang Zeng, South China University of Technology, CHINA

Received: August 23, 2020

Accepted: February 23, 2021

Published: March 16, 2021

Copyright: © 2021 Huang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) file.

Funding: The Taiwan Ministry of Science and Technology provided funding for this study in the form of a grant awarded to author LH (MOST 107-2118-M-007-002-MY2). Indirect support was provided through salaries/facilities use by the home institutions of LH, LD, and MH. AT&T provided support in the form of salary for author LL. The specific roles of these authors are

Introduction

The COVID-19 outbreak has developed into an international public health emergency. The reproductive number (R_0) of COVID-19 is a key piece of information for understanding an epidemic. Current intervention methods focus on quarantine methods with either mitigation or suppression strategies aimed at reducing the reproduction number R_0 and flattening the curve [1]. Asymptomatic or pre-symptomatic infectious cases are less likely to seek medical care or to be tested and quarantined, contributing to the infectious potential of a respiratory virus [2,3]. Clinical findings have suggested that the viral load in asymptomatic patients is similar to that in symptomatic patients [4]. Evidence suggests that these pre/asymptomatic patients can infect others before they manifest any symptoms [5–7]. In an earlier study [8] in

articulated in the 'author contribution' section. The funders did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. There was no additional external funding received for this study.

Competing interests: The authors have read the journal's policy. We wish to confirm that there are no known conflicts of interest associated with this paper and there has been no significant financial support for this work that could have influenced its outcome. AT&T provided the salary for LL but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Wuhan, China, 200 (83%) individuals out of 240 reported no exposure to an individual with respiratory symptoms, which suggests pre/asymptomatic infection is common [9]. Little is known on the implications of asymptomatic COVID-19 transmission on disease dynamics [10]. The Diamond Princess (DP) data [11–14] with reported asymptomatic cases may be considered as an “accidental” trial in an isolated environment. Based on the DP data [11–13], we estimate the R_0 as a function of time, and our approaches take explicit account of possibly infectious contacts between quarantined passengers in cabins and pre/asymptomatic crew, which has not been explored in the literature. A US CDC report states that “a high proportion of asymptomatic infections could partially explain the high attack rate among cruise ship passengers and crew” [15].

On January 20, 2020, the DP departed Yokohama, Japan, making stops in Hong Kong, Vietnam, Taiwan, Japan, and was scheduled to return to Yokohama on February 4 [12]. The DP [11–15], with 3,711 people (2,666 passengers and 1,045 crew members) on board as of February 5, 2020, was found to have an outbreak of COVID-19 from one traceable passenger from Hong Kong. This passenger became symptomatic on January 23 and disembarked on January 25 in Hong Kong. On February 1, six days after leaving the ship, he tested positive for SARS-CoV-2 at a Hong Kong hospital [12]. Japanese authorities were informed about this test result. Group activities continued on board through February 4, when the authorities announced positive test results for SARS-CoV-2 for another ten people on board. The ship was quarantined by the Japanese Ministry of Health, Labour and Welfare for what was expected to be a 14-day period (5–19 February), off the Port of Yokohama [12]. Initially, passengers were quarantined in their cabins while the crew continued to work [14]. Only symptomatic cases and close contacts were tested for COVID-19 and PCR-confirmed positive passengers were removed and isolated in Japanese hospitals. As reported [11,15], phased attempts were made to test all passengers including asymptomatic cases starting on February 11. As of February 20, 619 cases had been confirmed (16.7% of the population on board), including 82 crew and 537 passengers [11]; and 50.5% of the COVID-19 cases on the DP were asymptomatic (consisting of both true asymptomatic and pre-symptomatic infections) [13], while an estimated proportion 17.9% (95% credible interval: 15.5–20.2%) never developed symptoms [13]. Of 66 SARS-CoV-2 positive American DP travelers with complete symptom information, 14 (21%) were pre-symptomatic while on the ship [16]. Overall, 712 (19.2%) of the crew and passengers tested positive; of these, 331 (46.5%) were asymptomatic at the time of testing [15].

The R_0 of COVID-19 on DP has been estimated previously [17]; this research identified the R_0 as 14.8 initially and then declining to a stable 1.78 after the quarantine and removal interventions, assuming a 70% reduction in contact rate. That research does not take account of asymptomatic infections. Other researchers using the DP data up to February 16 have estimated the median R_0 as 2.28 [18]. They found R_0 remained high despite quarantine measures, while concluding that estimating R_0 was challenging due to the difficulty in identifying the exact number of infected cases. The R_0 values have important implications for predicting the effects of interventions. The threshold for combined vaccine efficacy and herd immunity needed for disease extinction is $1-1/R_0$. At $R_0 = 2$, the threshold is 50%, while at $R_0 = 4$, this threshold increases to 75%.

We investigated the changes in R_0 for COVID-19 on the DP from January 21 to February 19 with a chain binomial model [19–21] at different times under two scenarios: no quarantine assuming a random mixing condition before February 5, and quarantine of passengers in cabins from February 5 to 19—passengers may get infected either by an infectious case in a shared cabin or by pre/asymptomatic crew who continued to work. This work adds to the growing knowledge gained from the DP data in that we estimate R_0 by (1) mimicking the quarantine

conditions in practice, and (2) taking explicit account of the presence of the pre/asymptomatic crew and phased removal of infectious cases.

The chain binomial model originally proposed in [19], belongs to the broader class of stochastic discrete-time SIR models, and has been commonly used in the analysis of infectious disease spread [20,21] such as measles [20,22] and ebola [20]. The chain binomial model occurs naturally in cohort studies such as the DP case considered here where the number of individuals at risk in one serial interval are the survivors from the previous serial interval so that the conditional distributions are binomial. Each binomial probability mass function is conditioned on previous function and “products of these probability mass functions give the probabilities of particular sequences of binomial realizations” [21]. “The simple chain structure allows for statistical inference based on likelihood theory” [21]. In our analysis for the DP data, the novelty of the work is that we incorporate the asymptomatic ratio in estimating the R_0 based on the chain binomial model, which has not been explored in the literature.

Materials and methods

Data

We collected publicly available data on the outbreak on the DP from January 21 to February 19 [11–13]. We set January 21 as day 1, since January 20 was the start date (day 0) of the cruise. February 19 (day 30) was the date that most passengers were allowed to leave the ship. For those dates that Y , the number of new COVID-19 cases, was not reported, linear interpolation was used. As an example, there were 67 new cases on February 15, but no data were reported on February 14. After linear interpolation, Y on a daily basis became 33 and 34 for February 14 and 15 respectively. Based on the documented onset dates [11], there were 34 cases with onset dates before February 6, and we further adjusted the number of confirmed cases on February 3, 6, and 7, from 10, 10, and 41 cases to 17, 17, and 27 cases respectively. We chose serial intervals τ of 5 and 6 days as these are factors of 30 and are close to 7.5 days (95% CI 5.3–19) [8] and 4 days [23,24]. Then daily data were aggregated into 5- and 6-day intervals as binomial realizations. A [S1 File](#) contains R-code [25] for reproducing our calculations.

Chain binomial model with asymptomatic ratio

The chain binomial model assumes that an epidemic is formed from a succession of generations of infectious individuals from a binomial distribution [20,21]. For the DP data, the initial population size is $N_{t=0} = 3711$, where time t is the duration measured in units of the serial interval. To model the dynamics on the ship for the case $\tau = 6$, we make the following assumptions, which are stated in the order of time.

- (a) From January 21 to 26 ($t = 1$, the first serial interval), infection contacts happened at random following the random mixing assumption. Let I_t be the number of persons infected at time t . Then $I_{t=1}$ is a binomial random variable $B(N_{t=0}, p_1)$ with binomial transmission probability $p_1 = 1 - \exp(-\beta \times I_{t=0}/N_{t=0})$, where β is the transmission rate and $I_{t=0} = 1$ (the first case who disembarked on January 25). As in the SIR model, the probability that a subject escapes infectious contact is assumed to be $\exp(-\beta \times I_{t=0}/N_{t=0})$.
- (b) From January 27 to February 1 ($t = 2$), infection contacts again followed the random mixing assumption. Hence $I_{t=2}$ is a binomial random variable $B(N_{t=1}, p_2)$ with $p_2 = 1 - \exp(-\beta \times I_{t=1}/N_{t=0})$, and the number of persons at risk of infection is $N_{t=1} = 3711 - I_{t=1}$.
- (c) For the period February 2 to 7 ($t = 3$), $I_{t=3}$ is a binomial random variable $B(N_{t=2}, p_3)$ with $N_{t=2} = N_{t=1} - I_{t=2}$ and $p_3 = 1 - \exp(-\beta \times (I_{t=1} + I_{t=2})/N_{t=0})$. As the quarantine started

on February 5 and confirmed cases were removed, the number of persons infected, removed, and at risk of infection at the end of the $t = 3$ period were $I_{t=3}$, $(I_{t=1} + I_{t=2})$, and $N_{t=3} = N_{t=2} - I_{t=3}$ respectively.

- (d) For $t = 4$ and $t = 5$ during the quarantine of passengers, $N_t = N_{t-1} - I_t$, and infectious cases were removed. We further make the following assumptions. (i) Of all infected cases, 86.8% ($= 537/619$) were passengers and 13.2% ($= 82/619$) crew [11]. We use these proportions to calculate the number of infected persons in each group, $I_{p,t}$ and $I_{c,t}$, respectively, and $I_t = I_{p,t} + I_{c,t}$. This assumption is imposed since there is no public data available for the time course of $I_{p,t}$ and $I_{c,t}$. (ii) Crew members continued to work unless showing symptoms; hence the binomial transmission probability of crew remained the same as $1 - \exp(-\beta \times I_{t-1}/N_{t-1})$, $t = 4$ and 5 . In other words, crew were randomly mixing in the population on board. (iii) Passengers stayed in cabins most of the time. Assume that among infected passengers $I_{p,t}$, the proportion of infections that occurred in cabins is r_p , and that the average occupancy per cabin is 2. For those $I_{p,t-1}$ cases, $t = 4, 5$, the binomial transmission probability to infect $I_{p,t} \times r_p$ passengers in cabins is $1 - \exp(-\beta/2)$. In [11], $r_p = 0.2$ ($= 23/115$), while we assume $r_p = 0.2$ and 0.3 when $\tau = 6$. For this assumption, $r_p \leq I_{p,t-1}/I_{p,t}$, and thus r_p cannot be made arbitrarily large. (iv) The other $(1 - r_p)$ proportion of infected passengers' cases was possibly due to pre/asymptomatic crew members who continued to perform service [11], and their binomial transmission probability is assumed to be $p_t = 1 - \exp(-\beta \times aratio \times I_{c,t-1}/C_{t-1})$, where *aratio* is the pre/asymptomatic ratio and C_{t-1} is the number of crew members on board at time $t - 1$. That is to say, these passengers were randomly mixing in the crew population with possible infectious contact with pre/asymptomatic crew. In our calculations, *aratio* = 0.4, 0.465 [15], 0.505 [13], and 0.6.

Assumptions (a)–(c) correspond to no quarantine assuming a random mixing condition, and assumption (d) to quarantine of passengers in cabins in which passengers may either get infected (d)(iii) by an infectious case in a shared cabin or (d)(iv) by pre/asymptomatic crew who continued to work. The maximum likelihood (ML) approach was used to estimate β . We developed some R [25] code by modifying some R-functions in [20] for the chain binomial model and the ML step was carried out using the R-package bbmle [26]. The associated R-code is provided in the S1 File. For $t = 4, 5$, R_0 is the number of persons (passengers or crew) at risk (at time t) times either (d)(iv) $1 - \exp(-\beta \times aratio/C_{t-1})$ for passengers potentially infected by pre/asymptomatic crew, or (d)(ii) $1 - \exp(-\beta/N_{t-1})$ for crew. In the case of $\tau = 6$, $r_p = 0.2$ [11], and *aratio* = 0.505 [13], 100 stochastic simulations of the chain binomial model with $N = 3,700$ and the estimated β are performed to visually compare the simulated uncontrolled epidemics and observed DP data.

The calculation for the case $\tau = 5$ is analogous: period January 21 to 25 follows (a); period January 26 to 30 follows (b); period January 31 to February 4 follows (c); and periods February 5 to 9, February 10 to 14, and February 15 to 19 follow (d), and $r_p = 0.2$, which is the maximum value given the constraint $r_p \leq I_{p,t-1}/I_{p,t}$.

Results

Estimation of R_0

For the DP COVID-19 outbreak, Table 1 gives the estimates of β and their 95% confidence intervals. Since β is the basic reproductive number R_0 at the beginning of the epidemic ($t = 1, 2, 3$ when $\tau = 5$, and $t = 1, 2$ when $\tau = 6$), we observe from Table 1 that the estimated R_0 for the initial period is greater than 3 in every one of the scenarios that we considered. In addition, given τ and r_p , the estimated β increases as *aratio* decreases. Thus if the *aratio* is smaller than

Table 1. Estimates of β and their 95% confidence intervals for the Diamond Princess COVID-19 outbreak data. $\beta = R_0$ at $t = 1, 2$ when $\tau = 6$; $\beta = R_0$ at $t = 1, 2, 3$ when $\tau = 5$.

<i>aratio</i>	estimates of β (95% CI)		
	$\tau = 6$; $r_p = 0.2^c$	$\tau = 6$; $r_p = 0.3$	$\tau = 5$; $r_p = 0.2^c$
40%	3.94 (3.64, 4.27)	4.41 (4.06, 4.78)	3.41 (3.15, 3.69)
46.50% ^a	3.84 (3.54, 4.16)	4.28 (3.94, 4.64)	3.33 (3.07, 3.60)
50.50% ^b	3.78 (3.49, 4.09)	4.20 (3.87, 4.55)	3.27 (3.02, 3.54)
60%	3.64 (3.36, 3.94)	4.03 (3.71, 4.36)	3.16 (2.91, 3.42)

^a Taken from [15].

^b Taken from [13].

^c Taken from [11].

<https://doi.org/10.1371/journal.pone.0248273.t001>

40%, the estimated β would be larger than those in Table 1. With $r_p = 0.2$ in Table 1, comparing the estimated R_0 between $\tau = 5$ and $\tau = 6$, a longer interval leads to a larger estimate of R_0 and vice versa.

When *aratio* = 0.505 [13], the estimated R_0 as a function of t and its 95% CI are given in Table 2 and illustrated in Fig 1 for the case of $r_p = 0.2$. We observe that when $\tau = 6$ and $r_p = 0.2$, the R_0 for passengers in (d)(iv) is increased from 3.78 at $t = 3$ to 4.73 and 4.39 at $t = 4, 5$, respectively, and the R_0 is decreased to 1.06 and 1.05 respectively for crew in (d)(ii). This shows that R_0 for some passengers increased from $t = 3$ to $t = 4, 5$ if they were in contact with pre/asymptomatic crew. On the other hand, the R_0 for crew at $t = 4, 5$ is small and close to 1, since infected passengers were removed and crew were exposed to fewer cases. With a higher $r_p = 0.3$, the same $\tau = 6$ and *aratio* = 0.505, the estimated $\beta = 4.20$ (Table 1) is larger than the case with $r_p = 0.2$, and the R_0 for passengers in (d)(iv) is increased to 5.26 and 4.88 at $t = 4$ and 5 respectively (Table 2); and the R_0 for crew in (d)(ii) is decreased to 1.18 and 1.17 respectively. For the case of $\tau = 5$, $r_p = 0.2$, *aratio* = 0.505, and estimated $\beta = 3.27$ (Table 1), Table 2 shows that the R_0 for passengers in (d)(iv) is again increased to 4.18, 4.08, and 3.74 at $t = 4, 5$, and 6 respectively, and the R_0 for crew in (d)(ii) is below 1, 0.92, 0.92, and 0.91 respectively.

Table 2. Estimates of R_0 as a function of t and their 95% confidence intervals for the Diamond Princess COVID-19 outbreak when *aratio* = 0.505^a.

		R_0 as a function of t ; 95%CI		
time interval		2/2-2/7 ($t = 3$)	2/8-2/13 ($t = 4$)	2/14-2/19 ($t = 5$)
$\tau = 6$; $r_p = 0.2^b$	passengers in (d)(iv)	3.78 (3.49, 4.09)	4.73 (4.37, 5.12)	4.39 (4.06, 4.75)
	crew		1.06 (0.98, 1.15)	1.05 (0.97, 1.14)
	combined		2.90 (2.67, 3.13)	2.73 (2.52, 2.95)
$\tau = 6$; $r_p = 0.3$	passengers in (d)(iv)	4.20 (3.87, 4.55)	5.26 (4.84, 5.69)	4.88 (4.50, 5.29)
	crew		1.18 (1.08, 1.28)	1.17 (1.08, 1.27)
	combined		3.22 (2.97, 3.49)	3.03 (2.79, 3.28)
time interval		2/5-2/9 ($t = 4$)	2/10-2/14 ($t = 5$)	2/15-2/19 ($t = 6$)
$\tau = 5$; $r_p = 0.2^b$	passengers in (d)(iv)	4.18 (3.86, 4.52)	4.08 (3.77, 4.42)	3.74 (3.45, 4.04)
	crew	0.92 (0.85, 1.00)	0.92 (0.85, 0.99)	0.91 (0.84, 0.98)
	combined	2.55 (2.36, 2.76)	2.50 (2.31, 2.71)	2.32 (2.15, 2.52)

^a Taken from [13].

^b Taken from [11].

<https://doi.org/10.1371/journal.pone.0248273.t002>

Fig 1. Time-dependent effective reproduction number R_0 (solid lines) of COVID-19 on board the Diamond Princess ship January 21 (day 1) to February 19 (day 30) and their 95% confidence intervals (dash lines), assuming $\tau = 5$ and 6 days, $r_p = 0.2$ and $aratio = 0.505$. Blue: R_0 for passengers in contact with asymptomatic crew members. Red: R_0 for crew members.

<https://doi.org/10.1371/journal.pone.0248273.g001>

Other than those infections between passengers sharing the same cabin, the combined R_0 for passengers and crew are also given in Table 2, 2.90 and 2.73 respectively for $t = 4$ and 5 when $\tau = 6$, $r_p = 0.2$, and $aratio = 0.505$, decreasing from the initial $R_0 = 3.78$, which illustrates the limited effects of quarantine if pre/asymptomatic cases were present. Similarly, when $\tau = 5$, $r_p = 0.2$, and $aratio = 0.505$, the combined R_0 for passengers and crew is 2.55, 2.50, and 2.32 respectively for $t = 4, 5$ and 6. To understand the dynamics of no quarantine with a high R_0 , Fig 2 shows 100 stochastic simulations of the chain binomial model based on $N = 3,700$ and $\beta = 3.78$, assuming no quarantine, infected cases removed, $\tau = 6$, and extrapolation to 90 days, with the observed epidemic (red line). It suggests that the quarantine on DP did prevent a more serious outbreak. If there was no quarantine, the cumulative number of cases at the end of 30 days has a mean 856 (SD = 440), and median 791 (IQR = 538), while the observed DP data of 621 cases [11] is at the 35th percentile. Among 99 of 100 simulations, the entire population is infected at the end of 54 days, and in the remaining one simulation, the entire population is immune, not infected at all.

Mathematical explanation

Let us explain mathematically why the R_0 increased for passengers in contact with pre/asymptomatic crew. Denote the number of passengers in assumption (d)(iv) as Pa_t . To reduce their R_0 to a number smaller than the initial β , a sufficient condition is $aratio \leq C_{t-1}/Pa_t$, and the DP crew-passenger ratio at $t = 0$ is $1045/2666 = 0.39$ [12]. This suggests that as long as the $aratio$ is $\geq 40\%$, R_0 for passengers in (d)(iv) would remain high due to pre/asymptomatic crew. The higher the $aratio$ is, the more infectious contacts would happen. Thus, for a virus with a high $aratio$, the conventional quarantine procedure may not be effective to stop the spread of virus, highlighting the importance of early and effective surveillance.

We then derived a sufficient condition to achieve $R_0 \leq 1$ for both passengers and crew: $\beta \times aratio \times Pa_t \leq C_{t-1}$ and $C_t \leq N_{t-1}/\beta$. We attempt to interpret this condition as follows. For a β

Fig 2. 99 of 100 stochastic simulations of the chain binomial model based on $N = 3700$ and $\beta = 3.78$, assuming no quarantine, serial interval $\tau = 6$ days, infected cases removed, and extrapolation to 90 days, with the observed epidemic (red line). For the remaining one simulation, the entire population is immune, not infected at all.

<https://doi.org/10.1371/journal.pone.0248273.g002>

about 3, $C_t \leq N_{t-1}/\beta$ means that the number of people who continued to work during quarantine is less than one-third of the population, which is generally the case during quarantine. However, $\beta \times aratio \times Pa_t \leq C_{t-1}$ may not be satisfied depending on the *aratio* value. When the *aratio* is close to 0, this condition is satisfied, but when *aratio* is high, the pre/asymptomatic crew continue to spread the virus and passengers staying in cabins could not escape infectious contacts. This suggests that if the true *aratio* value is high, the current “stay-at-home” quarantine procedure may not be sufficient to reduce $R_0 \leq 1$ and to eliminate the virus completely.

Discussion

The effects of pre/asymptomatic population on the spread of COVID-19 during quarantine continues to provide a research case with great possibilities for gaining a better understanding of the pandemic. Clinical observations and lab tests have confirmed the existence of a pre/asymptomatic population infecting others [5–7]. It is not easy to give estimates of the size of this population, yet the DP outbreak provides useful real world data for this. 50.5% of passengers and crew members on the DP were pre/asymptomatic and an estimated 17.9% of the infected individuals never developed symptoms [13]. In a subsample of American DP travelers [16], 14 (21%) cases were pre-symptomatic while on the ship. In a retrospective study [27] of 104 DP COVID-19 cases, asymptomatic cases showed milder CT severity score than symptomatic cases.

In this study, the estimated R_0 for the initial period (Table 1) are all greater than 3, consistent with most estimates of R_0 reported earlier, showing that the COVID-19 virus is highly contagious [28–30]. The novelty of our approach has been to incorporate the pre/asymptomatic ratio into the chain binomial model to account for the possibly infectious contacts between quarantined passengers and pre/asymptomatic crew. The results show that with a serial interval of 6 days, R_0 is similar for $t = 1–3$, yet R_0 for some passengers in assumption (d)(iv) is higher for $t = 4, 5$. The results suggest that the observed proportions of infections, 86.8% (= 537/619) for passengers and 13.2% (= 82/619) for crew [11], is possible and we find evidence

to support a US CDC report that “a high proportion of asymptomatic infections could partially explain the high attack rate among cruise ship passengers and crew” [15].

Some research has suggested that the pre/asymptomatic population, “silent carriers,” are the main driving force behind this pandemic. A group [3] has estimated that the proportion of undocumented infections in China—including those who experience mild, limited or no symptoms and go undiagnosed—could be as high as 86% prior to January 23, 2020. They estimated the transmission rate of undocumented infections as 55% of the rate for documented infections, and yet that undocumented infections contributed to 79% of documented cases. Another group of researchers found that the total contribution from the pre/asymptomatic population is more than that of symptomatic patients [9]. Future studies to estimate the pre/asymptomatic ratio and the time when asymptomatic persons become infectious are needed to evaluate the effects of various control strategies [10,31,32].

The strength of this analysis is that it incorporates pre/asymptomatic infections in the DP data in a way not explored earlier. However, there are also limitations. First, due to inadequate data on the time course of infection cases among crew and passengers, assumption (d)(i) assumes a constant proportion, which may vary with time in practice. Second, the values of the parameter r_p assumed in the present study may not be sufficiently large due to mathematical constraints. A study [16] based on a subsample of American passengers on DP reported a high attack of 63% (27/43) for those sharing a cabin with an asymptomatic infected cabinmate. Third, the assumptions (a)-(d) under chain binomial models may not be sufficient to capture the complexity of the COVID-19 epidemics. Fuller data reporting is important for researchers to develop statistical methodology to help combat this pandemic.

Almost all of the passengers on DP were tested before they were evacuated. However, it is impractical to test everyone in the real world, especially for those pre/asymptomatic cases. On DP, crew members continued to perform service unless they showed symptoms. This provides a parallel to people doing “essential work” in society and thus exempt from shelter-in-place rules. Our study suggests that if the pre/asymptomatic ratio is high, the conventional quarantine might not be sufficient to reduce R_0 to below 1, implying that a combination of preventive measures is needed to stop the spread of virus [32]. The DP was docked in Vietnam and Taiwan on January 27–28 and January 31, 2020, respectively, and yet both reported low COVID-19 incidence rates [33], suggesting that the virus can be contained with early and appropriate measures.

Supporting information

S1 File. File containing data and R-code related to this article.
(PDF)

Author Contributions

Conceptualization: Li Li, Lucia Dunn, Mai He.

Data curation: Li-Shan Huang, Li Li.

Formal analysis: Li-Shan Huang.

Funding acquisition: Li-Shan Huang.

Investigation: Li-Shan Huang, Li Li, Lucia Dunn, Mai He.

Methodology: Li-Shan Huang, Li Li, Lucia Dunn.

Project administration: Li Li, Mai He.

Software: Li-Shan Huang.

Writing – original draft: Li-Shan Huang, Li Li, Lucia Dunn.

Writing – review & editing: Li-Shan Huang, Li Li, Lucia Dunn, Mai He.

References

1. Ferguson N, Laydon D, Nedjati-Gilani G, Imai N, Ainslie K, Baguelin M, et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. Imperial College London. 2020. Available from: <https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-NPI-modelling-16-03-2020.pdf>.
2. Munster VJ, Koopmans M, van Doremalen N, van Riel D, and de Wit E. A novel coronavirus emerging in China—key questions for impact assessment. *N Engl J Med* 2020; 382(8):692–694. <https://doi.org/10.1056/NEJMp2000929> PMID: 31978293
3. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science* 2020; 368(6490):489–493. <https://doi.org/10.1126/science.abb3221> PMID: 32179701
4. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020; 382(12):1177–1179. <https://doi.org/10.1056/NEJMc2001737> PMID: 32074444
5. Zhang J, Tian S, Lou J, Chen Y. Familial cluster of COVID-19 infection from an asymptomatic. *Critical Care* 2020; 24(1):119. <https://doi.org/10.1186/s13054-020-2817-7> PMID: 32220236
6. Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *N Engl J Med* 2020; 382(10):970–971. <https://doi.org/10.1056/NEJMc2001468> PMID: 32003551
7. Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, et al. Presumed asymptomatic carrier transmission of COVID-19. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.2565> PMID: 32083643
8. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020; 382(13):1199–1207. <https://doi.org/10.1056/NEJMoa2001316> PMID: 31995857
9. Ferretti L, Wymant C, Kendall M, Zhao L, Nurtay A, Abeler-Dörner L, et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science* 2020:eabb6936. <https://doi.org/10.1126/science.abb6936> PMID: 32234805
10. Park SW, Cornforth DM, Dushoff J, Weitz JS. The time scale of asymptomatic transmission affects estimates of epidemic potential in the COVID-19 outbreak. *Epidemics* 2020, 100392. <https://doi.org/10.1016/j.epidem.2020.100392> PMID: 32446187
11. National Institute of Infectious Disease. Field briefing: Diamond Princess COVID-19 cases, 20 Feb update. <https://www.niid.go.jp/niid/en/2019-ncov-e/9417-covid-dp-fe-02.html>. Published February 20, 2020. Accessed August 5, 2020.
12. National Institute of Infectious Disease. Field briefing: Diamond Princess COVID-19 cases. <https://www.niid.go.jp/niid/en/2019-ncov-e/9407-covid-dp-fe-01.html>. Published February 19, 2020. Accessed August 5, 2020.
13. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill* 2020; 25(10). <https://doi.org/10.2807/1560-7917.ES.2020.25.10.2000180>
14. Kakimoto K, Kamiya H, Yamagishi T, Matsui T, Suzuki M, Wakita T. Initial investigation of transmission of COVID-19 among crew members during quarantine of a cruise ship—Yokohama, Japan, February 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69:312–313. <https://doi.org/10.15585/mmwr.mm6911e2> PMID: 32191689
15. Moriarty LF, Plucinski MM, Marston BJ, Kurbatova EV, Knust B, Murray EL, et al. Public health responses to COVID-19 outbreaks on cruise ships—worldwide, February–March 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69(12). <https://doi.org/10.15585/mmwr.mm6912e3> PMID: 32214086
16. Plucinski MM, Wallace M, Uehara A, Kurbatova EV, Tobolowsky FA, Schneider ZD, et al. Coronavirus disease 2019 (COVID-19) in Americans aboard the Diamond Princess cruise ship. *Clin Infect Dis* 2020; ciaa1180. <https://doi.org/10.1093/cid/ciaa1180> PMID: 32785683
17. Rocklöv J, Sjödin H, Wilder-Smith A. COVID-19 outbreak on the Diamond Princess cruise ship: estimating the epidemic potential and effectiveness of public health countermeasures. *J Travel Med* 2020; 27(3),taaa030. <https://doi.org/10.1093/jtm/taaa030> PMID: 32109273

18. Zhang S, Diao M, Yu W, Pei L, Lin Z, and Chen D. Estimation of the reproductive number of novel coronavirus (COVID-19) and the probable outbreak size on the Diamond Princess cruise ship: a data-driven analysis. *Int J Infect Dis* 2020; 93:201–204. <https://doi.org/10.1016/j.ijid.2020.02.033> PMID: 32097725
19. Bailey NTJ. *The Mathematical Theory of Epidemics*. London: Griffin; 1957.
20. Bjørnstad ON. *Epidemics: Models and Data using R*. Springer; 2018.
21. Longini IM. Chain binomial model. In Armitage P, Colton T, editors. *The Encyclopedia of Biostatistics*, Volume 1. New York: Wiley, 1998. pp. 593–598.
22. Lytras T, Theano G, Sotirios T. A modified chain binomial model to analyse the ongoing measles epidemic in Greece, July 2017 to February 2018. *Euro Surveill* 2018; 23(17) 18–00165. <https://doi.org/10.2807/1560-7917.ES.2018.23.17.18-00165> PMID: 29717695
23. Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-19) infections. *Int J Infect Dis* 2020; 93:284–286. <https://doi.org/10.1016/j.ijid.2020.02.060> PMID: 32145466
24. Du Z, Xu X, Wu Y, Wang L, Cowling BJ, and Meyers LA. Serial interval of COVID-19 among publicly reported confirmed cases. *Emerg Infect Dis* 2020; 26(6):1341–1343. <https://doi.org/10.3201/eid2606.200357> PMID: 32191173
25. R Core Team. *R: A Language and Environment for Statistical Computing*. 2020. Vienna, Austria: R Foundation for Statistical Computing.
26. Bolker B. *bbmle: Tools for general maximum likelihood estimation*. <https://cran.r-project.org/web/packages/bbmle/>. Published February 3, 2020. Accessed March 3, 2020.
27. Inui S, Fujikawa A, Jitsu M, Kunishima N, Watanabe S, Suzuki Y et al. Chest CT findings in cases from the cruise ship “Diamond Princess” with coronavirus disease 2019 (COVID-19). *Radiol Cardiothorac Imaging* 2020; 2(2):e200110.
28. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med* 2020; 27(2). <https://doi.org/10.1093/jtm/taaa021> PMID: 32052846
29. Ranjan R. Predictions for COVID-19 outbreak in India using epidemiological models. *medRxiv* 2020. <https://doi.org/10.1101/2020.04.02.20051466>
30. Zhao S, Lin Q, Ran J, Musa SS, Yang G, Wang W, et al. Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. *Int J Infect Dis* 2020; 92:214–217. <https://doi.org/10.1016/j.ijid.2020.01.050> PMID: 32007643
31. Dobrovolny HM. Modeling the role of asymptomatics in infection spread with application to SARS-CoV-2. *PloS ONE*. 2020; 15(8):e0236976. <https://doi.org/10.1371/journal.pone.0236976> PMID: 32776963
32. Johansson MA, Quandelacy TM, Kada S, et al. SARS-CoV-2 Transmission From People Without COVID-19 Symptoms. *JAMA Netw Open*. 2021; 4(1):e2035057. <https://doi.org/10.1001/jamanetworkopen.2020.35057> PMID: 33410879
33. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis*. 2021. [https://doi.org/10.1016/S1473-3099\(20\)30120-1](https://doi.org/10.1016/S1473-3099(20)30120-1) PMID: 32087114