

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2009

Longitudinal trends in the association of metabolic syndrome with 550 k single-nucleotide polymorphisms in the Framingham Heart Study

Yong-Moon Park
Washington University School of Medicine

Michael A. Province
Washington University School of Medicine

Xiaoyi Gao
Washington University School of Medicine

Mary Feitosa
Washington University School of Medicine

Jun Wu
Washington University School of Medicine

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs



Part of the [Medicine and Health Sciences Commons](#)

Please let us know how this document benefits you.

Recommended Citation

Park, Yong-Moon; Province, Michael A.; Gao, Xiaoyi; Feitosa, Mary; Wu, Jun; Ma, Duanduan; Rao, DC; and Kraja, Aldi T., "Longitudinal trends in the association of metabolic syndrome with 550 k single-nucleotide polymorphisms in the Framingham Heart Study." *BMC Proceedings*. 3, Suppl 7. S1. (2009).
https://digitalcommons.wustl.edu/open_access_pubs/344

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Yong-Moon Park, Michael A. Province, Xiaoyi Gao, Mary Feitosa, Jun Wu, Duanduan Ma, DC Rao, and Aldi T. Kraja

Longitudinal trends in the association of metabolic syndrome with 550 k single-nucleotide polymorphisms in the Framingham Heart Study

Yong-Moon Park^{1,2}, Michael A Province³, Xiaoyi Gao³, Mary Feitosa³, Jun Wu³, Duanduan Ma³, DC Rao⁴ and Aldi T Kraja^{*3}

Addresses: ¹GEMS Training Program, Washington University School of Medicine, 660 South Euclid, St. Louis, Missouri 63110, USA, ²Department of Preventive Medicine, College of Medicine, The Catholic University of Korea, Seoul, Korea, ³Division of Statistical Genomics, Washington University School of Medicine, 4444 Forest Park Boulevard, St. Louis, Missouri 63108, USA and ⁴Division of Biostatistics, Washington University School of Medicine, 660 South Euclid Avenue, St. Louis, Missouri 63110, USA

E-mail: Yong-Moon Park - mark@catholic.ac.kr; Michael A Province - mprovince@wustl.edu; Xiaoyi Gao - xgao23@wustl.edu; Mary Feitosa - mfeitosa@wustl.edu; Jun Wu - jwu@wustl.edu; Duanduan Ma - ma@wustl.edu; DC Rao - rao@wubios.wustl.edu; Aldi T Kraja* - aldi@wustl.edu

*Corresponding author

from Genetic Analysis Workshop 16
St Louis, MO, USA 17-20 September 2009

Published: 15 December 2009

BMC Proceedings 2009, 3(Suppl 7):S116 doi: 10.1186/1753-6561-3-S7-S116

This article is available from: <http://www.biomedcentral.com/1753-6561/3/S7/S116>

© 2009 Park et al; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

We investigated the association of metabolic syndrome (MetS) with a 500 k and a 50 k single-nucleotide polymorphism (SNP) gene chip in the Framingham Heart Study. We cross-sectionally evaluated the MetS longitudinal trends. Data analyzed were from the Offspring Cohort (four exams: first ($n = 2,441$), third ($n = 2,185$), fifth ($n = 2,308$), and seventh ($n = 2,328$)) and the Generation 3 Cohort (one exam: the first exam ($n = 3,997$)). The prevalence of MetS was determined using the National Cholesterol Education Program Adult Treatment Panel III diagnostic criteria, modified with a newly developed correction for medication use. The association test between an SNP and MetS was performed with a generalized estimating equations method under the additive genetic model. Multiple-testing corrections were also performed. The prevalence of MetS in the offspring cohort increased from one visit to the next, and reached the highest point by the seventh exam comparable with the prevalence reported for the general US population. The pattern of the MetS prevalence over time also reflected itself in the association tests, in which the highest significances were seen in the fifth and seventh exams. The association tests showed that SNPs within genes *PRDM16*, *CETP*, *PTHBI*, *PAPPA*, and *FBN3*, and also some SNPs not in genes were significant or close to significance at the genome-wide thresholds. These findings are important in terms of eventually identifying with the causal loci for MetS.

Background

Metabolic syndrome (MetS) is characterized by abdominal obesity, dyslipidemia, elevated blood pressure, insulin resistance, glucose intolerance, and possibly a prothrombotic and proinflammatory state [1]. MetS is a rising global public health problem because of its role in increasing the risk of cardiovascular disease and diabetes mellitus [2]. In addition to the lifestyle or environmental risk factors, much evidence has shown that common genetic variants predispose individuals to development of MetS [3]. Framingham Heart Study (FHS) is one of the best-known cohort studies of cardiovascular disease that has demonstrated an association between cardiovascular risk factors and cardiovascular disease [4]. In the current study we investigated the longitudinal trends in MetS association with a 500 k and 50 k single-nucleotide polymorphisms (SNPs) in the FHS datasets.

Methods

Sampled data and MetS definition

The FHS data comprised three generations of longitudinal measurements. The first dataset is the "Original Cohort", the second dataset is the "Offspring Cohort" and the third dataset is the "Generation 3 Cohort". We analyzed data from the Offspring Cohort and the Generation 3 Cohort. The data of Original Cohort were excluded due to missing variables related to MetS. Longitudinal analysis for the prevalence of MetS and its components was performed in each of the four exams (first ($n = 2,441$), third ($n = 2,185$), fifth ($n = 2,308$), and seventh ($n = 2,328$) exams) of the Offspring Cohort, and a cross-sectional analysis was performed in the first exam ($n = 3,997$) of the Generation 3 Cohort. The prevalence of MetS was identified using the National Cholesterol Education Program (NCEP) modified diagnostic criteria with a newly developed correction for medication use [1]. One exception was that the waist circumference was not available in the FHS data distributed through Genetic Analysis Workshop 16. Therefore, we substituted it with the body mass index (BMI) criteria for obesity. Previous publications have shown that BMI is highly correlated with the waist circumference [5]. An individual with a combination of any three or more of the following risk factors was classified as having MetS: BMI ≥ 30 kg/m²; triglyceride (TG) ≥ 150 mg/dl; high-density lipoprotein cholesterol (HDL) < 40 mg/dl in men and < 50 mg/dl in women; systolic blood pressure (SBP) or diastolic blood pressure (DBP) $\geq 130/85$ mm Hg or use of anti-hypertensive medications; and fasting blood glucose (GLUC) ≥ 110 mg/dl or use of anti-diabetic medications with the age of diagnosis with diabetes mellitus ≥ 40 years. The modified NCEP diagnostic criteria were applied to consider the participants medication use. Thus, TG and HDL for the subjects

treated with antihyperlipidemics were corrected to new values based on the following formula: TG/(1-15.2/100), HDLC/(1+6.1/100) (mg/dl). The SBP and DBP for subjects using antihypertensives were corrected to new values based on the following formula: SBP+14.8 and DBP+10.5 (mm Hg). These formulas were based on previous research, which represent corrections of a participant's medicated traits based on the mean values of many summarized treatment clinical trials [6-8].

Statistical analysis

The qualitative MetS variable, which took medication use into account, was used to test its association with both the genome-wide scan 500 k SNPs and the additional 50 k gene SNPs, for a total of approximately 550 k SNPs (GeneChip® Human Mapping 500 k Array Set and the 50 k Human Gene Focused Panel). The genotype data were recoded based on the additive model. A logistic regression of recoded genotype on MetS, based on the generalized estimating equation statistical model to account for the familial relationships among subjects within a pedigree, was performed using PROC GENMOD of SAS v 9.1.3 applying parallel computing with Platform SAS under Linux OS. In order to address multiple testing correction issues, we applied the *simpleM* method, which is an effective number of independent tests approach based on principal-component analysis that takes into account linkage disequilibrium (LD) information among SNPs [9]. The inferred effective number of independent tests for the 500 k chip was 281,502. Therefore, the genome-wide threshold for declaring significance for this data was a p -value $\leq 1.78 \times 10^{-7}$. For the 50 k chip, based on the same method the significance threshold was 1.29×10^{-6} . We considered these two chip results separately because they represent independent chips and also our analyses per type of chip were performed independent of each other.

Results

MetS: trends in prevalence

The mean (range) age of subjects at the first, third, fifth, and seventh exam in the Offspring Cohort was 33.6 (9-60), 46.2 (21-72), 53.1 (28-79), and 60.1 (35-85) years, respectively. The mean (range) age of subjects in the Generation 3 Cohort was 40.2 (19-72) years. The prevalence of MetS at the first, third, fifth, and seventh exam in the offspring cohort was 10.9%, 15%, 23.7%, and 23.6%, respectively. The prevalence of MetS at first exam in the Generation 3 Cohort was 14.2%. The prevalence of subjects having four and five components of MetS beyond their threshold increased with the number of exams, while the prevalence of subjects having three components beyond the threshold decreased on the seventh exam (Figure 1). Particularly, the prevalence of high BMI, high




Figure 1

Trends of MetS prevalence in the FHS. The prevalence of MetS in the Offspring Cohort tended to increase with the number of exams and reached the highest point on the fifth and seventh exam, comparable with the MetS prevalence of the general US population.

TG, and high BP increased with exams. The prevalence of high GLUC increased with the exams, except for the GLUC at first exam. The changes in HDL prevalence had no particular trend.

Association analysis

The association tests showed that a few SNPs within genes *PRDM16*, *CETP*, *PTHB1*, *PAPPA*, and *FBN3*, and also some SNPs not in genes, were significant or close to significance at the genome-wide thresholds for different exams in the Offspring Cohort and a few in the Generation 3 Cohort (Table 1). If the primary significant association MetS-SNP for a particular gene was from the 50 k chip, then we found that the 500 k chip for each gene reported had other SNPs that also were significant but did not reach the ultimate genome-wide thresholds. The same was observed when the primary significant association was from the 500 k. The 500 k significant SNPs findings were supported from other SNPs of the same gene on 50 k chip, but they did not pass the genome-wide thresholds. Due to space restriction, we do not report these findings.

Discussion

The longitudinal trends of MetS prevalence and of the association tests were evaluated by using the cross-sectional phenotypic measurements. The prevalence of MetS in the Offspring Cohort increased from one visit to the next, and reached the highest point by the seventh exam, comparable with reported MetS prevalence of the US population [2]. The prevalence of MetS was also high in the young people (Generation 3 Cohort), driven mostly by the increase in obesity. This pattern was also seen in the association results, where the highest significance values were found in analysis of the later exams (fifth and seventh). SNP rs17390167, part of the gene *PRDM16* on chromosome 1, was close to the significance threshold of the 50 k chip. This gene recently is reported to be the principal regulator of brown adipocyte tissue formation and function [10,11]. Another interesting finding was the association of rs11508026, a SNP on the *CETP* (cholesterol ester transfer protein) gene, with the MetS. This gene is already studied extensively for its inhibitors, because inhibition of *CETP* elevates the fraction of plasma

Table 1: SNPs associated significantly with the qualitative MetS corrected for medication use (4 exams of offspring cohort and 1 exam of generation 3 cohort data)

Marker	Chr	Position (bp)	Hugo	Role	Exam 21a			Exam 23			Exam 25			Exam 27			Exam 31			
					MAF (%)	Estimate	STD error	Estimate	STD error	Estimate	STD error	Estimate	STD error	Estimate	STD error	Estimate	STD error			
50 k																				
rs17390167	1	3084691	PRDM16	Intron	5.2	-0.01	0.02	4.9×10^{-1}	-0.06	0.03	1.8×10^{-2}	-0.14	0.03	3.8×10^{-6}	-0.04	0.03	1.4×10^{-1}	-0.01	0.02	4.6×10^{-1}
rs11508026	16	55556829	CETP	Intron	42.5	-0.02	0.01	1.1×10^{-1}	-0.04	0.01	2.0×10^{-4}	-0.04	0.01	2.7×10^{-3}	-0.07	0.01	7.2×10^{-7}	-0.01	0.01	1.6×10^{-1}
500 k																				
(in genes)																				
rs13241465	7	33473074	PTHBI	Intron	32.6	0.00	0.01	8.8×10^{-1}	0.02	0.01	1.8×10^{-1}	0.08	0.01	1.4×10^{-8}	0.03	0.01	1.3×10^{-2}	0.03	0.01	1.3×10^{-3}
rs4236337	7	33473431	PTHBI	Intron	32.4	0.00	0.01	9.2×10^{-1}	0.02	0.01	1.6×10^{-1}	0.08	0.01	2.8×10^{-8}	0.03	0.01	1.9×10^{-2}	0.03	0.01	2.4×10^{-3}
rs4509212	7	33473170	PTHBI	Intron	31.4	0.00	0.01	9.4×10^{-1}	0.02	0.01	7.2×10^{-2}	0.07	0.01	6.2×10^{-8}	0.03	0.01	1.5×10^{-2}	0.02	0.01	1.3×10^{-2}
rs2418441	9	118114658	PAPPA	Intron	1.0	-0.02	0.04	6.7×10^{-1}	0.01	0.05	8.4×10^{-1}	0.06	0.06	3.3×10^{-1}	-0.03	0.06	5.9×10^{-1}	-0.25	0.04	1.2×10^{-8}
rs10408896	19	8124201	FBN3	Promoter	0.1	0.03	0.14	8.3×10^{-1}	0.41	0.16	1.0×10^{-2}	0.12	0.19	5.4×10^{-1}	0.12	0.19	5.4×10^{-1}	0.49	0.09	8.8×10^{-8}
500 k																				
(not in genes)																				
rs318256	8				8.2	-0.01	0.02	4.6×10^{-1}	0.00	0.02	9.2×10^{-1}	0.02	0.02	4.4×10^{-1}	0.01	0.02	7.2×10^{-1}	0.08	0.01	1.5×10^{-7}
rs1431573	11	126981969			43.9	-0.01	0.01	5.5×10^{-1}	-0.06	0.01	1.4×10^{-7c}	-0.03	0.01	7.8×10^{-3}	-0.02	0.01	1.9×10^{-1}	0.00	0.01	9.4×10^{-1}
rs17037068	12	104333075			1.7	0.10	0.04	3.7×10^{-3b}	0.27	0.04	5.3×10^{-10}	0.14	0.05	9.1×10^{-3}	0.14	0.05	4.2×10^{-3}	-0.03	0.03	2.9×10^{-1}
rs12437159	14	45882211			44.6	-0.02	0.01	6.7×10^{-3}	-0.04	0.01	6.2×10^{-5}	-0.06	0.01	4.3×10^{-7}	-0.08	0.01	1.0×10^{-9}	0.00	0.01	9.1×10^{-1}
rs12147964	14	45888772			42.5	-0.02	0.01	9.8×10^{-3}	-0.04	0.01	2.7×10^{-4}	-0.07	0.01	2.7×10^{-8}	-0.08	0.01	7.7×10^{-11}	0.00	0.01	8.6×10^{-1}
rs2899976	14	45893237			41.8	-0.02	0.01	1.3×10^{-2}	-0.04	0.01	9.5×10^{-4}	-0.07	0.01	2.0×10^{-7}	-0.08	0.01	2.3×10^{-10}	0.00	0.01	9.1×10^{-1}
rs11625735	14	45895075			41.8	-0.02	0.01	1.4×10^{-2}	-0.04	0.01	4.7×10^{-4}	-0.07	0.01	9.4×10^{-8}	-0.08	0.01	1.8×10^{-10}	0.00	0.01	9.0×10^{-1}
rs2010338	14	45895631			41.8	-0.02	0.01	8.9×10^{-3}	-0.04	0.01	3.4×10^{-4}	-0.07	0.01	4.2×10^{-8}	-0.08	0.01	1.5×10^{-10}	0.00	0.01	7.4×10^{-1}
rs2415974	14	45906862			42.0	-0.02	0.01	7.1×10^{-3}	-0.04	0.01	3.0×10^{-4}	-0.07	0.01	1.2×10^{-7}	-0.08	0.01	6.8×10^{-11}	0.00	0.01	7.8×10^{-1}
rs8009432	14	45917176			41.7	-0.02	0.01	1.4×10^{-2}	-0.04	0.01	4.4×10^{-4}	-0.07	0.01	1.2×10^{-7}	-0.08	0.01	3.1×10^{-10}	0.00	0.01	8.6×10^{-1}
rs10498398	14	45937239			41.1	0.02	0.01	9.8×10^{-3}	0.04	0.01	1.0×10^{-3}	0.07	0.01	8.6×10^{-8}	0.08	0.01	1.7×10^{-9}	0.00	0.01	8.4×10^{-1}
rs8019899	14	45952022			41.3	0.02	0.01	5.6×10^{-3}	0.04	0.01	1.1×10^{-3}	0.07	0.01	1.2×10^{-7}	0.07	0.01	3.0×10^{-9}	0.00	0.01	8.1×10^{-1}

^aExam 21, 2 is the Offspring Cohort, and 1 is the first exam.

^b**Bold**, *p*-values support a significance trend, but do not reach the threshold.

^c*Italic*, *p*-values that pass the significance threshold and support genome-wide significance.

cholesterol associated with HDL [12]. SNPs rs13241465, rs4236337, and rs4509212 are part of the *PTHB1* gene. This gene is known also as *BBS9* gene and recognized as one of the recent Bardet-Biedl syndrome genes. Such gene polymorphisms are hypothesized to contribute to human obesity and diabetes mellitus [13].

In the Generation 3 Cohort first exam, the prevalence of MetS was moderate and comparable with those of the first and third exams of the Offspring Cohort. As a result, only a few associations were highly significant. The SNPs rs2418441 (*PAPPA*) and rs10408896 (*FBN3*) were highly significant, but at the same time one must be cautious when considering them as candidate genes for MetS, because of the rare minor allele frequency. Despite the rarity of minor allele frequency for the above two SNPs, *PAPPA* gene, a metalloproteinase, is reported to regulate the human atherosclerotic plaque [14]; *FBN3* (Fibrillin 3) is reported to contribute in the polycystic ovary syndrome, which is associated with obesity [15]. Several other SNPs with *p*-values close to significance from the same genes mentioned in the Results and Discussion were present in the FHS data. For space reasons we did not report any of them. Finally, we would like to draw readers attention to a number of SNPs located on chromosome 14 (rs12437159, rs12147964, rs2899976, rs11625735, rs2010338, rs2415974, rs8009432, rs10498398, and rs8019899), which although not located on genes, still were associated highly with MetS.

Conclusion

These findings are important for identifying the role of the genes reported to be associated with MetS. As such, they need to be further investigated in conjunction with the quantitative risk factors of MetS.

List of abbreviations used

BMI: Body mass index; DBP: Diastolic blood pressure; FHS: Framingham Heart Study; GLUC: Fasting blood glucose; HDLC: High-density lipoprotein cholesterol; LD: Linkage disequilibrium; MetS: Metabolic syndrome; NCEP: National Cholesterol Education Program; SBP: Systolic blood pressure; SNP: Single-nucleotide polymorphism; TG: Triglyceride.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Y-MP, ATK, MAP, and DCR conceived of the idea of this analysis. Y-MP and ATK performed the analysis and wrote the manuscript. MAP, DCR, MF, XG, JW, and DM

contributed ideas to improve the manuscript. XG performed the thresholds' *p*-value calculations.

Acknowledgements

The Genetic Analysis Workshops are supported by NIH grant R01 GM031575 from the National Institute of General Medical Sciences.

This article has been published as part of *BMC Proceedings* Volume 3 Supplement 7, 2009: Genetic Analysis Workshop 16. The full contents of the supplement are available online at <http://www.biomedcentral.com/1753-6561/3?issue=S7>.

References

- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults: **Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III)**. *JAMA* 2001, **285**:2486–2497.
- Eckel RH, Grundy SM and Zimmet PZ: **The metabolic syndrome**. *Lancet* 2005, **365**:1415–1428.
- Pollex RL and Hegele RA: **Genetic determinants of the metabolic syndrome**. *Nat Clin Pract Cardiovasc Med* 2006, **3**:482–489.
- Kannel WB: **CHD risk factors: a Framingham study update**. *Hosp Pract* 1990, **25**:119–127.
- Kraja AT, Hunt SC, Pankow JS, Myers RH, Heiss G, Lewis CE, Rao DC and Province MA: **An evaluation of the metabolic syndrome in the HyperGEN study**. *Nutr Metab (Lond)* 2005, **2**:17.
- Kraja AT, Borecki IB, North K, Tang W, Myers RH, Hopkins PN, Arnett DK, Corbett J, Adelman A and Province MA: **Longitudinal and age trends of metabolic syndrome and its risk factors: the Family Heart Study**. *Nutr Metab (Lond)* 2006, **3**:41.
- Wu J, Kraja AT, Oberman A, Lewis CE, Ellison RC, Arnett DK, Heiss G, Lalouel JM, Turner ST, Hunt SC, Province MA and Rao DC: **A summary of antihypertensive medication effects on measured blood pressure**. *Am J Hypertens* 2005, **18**:934–941.
- Wu J, Province MA, Coon H, Hunt SC, Eckfeldt JH, Arnett DK, Heiss G, Lewis CE, Ellison RC, Rao DC, Rice T and Kraja AT: **An investigation of the effects of lipid-lowering medications: genome-wide linkage analysis of lipids in the HyperGEN study**. *BMC Genet* 2007, **8**:60.
- Gao X, Starmer J and Martin ER: **A multiple testing correction method for genetic association studies using correlated single nucleotide polymorphisms**. *Genet Epidemiol* 2008, **32**:361–369.
- Farmer SR: **Molecular determinants of brown adipocyte formation and function**. *Genes Dev* 2008, **22**:1269–1275.
- Kajimura S, Seale P, Tomaru T, Erdjument Bromage H, Cooper MP, Ruas JL, Chin S, Tempst P, Lazar MA and Spiegelman BM: **Regulation of the brown and white fat gene programs through a PRDM16/CtBP transcriptional complex**. *Genes Dev* 2008, **22**:1397–1409.
- Cunningham D, Lin W, Hoth LR, Danley DE, Ruggeri RB, Geoghegan KF, Chrnyk BA and Boyd JG: **Biophysical and biochemical approach to locating an inhibitor binding site on cholesteryl ester transfer protein**. *Bioconjug Chem* 2008, **19**:1604–1613.
- Sun G: **Application of DNA microarrays in the study of human obesity and type 2 diabetes**. *OMICs* 2007, **11**:25–40.
- Conover CA, Harrington SC and Bale LK: **Differential regulation of pregnancy associated plasma protein-A in human coronary artery endothelial cells and smooth muscle cells**. *Growth Horm IGF Res* 2008, **18**:213–220.
- Urbanek M, Sam S, Legro RS and Dunaif A: **Identification of a polycystic ovary syndrome susceptibility variant in fibrillin-3 and association with a metabolic phenotype**. *J Clin Endocrinol Metab* 2007, **92**:4191–4198.