

1-5-2004

Amyloid-beta peptide induces oligodendrocyte death by activating the neutral sphingomyelinase-ceramide pathway

Jiunn-Tay Lee
National Defense Medical Center, Republic of China

Jan Xu
Washington University School of Medicine in St. Louis

Jin-Moo Lee
Washington University School of Medicine in St. Louis

Grace Ku
Washington University School of Medicine in St. Louis

Xianlin Han
Washington University School of Medicine in St. Louis

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs



Part of the [Medicine and Health Sciences Commons](#)

Please let us know how this document benefits you.

Recommended Citation

Lee, Jiunn-Tay; Xu, Jan; Lee, Jin-Moo; Ku, Grace; Han, Xianlin; Yang, Ding-I; Chen, Shawei; and Hsu, Chung Y., "Amyloid-beta peptide induces oligodendrocyte death by activating the neutral sphingomyelinase-ceramide pathway." *The Journal of Cell Biology*. 164, 1. 123-131. (2004).
https://digitalcommons.wustl.edu/open_access_pubs/606

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Jiunn-Tay Lee, Jan Xu, Jin-Moo Lee, Grace Ku, Xianlin Han, Ding-I Yang, Shawei Chen, and Chung Y. Hsu

Amyloid- β peptide induces oligodendrocyte death by activating the neutral sphingomyelinase–ceramide pathway

Giunn-Tay Lee,³ Jan Xu,¹ Jin-Moo Lee,¹ Grace Ku,¹ Xianlin Han,² Ding-I Yang,¹ Shawei Chen,¹ and Chung Y. Hsu^{1,4}

¹Center for the Study of Nervous System Injury, Department of Neurology, and ²Division of Bioorganic Chemistry and Molecular Pharmacology, Department of Internal Medicine, Washington University School of Medicine, St. Louis, MO 63110

³Department of Neurology, Tri-Service General Hospital, National Defense Medical Center, Taipei 114, Taiwan, Republic of China

⁴Taipei Medical University, Taipei 110, Taiwan, Republic of China

Amyloid- β peptide (A β) accumulation in senile plaques, a pathological hallmark of Alzheimer's disease (AD), has been implicated in neuronal degeneration. We have recently demonstrated that A β induced oligodendrocyte (OLG) apoptosis, suggesting a role in white matter pathology in AD. Here, we explore the molecular mechanisms involved in A β -induced OLG death, examining the potential role of ceramide, a known apoptogenic mediator. Both A β and ceramide induced OLG death. In addition, A β activated neutral sphingomyelinase (nSMase), but not acidic sphingomyelinase, resulting in increased ceramide generation. Blocking ceramide degradation with

N-oleoyl-ethanolamine exacerbated A β cytotoxicity; and addition of bacterial sphingomyelinase (mimicking cellular nSMase activity) induced OLG death. Furthermore, nSMase inhibition by 3-*O*-methyl-sphingomyelin or by gene knockdown using antisense oligonucleotides attenuated A β -induced OLG death. Glutathione (GSH) precursors inhibited A β activation of nSMase and prevented OLG death, whereas GSH depletors increased nSMase activity and A β -induced death. These results suggest that A β induces OLG death by activating the nSMase–ceramide cascade via an oxidative mechanism.

Introduction

The amyloid- β peptide (A β), a 39–43–amino acid cleavage product of amyloid precursor protein (Estus et al., 1992; Haass et al., 1992), has been implicated as the primary neurotoxic factor in Alzheimer's disease (AD) pathogenesis (Yankner et al., 1989). In vitro, A β is toxic to neurons (Yankner et al., 1989; Behl et al., 1994; Yu et al., 1998), endothelial cells (Thomas et al., 1996; Huang et al., 1998), astrocytes (Brera et al., 2000), vascular smooth muscle cells (Kawai et al., 1993; Davis-Salinas et al., 1995), and oligodendrocytes (OLGs; Xu et al., 2001); and A β depositions in senile plaques are postulated to cause neuronal and vascular degeneration in AD brains (Masters et al., 1985; Yankner et al., 1989; Thomas et al., 1996). Although A β -mediated cell

death demonstrates morphological, biochemical, and molecular features of apoptosis, the molecular mechanism underlying A β cytotoxicity remains largely undefined but may involve oxidative stress (Behl et al., 1994; Schapira, 1996). NF- κ B and AP-1, redox-sensitive transcription factors, are activated in A β -treated OLGs, and *N*-acetylcysteine (NAC), an antioxidant, prevents A β -mediated OLG apoptosis (Xu et al., 2001). OLGs are susceptible to oxidative stress because they have low levels of reduced glutathione (GSH) and high concentrations of iron, resulting in a compromised ability to scavenge peroxides (Thorburne and Juurlink, 1996; Back et al., 1998; Juurlink et al., 1998).

J.-T. Lee and J. Xu contributed equally to this paper.

Address correspondence to Chung Y. Hsu, Center for the Study of Nervous System Injury, Dept. of Neurology, Washington University School of Medicine, 660 S. Euclid Ave., Box 8111, St. Louis, MO 63110. Tel.: (314) 362-3304. Fax: (314) 362-9462. email: hsuc@neuro.wustl.edu

Key words: Alzheimer's disease; apoptosis; cell death; oxidative stress; white matter

Abbreviations used in this paper: 3-OMe-SM, 3-*O*-methyl-sphingomyelin; A β , amyloid- β peptide; AD, Alzheimer's disease; aSMase, acidic sphingomyelinase; bSMase, bacterial sphingomyelinase; BSO, buthionine sulfoximine; DEM, diethyl maleate; ESI/MS, electrospray ionization/mass spectrometry; GSH, glutathione; LDH, lactate dehydrogenase; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; NAC, *N*-acetylcysteine; nSMase, neutral sphingomyelinase; NOE, *N*-oleoyl-ethanolamine; OLG, oligodendrocyte; PLP, proteolipid protein.




Figure 1. Characterization of OLG cultures. Immunofluorescent (A, C, E, and G) and corresponding phase-contrast (B, D, F, and H) photomicrographs of neurosphere-derived differentiated OLGs in culture. Cells demonstrate characteristic morphology of OLGs, plump cell body with multiple branching processes. Virtually all cells in culture immunostained for the OLG-specific antigens, CNP (A), Rip (C), GalC (E), and PLP (G). Bar, 30 μ m.

Ceramide, a lipid second messenger that increases the cellular oxidative state, has been implicated in several apoptosis paradigms including trophic factor withdrawal and exposure to proinflammatory molecules (Coroneos et al., 1995; Kyriakis and Avruch, 1996; Kolesnick and Kronke, 1998). Cellular ceramide synthesis increases in response to stress or death signals (Haimovitz-Friedman et al., 1994; Tepper et al., 1995; Verheij et al., 1996). One pathway of ceramide formation involves sphingomyelin hydrolysis by either neutral sphingomyelinase (nSMase) or acidic sphingomyelinase (aSMase; Testi, 1996); both enzymes are involved in several cell death paradigms (Kolesnick and Kronke, 1998; Levade and Jaffrezou, 1999). Another pathway involves ceramide synthase-catalyzed de novo ceramide synthesis (Bose et al., 1995; Spiegel and Merrill, 1996; Xu et al., 1998).

$A\beta$ and ceramide share cell death signaling characteristics. $A\beta$ -induced apoptosis involves TNF- α , p75 neurotrophin receptor, and Fas ligand (Blasko et al., 1997; de la Monte et al., 1997; Yaar et al., 1997), which are cell surface receptors that relay death signals through the sphingomyelin-ceramide pathway (Dobrowsky et al., 1995; Hannun, 1996). Moreover, both $A\beta$ (Kaneko et al., 1995; Askanas et al., 1996; Bruce-Keller et al., 1998; Xu et al., 2001) and ceramide (Garcia-Ruiz et al., 1997; Singh et al.,




Figure 2. $A\beta$ 25-35 and C2-ceramide induced OLG death. (A) OLGs treated with 10–100 μ M $A\beta$ 25-35 for 48 h. (B) OLGs exposed to 10–100 μ M C2-ceramide for 24 h. The assessment of cell survival based on MTT assay or cell death based on LDH assay showed a dose-dependent cytotoxic effect of $A\beta$ 25-35 and C2-ceramide. Data shown are representative of three separate experiments of triplicate samples with similar results; error bars represent SD.

1998) cause mitochondrial dysfunction and induce oxidative stress. In vitro, OLG death induced by $A\beta$ (Xu et al., 2001) or ceramide (Larocca et al., 1997; Singh et al., 1998; Scurlock and Dawson, 1999) share similar apoptotic characteristics. Lower sphingomyelin levels and higher ceramide levels in AD brains have been reported (Soderberg et al., 1992), thereby implying that increased sphingomyelin degradation and ceramide accumulation contribute to AD pathogenesis. We have previously shown that $A\beta$ induced OLG death with characteristic features of apoptosis (Xu et al., 2001). In this paper, we demonstrate that ceramide mediates $A\beta$ -induced OLG death by activating the nSMase-ceramide cascade.

Results


$A\beta$ and C2-ceramide are cytotoxic to OLGs

We have previously shown that $A\beta$ induced apoptosis in primary OLG cultures derived from neonatal rat brains, characterized by nuclear and cytoskeletal disintegration, DNA fragmentation, and mitochondrial dysfunction (Xu et al., 2001). In this paper, we used neurosphere-derived differentiated OLGs because of their ease of preparation compared with primary OLG isolation. The neurosphere-derived differentiated OLGs exhibited characteristic OLG morphology, which is composed of large cell bodies with multiple branching processes (Fig. 1, A–H). Immunocytochemical analysis revealed that virtually all cells in culture expressed four OLG-specific surface markers including cyclic nucleotide 3'-phosphodiesterase (Fig. 1 A, CNP), Rip (Fig. 1 C), galactocerebroside (Fig. 1 E, GalC), and PLP (myelin proteolipid protein; Fig. 1 G). In addition, differentiated OLGs showed similar characteristic electrophysiological membrane potentials (unpublished data) as reported in previous studies (McDonald et al., 1999).

As demonstrated previously, $A\beta$ 1-40 and $A\beta$ 25-35 peptides have equal potencies for inducing death in primary cultures of OLGs (Xu et al., 2001); thus, most of our studies were conducted using $A\beta$ 25-35 with key experiments confirmed with $A\beta$ 1-40. $A\beta$ 25-35 treatment for 48 h caused OLG death in a concentration-dependent manner with an

Figure 3. Time-dependent cell death induced by A β 25-35 and C2-ceramide.

(A) Cell survival as measured by MTT assay. (B) Cell death as determined by LDH assay. OLGs were treated with 20 μ M A β 25-35 or 30 μ M C2-ceramide for the time periods indicated. The cells and culture medium were used for the MTT or LDH assay, respectively. Note the faster pace of ceramide in inducing OLG death, 50% OLG death was noted 24 and 48 h after ceramide and A β treatment, respectively. Data shown are representative of three separate experiments of triplicate samples with similar results; error bars represent SD.



approximate EC₅₀ = 20 μ M (Fig. 2 A). C2-ceramide, a cell-permeable ceramide analogue, also induced dose-dependent OLG death with an approximate EC₅₀ = 30 μ M (Fig. 2 B). C2-dihydroceramide, a cell-impermeable and inactive form of ceramide, was not cytotoxic at concentrations up to 100 μ M (unpublished data). The toxic effect of ceramide was apparent at an earlier time point than that of A β (Fig. 3, A and B), raising the possibility that ceramide may be a mediator of A β -induced cell death.

Ceramide in A β 25-35-induced OLG death

To support the contention that ceramide is involved in A β -induced OLG death, TLC and electrospray ionization/mass spectrometry (ESI/MS) were used to measure ceramide content in OLGs with and without A β treatment. A β 25-35 treatment resulted in a fivefold increase in ceramide synthesis as determined by TLC (Fig. 4 A). An increase in cellular ceramide content was confirmed by ESI/MS, which demonstrated peak ceramide content 10 h after A β treatment (Fig. 4 B). *N*-oleoyl-ethanolamine (NOE), a specific ceramidase inhibitor that prevents the degradation of cellular ceramide at subtoxic doses (Sugita et al., 1975; Pahan et al., 1998), augmented A β 25-35-induced OLG death (Fig. 4 C). Together, these results suggest a strong correlation between increases in cellular ceramide levels and A β -induced OLG death.

nSMase activation in A β 25-35-induced OLG death

Ceramide is derived from sphingomyelin hydrolysis catalyzed by nSMase or aSMase (Hannun, 1996). A β 25-35 treatment increased nSMase activity in OLGs as early as 2.5 min after exposure and reached maximal levels at \sim 16 h (Fig. 5 A). A β 1-40 treatment also increased OLG nSMase activity in a similar manner (unpublished data). In contrast, A β 25-35 treatment did not alter aSMase activity (Fig. 5 A). The addition of recombinant bacterial sphingomyelinase (bSMase), an exogenous source of sphingomyelinase that mimics nSMase action by degrading membrane sphingomyelin to increase cellular ceramide levels (Okazaki et al., 1989; Jarvis et al., 1994; Zhang et al., 1997; Tonnetti et al., 1999), caused OLG death (Fig. 5 B). Furthermore, the




Figure 4. A β 25-35-induced ceramide production and enhancement of A β cytotoxicity by a ceramidase inhibitor. (A) An increase in cellular ceramide synthesis in OLGs prelabeled with [³H]palmitate and treated with 10 μ M A β 25-35 for 24 h. (B) Quantitative determination of endogenous ceramide concentration by ESI/MS at various time points after A β 25-35 treatment. Note the elevated ceramide content at 5, 10, and 16 h after A β 25-35 exposure. (C) Effect of NOE, a ceramidase inhibitor, on A β -induced OLG death. OLGs were treated with 10 μ M A β 25-35 plus 1 μ M NOE for 24 h. The culture medium was collected for the LDH assay. Data shown are mean \pm SD of three separate experiments in triplicate. *, Significant difference from control; **, significant difference from A β treatment; P < 0.05.

nSMase inhibitors, 3-O-methyl-sphingomyelin (3-OME-SM; Lister et al., 1995) and NAC (Liu et al., 1998a; Yoshimura et al., 1999), were effective in protecting OLGs against A β cytotoxicity (Fig. 5, C and D). Confirming the effects of the nSMase inhibitors, a marked reduction in nSMase enzymatic activity was observed in OLGs treated with 3-OME-SM or NAC (Fig. 6 A). Moreover, significant decreases in endogenous ceramide content and increases in sphingomyelin levels were detected in OLGs treated with the inhibitors (Fig. 6, B and C). To further confirm the contribution of nSMase in A β -mediated cell death, sense and antisense oligonucleotides specific for nSMase were generated. Antisense oligonucleotides reduced nSMase activity (Fig. 7 A), reduced ceramide content in cell lysates (Fig. 7 B), and attenuated A β -induced cell death (Fig. 7 C), but sense oligonucleotides had no effect on nSMase activity, ceramide content, or cell survival.




Figure 5. Spingomyelinase in A β -induced OLG death. (A) nSMase and aSMase activity in OLGs treated with 10 μ M A β 25-35 for various time periods (from 2.5 min to 36 h). (B) A β or bSMase induced OLG death as determined by LDH assay. OLGs were exposed to 10 μ M A β 25-35 or bSMase (1 U/ml) for 24 h. (C) LDH and (D) trypan blue assays on the effects of nSMase inhibitors on A β -mediated OLG death. OLGs were treated with 25 mM NAC or 1 μ M 3-OMe-SM for 2 h before 10 μ M A β 25-35 treatment for 24 h. Data shown are representative of three separate experiments of triplicate samples with similar results; error bars represent SD. *, Significant difference from control; **, significant difference from A β treatment; P < 0.05.

It has been shown that A β depletes GSH in cortical neurons in vitro (Muller et al., 1997), which correlates well with the ability of A β to heighten cellular oxidative stress (Cafe et al., 1996). GSH depletion has been shown to activate nSMase activity (Liu and Hannun, 1997; Liu et al., 1998a,b). The fact that NAC, a GSH precursor, inhibited A β 25-35 activation of nSMase and prevented A β -mediated OLG death suggests that A β 25-35 depletion of cellular GSH is involved in the A β -nSMase-ceramide OLG death signaling cascade. Buthionine sulfoximine (BSO) and diethyl maleate (DEM) have been shown to deplete GSH in various cell types (Anderson and Meister, 1983; Masukawa et al., 1983; Szekely and Lobreau, 1987). Both BSO and DEM selectively increased nSMase activity (Fig. 8 A), increased ceramide levels (Fig. 8 B), decreased cellular GSH levels (Fig. 8 C), and were cytotoxic to OLGs (Fig. 8 D).

Because A β 25-35 did not activate aSMase, it is unlikely that aSMase is involved in A β 25-35-induced OLG death. Selective aSMase inhibitors such as desipramine and chlorpromazine (Albouz et al., 1986) were ineffective in protecting OLGs from A β 25-35-induced death (unpublished data). Furthermore, fumonisins B₂, a ceramide synthase inhibitor, did not block A β 25-35 cytotoxicity in this cell death paradigm (unpublished data). Together, these results suggest that ceramide generation catalyzed by nSMase, but not aSMase or ceramide synthase, mediates the A β 25-35 death pathway in OLGs.




Figure 6. Effect of nSMase inhibitors on sphingomyelinase activity and ceramide/sphingomyelin content in A β 25-35-treated OLGs. nSMase and aSMase activity (A) and cellular ceramide (B) and sphingomyelin (C) content in OLGs treated with NAC or 3-OMe-SM nSMase inhibitors. OLGs were treated with 1 μ M 3-OMe-SM or 25 mM NAC for 2 h before 10 μ M A β 25-35 exposure for 16 h. Representative TLC data are shown from one of three experiments in the top panels in B and C. A and the bottom panels in B and C show composite results from three separate experiments; error bars represent SD. *, Significant difference from control; **, significant difference from A β treatment; P < 0.05.

Discussion

Several lines of evidence support the contention that ceramide mediates, at least in part, A β -induced OLG death. Both A β and C2-ceramide (but not the biologically inactive dihydroceramide) caused OLG death in a time-dependent manner. A β treatment increased ceramide formation in OLGs. In addition, increasing cellular ceramide release from sphingomyelin by exogenous bSMase, which mimics cellular nSMase action, also induced OLG death. Inhibition of ceramide degradation by NOE, a ceramidase inhibitor, enhanced A β cytotoxicity in OLGs.

Results from this study also support the contention that A β cytotoxicity is mediated via activation of nSMase leading to increased cellular ceramide generation. A β 25-35 and A β 1-40 activated nSMase, but not aSMase, in OLGs. Addition-




Figure 7. Treatment of OLGs with nSMase antisense oligonucleotides. OLGs were treated with sense or antisense oligonucleotides specific for nSMase 4 h before 20 μ M A β addition. After 24 h, the cells were harvested and assayed for nSMase activity (A), ceramide content (B), and MTT (C). The graph depicts a representative of three separate experiments with similar results; error bars represent SD. *, Significant differences compared with A β treatment alone; $P < 0.05$.

ally, nSMase inhibitors such as 3-OMe-SM and NAC (also an antioxidant) prevented A β 25-35-induced nSMase activity, which resulted in decreased ceramide synthesis from sphingomyelin and protected OLGs from A β 25-35 cytotoxicity. Antisense oligonucleotides specific for nSMase also attenuated A β -induced OLG cell death, further implicating nSMase as a mediator. Chemical agents such as BSO and DEM that deplete cellular GSH content also activated nSMase in OLGs and caused cell death. The specific role of nSMase in A β -induced OLG death is supported by the finding that pharmacological inhibition of nSMase, but not aSMase or ceramide synthase, prevented A β 25-35-induced OLG death.

The exact mechanism underlying A β -mediated nSMase activation remains to be elucidated but may involve changes in the cellular redox state and/or GSH metabolism (Sawai and Hannun, 1999); GSH is the most abundant thiol-containing compound in living cells. nSMase enzymatic activity is directly regulated by cellular GSH content (Liu and Hannun, 1997; Liu et al., 1998a,b). A β has been shown to deplete GSH in cultured cortical neurons (Muller et al., 1997), and depletion of cellular GSH stores by oxidative stress has been proposed as a prime mechanism underlying the A β cytotoxic action (Muller et al., 1997; Pereira et al., 1999). Thus, it is plausible that a decrease in GSH level subsequent to A β exposure may activate nSMase in OLGs. NAC, a GSH precursor,




Figure 8. Effects of GSH depleting agents compared with A β 25-35 on sphingomyelinase activity, cellular GSH content, and cell survival. OLGs were treated with 10 μ M A β 25-35, 100 μ M BSO, or 500 μ M DEM for 16 h before measurement of sphingomyelinase activity (A), ceramide synthesis (B), or GSH content (C), or 24 h before MTT assay (D). Data shown in A, C, and D are representative of three separate experiments of triplicate samples and, in B, of two separate experiments with similar findings; error bars represent SD. *, Significant difference from control; $P < 0.05$.

inhibited A β 25-35 activation of nSMase and protected OLGs against A β -induced death, whereas depletion of cellular GSH stores by BSO or DEM resulted in selective activation of nSMase and OLG death. Although the agents used to manipulate GSH levels may be relatively nonspecific, these findings raise the possibility that the activation of nSMase by A β may involve the depletion of cellular GSH content.

Oxidative stress plays a prominent role in A β -mediated neuronal and OLG death (Behl et al., 1994; Behl, 1999; Markesbery, 1999; Xu et al., 2001). Brain tissue is especially sensitive to oxidative injury because of its higher metabolic rate driven by glucose, lower concentrations of protective antioxidants, and higher levels of polyunsaturated fatty acids that are susceptible to lipid peroxidation (Behl and Sagara, 1997; Behl, 1999; Markesbery, 1999). Although A β -mediated oxidative stress induces mtDNA damage (Bozner et al., 1997; Xu et al., 2001) and activates selected transcription factors including NF- κ B and AP-1 (Abate et al., 1990; Schreck et al., 1991; Pinkus et al., 1996; Xu et al., 2001), the mechanism by which A β induces oxidative stress in the AD brain remains unknown. Ceramide has emerged as a potent second messenger in oxidative stress-induced apoptosis (Hannun and Luberto, 2000). Hydrogen peroxide (Goldkorn et al., 1998), 1- β -D-arabinofuranosylcytosine (Bradshaw et al., 1996), daunorubicin (Jaffrezou et al., 1996), TNF- α (Bezombes et al., 1998), γ -rays (Bruno et al., 1998), hypoxia (Yoshimura et al., 1998), CD40 activation (Segui et al., 1999), and sindbis virus infection (Jan et al., 2000) are among the agents that can mediate cell death via nSMase activation, thus emphasizing the central role of the nSMase-ceramide cascade. Results shown here provide a unique signaling pathway from cell surface A β engagement, induction

of oxidative stress, and activation of the nSMase–ceramide cascade culminating in OLG death.

In summary, this work demonstrates a novel mechanism for A β -induced OLG death. These results reveal a causal relationship between A β exposure and the activation of the nSMase–ceramide pathway, which is likely to involve heightened oxidative stress after depletion of cellular GSH stores. In addition, we have evidence that activation of the nSMase–ceramide cascade may also contribute to A β -induced death of cortical neurons and cerebral endothelial cells (unpublished data), thereby suggesting that this cascade may be operating in many cell types other than OLGs. A β →nSMase→ceramide cascade represents a novel signaling pathway that contributes at least in part to A β cytotoxicity to various types of brain cells. Identification of this pathway may lead to the development of more effective therapeutic strategies aimed at preventing A β -induced cell death. For instance, blockade of the A β -activated death signaling process can be achieved by pharmacological modulation of nSMase activity as demonstrated in this work.

Materials and methods

Reagents and cell culture

All chemicals were purchased from Sigma-Aldrich, and all cell culture reagents were purchased from Invitrogen unless otherwise specified. B104 cells (a gift from David Schubert, Salk Institute for Biological Studies, La Jolla, CA) were cultured in DME supplemented with 10% FBS, 100 U/ml penicillin, and 100 μ g/ml streptomycin.

OLG culture

Neurospheres were cultured using the methods of Zhang et al. (1999) with modifications. In brief, embryonic rat brains (E14–16 d) were dissected, homogenized gently in DME/Ham's F12, and centrifuged at 350 g for 5 min. The pellet was digested with 0.05% trypsin in 1.5 ml of 0.53 M EDTA for 30 min at 37°C, followed by the addition of 1.5 ml DME/Ham's F12 with 20% FBS, and filtered through 10- μ m nylon mesh. The filtrate was centrifuged at 350 g for 5 min, and the pellet was washed twice with DME/Ham's F12. Dissociated cells were layered on a pre-equilibrated Percoll gradient (formed by centrifuging 50% Percoll and 50% DME/Ham's F12 at 23,500 g for 1 h at 4°C) and centrifuged at 3,500 g for 15 min. The fraction containing glial progenitors banding between myelin and blood cell layers was recovered and washed twice with DME/Ham's F12 followed by another wash with neurosphere culture medium (DME/Ham's F12/Hepes, N1 supplement, 25 μ g/ml insulin, 130 ng/ml progesterone, 20 ng/ml of basic FGF, and 20 ng/ml EGF). The cell pellet was resuspended in 20 ml of neurosphere culture medium and seeded in 75-mm culture flasks. After 24 h, when neurospheres had formed, 5 ml of fresh medium was added to each culture every other day for 7 d, and then the neurosphere cultures were split (1:2). The neurospheres were dissociated gently 10 times with a syringe and a 25-gauge needle and centrifuged at 350 g. The resulting cell pellets were treated with 0.05% trypsin/0.53 mM EDTA and centrifuged at 350 g for 10 min. The cells were resuspended in progenitor medium (69% DME/Ham's F12/Hepes containing N1 supplement, 10 μ g/ml insulin, 20 nM progesterone, 30% conditioned medium from B104 cells, and 1% FBS) and plated on 100-mm culture dishes precoated with poly-L-ornithine. For differentiated OLG cultures, progenitor cells were detached with trypsin/EDTA and cultured on poly-L-ornithine-coated plates or coverslips in mature OLG medium (DME/Ham's F12; N1 supplement, 20 μ g/ml biotin; 20 μ g/ml triiodo-L-thyronine, T3, and 1% FBS).

Immunocytochemistry

Differentiated OLG cultures were fixed in 4% PFA, washed in PBS, and blocked with 5% goat serum. Fixed cells were incubated with the primary antibody overnight at 4°C. The following primary antibodies were used: anti-mouse CNP, anti-mouse Rip, anti-mouse PLP (Chemicon), and anti-mouse GalC (Cedarlane Laboratories) at a concentration of 1:100. The secondary antibody, anti-mouse IgG conjugated to FITC (Vector Laboratories), was added for 1 h at RT. The cells were washed in PBS and visual-

ized using a confocal microscope (model LSM 5 Pascal; Carl Zeiss MicroImaging, Inc.) equipped with a CCD camera (model Axiocam HR; Carl Zeiss MicroImaging, Inc.). Images were collected and processed using Adobe Photoshop software.

Sphingomyelinase assay

The cells were washed twice with PBS, pH 7.4, and lysed in 0.2% Triton X-100 for 10 min at 4°C. The lysates were sonicated for 30 s in ice-cold bath, and protein concentrations were determined by Lowry assay (Lowry et al., 1951). A sphingomyelinase substrate, [methyl-¹⁴C]sphingomyelin (55 mCi/mmol; Amersham Biosciences), was evaporated to dryness and resuspended in either 25 μ l of nSMase assay buffer (40 mM Hepes, 5 mM MgCl₂, and 0.2% Triton X-100, pH 7.4) or aSMase assay buffer (250 mM sodium acetate and 0.2% Triton X-100, pH 5.2) and sonicated to form micelles on ice until use. Each reaction containing 25 μ l of cell lysate protein (1 mg/ml) and 25 μ l [methyl-¹⁴C]sphingomyelin (0.23 nmol) in nSMase or aSMase assay buffer was incubated for 2 h at 37°C. The reaction was terminated with 200 μ l CHCl₃/methanol (1:1) and 90 μ l H₂O followed by vigorous agitation. The samples were centrifuged at 6,000 g for 5 min. [¹⁴C]Phosphocholine in the aqueous phase (120 μ l) was collected for liquid scintillation counting. Phosphocholine is the degraded moiety of sphingomyelin after ceramide is released by nSMase or aSMase. The aSMase or nSMase activity was calculated as picomoles of sphingomyelin hydrolyzed by 1 mg of total proteins per hour and expressed as a percentage of control values.

TLC

OLGs with or without A β 25-35 treatment were cultured with 10 μ Ci [³H]palmitate (1 mCi/ml; Amersham Biosciences; Kaneko et al., 1995). The labeled cells were collected and washed twice with PBS, pH 7.4, to remove free isotope before lipid extraction (Xu et al., 1998). The cell pellet was resuspended in 400 μ l methanol/1 N HCl (100:6, vol/vol) followed by 800 μ l chloroform and 240 μ l H₂O. The sample was mixed and centrifuged at 6,000 g for 5 min. The lipid fraction was reextracted with 1 ml chloroform/methanol (2:1, vol/vol) and applied to a TLC plate. The solvent was chloroform/methanol/acetic acid/H₂O (85:4.5:5:0.5, vol/vol) for ceramide and chloroform/methanol/acetic acid/water (65:25:8.8:4.5, vol/vol) for sphingomyelin. Plates were air dried and sprayed with 1 M sodium salicylate for autoradiography. Standard lipids were stained by rhodamine 6G (Sigma-Aldrich) and visualized by UV light.

Quantitative ceramide analysis by ESI/MS

After three washes with PBS, pH 7.4, OLGs harvested from 100-mm culture plates with or without A β 25-35 treatment were homogenized in 0.5 ml PBS, pH 7.4, with a glass tissue grinder. A bicinchoninic protein assay kit was used to determine the protein concentration before lipid extraction (Pierce Chemical Co.). Lipids from the homogenates were extracted as described previously with modifications (Bligh and Dyer, 1959) using 50 mM LiOH in the aqueous layer and C17:0 ceramide (2 nmol/mg protein) as an internal standard for quantitation of ceramide content. These molecular species represent <1% of the endogenous cellular lipid mass. The lipid extracts were dried under a nitrogen stream, dissolved in chloroform, desalted with Sep-Pak columns, and filtered with 0.2 μ m PTFE syringe filters (Fisher Scientific). Lipids were reextracted with 20 mM LiOH in the aqueous layer, dried under a nitrogen stream, and resuspended in 0.5 ml of chloroform/methanol (1:1, vol/vol) for ESI/MS analysis.

ESI/MS analysis was performed using a spectrometer (model TSQ-7000; Finnigan) equipped with an electrospray ion source as described previously (Han et al., 1996, 2001). A 5-min period of signal averaging in the profile mode was used for each spectrum of a lipid extract. All extracts were directly infused into the ESI chamber using a syringe pump at 1 μ l/min flow rate. Ceramide in the lipid extracts was quantitated directly as deprotonated ions ([M-H]⁻) in comparison with an internal standard (C17:0 ceramide) after correction for ¹³C isotope effects in the negative-ion mode. Ion peaks were identified using tandem mass spectroscopic analyses as described previously (Han and Gross, 1995).

Cell death assays

OLG viability was quantitated by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay and trypan blue exclusion method. Cell death was also assessed by the amount of lactate dehydrogenase (LDH) release into the culture medium after A β or C2-ceramide treatment (Koh and Choi, 1987; Shaikh et al., 1997; Xu et al., 1998). The amount of LDH released by cells killed with Triton X-100 was considered maximal cell death or "full kill" (Xu et al., 1998).

Assay for cellular GSH content

Cellular GSH levels were determined using a GSH-400 colorimetric assay kit (Calbiochem-Novabiochem). Triplicate samples (3×10^6 cells) were collected by centrifugation and washed twice with PBS, pH 7.4. The cell pellets were treated with 5% metaphosphoric acid (Sigma-Aldrich). A Teflon pestle was used to homogenize the cells. Protein concentrations were determined by Lowry assay (Lowry et al., 1951). The homogenates were centrifuged at 3,000 g for 10 min at 4°C. Supernatants were assayed for GSH according to the instructions provided with the kit. A standard curve was generated with graded concentrations of GSH (5–40 μ M). GSH concentration was measured by absorbance at 400 nm with a spectrophotometer.

nSMase antisense oligonucleotides

Morpholino sense (GCCGACAGAAAAGTTGTGCTTCAT) and antisense (CCTCTTACCTCAGTTACAATTATA) oligonucleotides were generated for nSMase (Gene Tools, LLC). OLGs in serum-free medium were treated with 1.4 μ M oligonucleotides in EPEI delivery solution (per manufacturer's instructions; Gene Tools) for 4 h. The medium was exchanged and A β was added for 24 h, after which cells were harvested for nSMase activity or cell death determination.

Statistical analysis

Results are expressed as mean \pm SD. Differences among groups were analyzed by one-way ANOVA followed by Bonferroni's post-hoc *t* test to determine statistical significance. Comparison between two experimental groups was based on two-tailed *t* test. *P* < 0.05 was considered statistically significant.

We would like to thank Dr. Yannan Ouyang for his technical expertise in confocal microscopy.

This work was supported by National Institutes of Health grants NS37230 and NS40525.

Submitted: 2 July 2003

Accepted: 19 November 2003

References

- Abate, C., L. Patel, F.J. Rauscher, III, and T. Curran. 1990. Redox regulation of fos and jun DNA-binding activity in vitro. *Science*. 249:1157–1161.
- Albouz, S., F. Le Saux, D. Wenger, J.J. Hauw, and N. Baumann. 1986. Modifications of sphingomyelin and phosphatidylcholine metabolism by tricyclic antidepressants and phenothiazines. *Life Sci*. 38:357–363.
- Anderson, M.E., and A. Meister. 1983. Transport and direct utilization of gamma-glutamylcyst(e)ine for glutathione synthesis. *Proc. Natl. Acad. Sci. USA*. 80:707–711.
- Askanas, V., J. McFerrin, S. Baque, R.B. Alvarez, E. Sarkozi, and W.K. Engel. 1996. Transfer of beta-amyloid precursor protein gene using adenovirus vector causes mitochondrial abnormalities in cultured normal human muscle. *Proc. Natl. Acad. Sci. USA*. 93:1314–1319.
- Back, S.A., X. Gan, Y. Li, P.A. Rosenberg, and J.J. Volpe. 1998. Maturation-dependent vulnerability of oligodendrocytes to oxidative stress-induced death caused by glutathione depletion. *J. Neurosci*. 18:6241–6253.
- Behl, C. 1999. Alzheimer's disease and oxidative stress: implications for novel therapeutic approaches. *Prog. Neurobiol*. 57:301–323.
- Behl, C., and Y. Sagara. 1997. Mechanism of amyloid beta protein induced neuronal cell death: current concepts and future perspectives. *J. Neural Transm. Suppl*. 49:125–134.
- Behl, C., J.B. Davis, R. Lesley, and D. Schubert. 1994. Hydrogen peroxide mediates amyloid beta protein toxicity. *Cell*. 77:817–827.
- Bezombes, C., N. Maestre, G. Laurent, T. Levade, A. Bettaieb, and J.P. Jaffrezou. 1998. Restoration of TNF-alpha-induced ceramide generation and apoptosis in resistant human leukemia KG1a cells by the P-glycoprotein blocker PSC833. *FASEB J*. 12:101–109.
- Blasko, I., T.L. Schmitt, E. Steiner, K. Trieb, and B. Grubeck-Loebenstein. 1997. Tumor necrosis factor alpha augments amyloid beta protein (25-35) induced apoptosis in human cells. *Neurosci. Lett*. 238:17–20.
- Bligh, E.G., and W.J. Dyer. 1959. A rapid method of total lipid extraction and purification. *Can. J. Med. Sci*. 37:911–917.
- Bose, R., M. Verheij, A. Haimovitz-Friedman, K. Scotto, Z. Fuks, and R. Kolesnick. 1995. Ceramide synthase mediates daunorubicin-induced apoptosis: an alternative mechanism for generating death signals. *Cell*. 82:405–414.
- Bozner, P., V. Grishko, S.P. LeDoux, G.L. Wilson, Y.C. Chyan, and M.A. Pap-polla. 1997. The amyloid beta protein induces oxidative damage of mitochondrial DNA. *J. Neuropathol. Exp. Neurol*. 56:1356–1362.
- Bradshaw, C.D., K.M. Ella, A.L. Thomas, C. Qi, and K.E. Meier. 1996. Effects of Ara-C on neutral sphingomyelinase and mitogen- and stress-activated protein kinases in T-lymphocyte cell lines. *Biochem. Mol. Biol. Int*. 40:709–719.
- Brera, B., A. Serrano, and M.L. de Ceballos. 2000. beta-amyloid peptides are cytotoxic to astrocytes in culture: a role for oxidative stress. *Neurobiol. Dis*. 7:395–405.
- Bruce-Keller, A.J., J.G. Begley, W. Fu, D.A. Butterfield, D.E. Bredesen, J.B. Hutchins, K. Hensley, and M.P. Mattson. 1998. Bcl-2 protects isolated plasma and mitochondrial membranes against lipid peroxidation induced by hydrogen peroxide and amyloid beta-peptide. *J. Neurochem*. 70:31–39.
- Bruno, A.P., G. Laurent, D. Averbeck, C. Demur, J. Bonnet, A. Bettaieb, T. Levade, and J.P. Jaffrezou. 1998. Lack of ceramide generation in TF-1 human myeloid leukemic cells resistant to ionizing radiation. *Cell Death Differ*. 5:172–182.
- Cafe, C., C. Torri, L. Bertorelli, N. Angeretti, E. Lucca, G. Forloni, and F. Marzatico. 1996. Oxidative stress after acute and chronic application of beta-amyloid fragment 25-35 in cortical cultures. *Neurosci. Lett*. 203:61–65.
- Coroneos, E., M. Martinez, S. McKenna, and M. Kester. 1995. Differential regulation of sphingomyelinase and ceramidase activities by growth factors and cytokines. Implications for cellular proliferation and differentiation. *J. Biol. Chem*. 270:23305–23309.
- Davis-Salinas, J., S.M. Saporito-Irwin, C.W. Cotman, and W.E. Van Nostrand. 1995. Amyloid beta-protein induces its own production in cultured degenerating cerebrovascular smooth muscle cells. *J. Neurochem*. 65:931–934.
- de la Monte, S.M., Y.K. Sohn, and J.R. Wands. 1997. Correlates of p53- and Fas (CD95)-mediated apoptosis in Alzheimer's disease. *J. Neurol. Sci*. 152:73–83.
- Dobrowsky, R.T., G.M. Jenkins, and Y.A. Hannun. 1995. Neurotrophins induce sphingomyelin hydrolysis. Modulation by co-expression of p75NTR with Trk receptors. *J. Biol. Chem*. 270:22135–22142.
- Estus, S., T.E. Golde, T. Kunishita, D. Blades, D. Lowery, M. Eisen, M. Usiak, X.M. Qu, T. Tabira, B.D. Greenberg, et al. 1992. Potentially amyloidogenic, carboxyl-terminal derivatives of the amyloid protein precursor. *Science*. 255:726–728.
- Garcia-Ruiz, C., A. Colell, M. Mari, A. Morales, and J.C. Fernandez-Checa. 1997. Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species. Role of mitochondrial glutathione. *J. Biol. Chem*. 272:11369–11377.
- Goldkorn, T., N. Balaban, M. Shannon, V. Chea, K. Matsukuma, D. Gilchrist, H. Wang, and C. Chan. 1998. H₂O₂ acts on cellular membranes to generate ceramide signaling and initiate apoptosis in tracheobronchial epithelial cells. *J. Cell Sci*. 111:3209–3220.
- Haass, C., M.G. Schlossmacher, A.Y. Hung, C. Vigo-Pelfrey, A. Mellon, B.L. Ostaszewski, I. Lieberburg, E.H. Koo, D. Schenk, D.B. Teplow, et al. 1992. Amyloid beta-peptide is produced by cultured cells during normal metabolism. *Nature*. 359:322–325.
- Haimovitz-Friedman, A., C.C. Kan, D. Ehleiter, R.S. Persaud, M. McLoughlin, Z. Fuks, and R.N. Kolesnick. 1994. Ionizing radiation acts on cellular membranes to generate ceramide and initiate apoptosis. *J. Exp. Med*. 180:525–535.
- Han, X., and R.W. Gross. 1995. Structural determination of picomole amounts of phospholipids via electrospray ionization tandem mass spectrometry. *J. Am. Soc. Mass Spectrom*. 6:1202–1210.
- Han, X., R.A. Gubitosi-Klug, B.J. Collins, and R.W. Gross. 1996. Alterations in individual molecular species of human platelet phospholipids during thrombin stimulation: electrospray ionization mass spectrometry-facilitated identification of the boundary conditions for the magnitude and selectivity of thrombin-induced platelet phospholipid hydrolysis. *Biochemistry*. 35:5822–5832.
- Han, X., D.M. Holtzman, and D.W. McKeel. 2001. Plasmalogen deficiency in early Alzheimer's disease subjects and in animal models: molecular characterization using electrospray ionization mass spectrometry. *J. Neurochem*. 77:1168–1180.
- Hannun, Y.A. 1996. Functions of ceramide in coordinating cellular responses to stress. *Science*. 274:1855–1859.
- Hannun, Y.A., and C. Luberto. 2000. Ceramide in the eukaryotic stress response. *Trends Cell Biol*. 10:73–80.
- Huang, S.S., F.W. Huang, J. Xu, S. Chen, C.Y. Hsu, and J.S. Huang. 1998. Amyloid beta-peptide possesses a transforming growth factor-beta activity. *J. Biol. Chem*. 273:27640–27644.
- Jaffrezou, J.P., T. Levade, A. Bettaieb, N. Andrieu, C. Bezombes, N. Maestre, S.

- Vermeersch, A. Rousse, and G. Laurent. 1996. Daunorubicin-induced apoptosis: triggering of ceramide generation through sphingomyelin hydrolysis. *EMBO J.* 15:2417–2424.
- Jan, J.T., S. Chatterjee, and D.E. Griffin. 2000. Sindbis virus entry into cells triggers apoptosis by activating sphingomyelinase, leading to the release of ceramide. *J. Virol.* 74:6425–6432.
- Jarvis, W.D., R.N. Kolesnick, F.A. Fornari, R.S. Traylor, D.A. Gewirtz, and S. Grant. 1994. Induction of apoptotic DNA damage and cell death by activation of the sphingomyelin pathway. *Proc. Natl. Acad. Sci. USA.* 91:73–77.
- Juurlink, B.H., S.K. Thorburne, and L. Hertz. 1998. Peroxide-scavenging deficit underlies oligodendrocyte susceptibility to oxidative stress. *Glia.* 22:371–378.
- Kaneko, I., N. Yamada, Y. Sakuraba, M. Kamenosono, and S. Tutumi. 1995. Suppression of mitochondrial succinate dehydrogenase, a primary target of beta-amyloid, and its derivative racemized at Ser residue. *J. Neurochem.* 65:2585–2593.
- Kawai, M., R.N. Kalaria, P. Cras, S.L. Siedlak, M.E. Velasco, E.R. Shelton, H.W. Chan, B.D. Greenberg, and G. Perry. 1993. Degeneration of vascular muscle cells in cerebral amyloid angiopathy of Alzheimer disease. *Brain Res.* 623:142–146.
- Koh, J.Y., and D.W. Choi. 1987. Quantitative determination of glutamate mediated cortical neuronal injury in cell culture by lactate dehydrogenase efflux assay. *J. Neurosci. Methods.* 20:83–90.
- Kolesnick, R.N., and M. Kronke. 1998. Regulation of ceramide production and apoptosis. *Annu. Rev. Physiol.* 60:643–665.
- Kyriakis, J.M., and J. Avruch. 1996. Protein kinase cascades activated by stress and inflammatory cytokines. *Bioessays.* 18:567–577.
- Larocca, J.N., M. Farooq, and W.T. Norton. 1997. Induction of oligodendrocyte apoptosis by C2-ceramide. *Neurochem. Res.* 22:529–534.
- Levade, T., and J.P. Jaffrezou. 1999. Signalling sphingomyelinases: which, where, how and why? *Biochim. Biophys. Acta.* 1438:1–17.
- Lister, M.D., Z.S. Ruan, and R. Bittman. 1995. Interaction of sphingomyelinase with sphingomyelin analogs modified at the C-1 and C-3 positions of the sphingosine backbone. *Biochim. Biophys. Acta.* 1256:25–30.
- Liu, B., and Y.A. Hannun. 1997. Inhibition of the neutral magnesium-dependent sphingomyelinase by glutathione. *J. Biol. Chem.* 272:16281–16287.
- Liu, B., N. Andrieu-Abadie, T. Levade, P. Zhang, L.M. Obeid, and Y.A. Hannun. 1998a. Glutathione regulation of neutral sphingomyelinase in tumor necrosis factor-alpha-induced cell death. *J. Biol. Chem.* 273:11313–11320.
- Liu, B., D.F. Hassler, G.K. Smith, K. Weaver, and Y.A. Hannun. 1998b. Purification and characterization of a membrane bound neutral pH optimum magnesium-dependent and phosphatidylserine-stimulated sphingomyelinase from rat brain. *J. Biol. Chem.* 273:34472–34479.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr, and R.J. Randall. 1951. Protein measurement with the Folin-Phenol reagents. *J. Biol. Chem.* 193:265–275.
- Markesbery, W.R. 1999. The role of oxidative stress in Alzheimer disease. *Arch. Neurol.* 56:1449–1452.
- Masters, C.L., G. Multhaup, G. Simms, J. Pottgiesser, R.N. Martins, and K. Beyreuther. 1985. Neuronal origin of a cerebral amyloid: neurofibrillary tangles of Alzheimer's disease contain the same protein as the amyloid of plaque cores and blood vessels. *EMBO J.* 4:2757–2763.
- Masukawa, T., T. Nishimura, H. Kito, and H. Iwata. 1983. Influence of diethylmaleate on the formation of bis(methylmercuric)selenide and methylmercury distribution in rats. *J. Pharmacobiodyn.* 6:950–953.
- McDonald, J.W., X.Z. Liu, Y. Qu, S. Liu, S.K. Mickey, D. Turetsky, D.I. Gottlieb, and D.W. Choi. 1999. Transplanted embryonic stem cells survive, differentiate and promote recovery in injured rat spinal cord. *Nat. Med.* 5:1410–1412.
- Muller, W.E., F.J. Romero, S. Perovic, G. Pergande, and P. Pialoglou. 1997. Protection of flupirtine on beta-amyloid-induced apoptosis in neuronal cells in vitro: prevention of amyloid-induced glutathione depletion. *J. Neurochem.* 68:2371–2377.
- Okazaki, T., R.M. Bell, and Y.A. Hannun. 1989. Sphingomyelin turnover induced by vitamin D3 in HL-60 cells. Role in cell differentiation. *J. Biol. Chem.* 264:19076–19080.
- Pahan, K., F.G. Sheikh, M. Khan, A.M. Namboodiri, and I. Singh. 1998. Sphingomyelinase and ceramide stimulate the expression of inducible nitric-oxide synthase in rat primary astrocytes. *J. Biol. Chem.* 273:2591–2600.
- Pereira, C., M.S. Santos, and C. Oliveira. 1999. Involvement of oxidative stress on the impairment of energy metabolism induced by A beta peptides on PC12 cells: protection by antioxidants. *Neurobiol. Dis.* 6:209–219.
- Pinkus, R., L.M. Weiner, and V. Daniel. 1996. Role of oxidants and antioxidants in the induction of AP-1, NF-kappaB, and glutathione S-transferase gene expression. *J. Biol. Chem.* 271:13422–13429.
- Sawai, H., and Y.A. Hannun. 1999. Ceramide and sphingomyelinases in the regulation of stress responses. *Chem Phys. Lipids.* 102:141–147.
- Schapira, A.H. 1996. Oxidative stress and mitochondrial dysfunction in neurodegeneration. *Curr. Opin. Neurol.* 9:260–264.
- Schreck, R., P. Rieber, and P.A. Baeuerle. 1991. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *EMBO J.* 10:2247–2258.
- Scurlock, B., and G. Dawson. 1999. Differential responses of oligodendrocytes to tumor necrosis factor and other pro-apoptotic agents: role of ceramide in apoptosis. *J. Neurosci. Res.* 55:514–522.
- Segui, B., N. Andrieu-Abadie, S. Adam-Klages, O. Meilhan, D. Kreder, V. Garcia, A.P. Bruno, J.P. Jaffrezou, R. Salvayre, M. Kronke, and T. Levade. 1999. CD40 signals apoptosis through FAN-regulated activation of the sphingomyelin-ceramide pathway. *J. Biol. Chem.* 274:37251–37258.
- Shaikh, A.Y., J. Xu, Y. Wu, L. He, and C.Y. Hsu. 1997. Melatonin protects bovine cerebral endothelial cells from hyperoxia-induced DNA damage and death. *Neurosci. Lett.* 229:193–197.
- Singh, I., K. Pahan, M. Khan, and A.K. Singh. 1998. Cytokine-mediated induction of ceramide production is redox-sensitive. Implications to proinflammatory cytokine-mediated apoptosis in demyelinating diseases. *J. Biol. Chem.* 273:20354–20362.
- Soderberg, M., C. Edlund, I. Alafuzoff, K. Kristensson, and G. Dallner. 1992. Lipid composition in different regions of the brain in Alzheimer's disease/senile dementia of Alzheimer's type. *J. Neurochem.* 59:1646–1653.
- Spiegel, S., and A.H. Merrill, Jr. 1996. Sphingolipid metabolism and cell growth regulation. *FASEB J.* 10:1388–1397.
- Sugita, M., M. Willians, J.T. Dulaney, and H.W. Moser. 1975. Ceramidase and ceramide synthesis in human kidney and cerebellum. Description of a new alkaline ceramidase. *Biochim. Biophys. Acta.* 398:125–131.
- Szekely, J.G., and A.U. Lobreau. 1987. The effect of glutathione depletion by diamide, diethyl maleate or buthione sulfoximine on the surface structure of mouse L-cells. *Scanning Microsc.* 1:273–281.
- Tepper, C.G., S. Jayadev, B. Liu, A. Bielawska, R. Wolff, S. Yonehara, Y.A. Hannun, and M.F. Seldin. 1995. Role for ceramide as an endogenous mediator of Fas-induced cytotoxicity. *Proc. Natl. Acad. Sci. USA.* 92:8443–8447.
- Testi, R. 1996. Sphingomyelin breakdown and cell fate. *Trends Biochem. Sci.* 21:468–471.
- Thomas, T., G. Thomas, C. McLendon, T. Sutton, and M. Mullan. 1996. beta-Amyloid-mediated vasoactivity and vascular endothelial damage. *Nature.* 380:168–171.
- Thorburne, S.K., and B.H. Juurlink. 1996. Low glutathione and high iron govern the susceptibility of oligodendroglial precursors to oxidative stress. *J. Neurochem.* 67:1014–1022.
- Tonnetti, L., M.C. Veri, E. Bonvini, and L. D'Adamo. 1999. A role for neutral sphingomyelinase-mediated ceramide production in T cell receptor-induced apoptosis and mitogen-activated protein kinase-mediated signal transduction. *J. Exp. Med.* 189:1581–1589.
- Verheij, M., R. Bose, X.H. Lin, B. Yao, W.D. Jarvis, S. Grant, M.J. Birrer, E. Szabo, L.I. Zon, J.M. Kyriakis, et al. 1996. Requirement for ceramide-initiated SAPK/JNK signalling in stress-induced apoptosis. *Nature.* 380:75–79.
- Xu, J., C.H. Yeh, S. Chen, L. He, S.L. Sensi, L.M. Canzoniero, D.W. Choi, and C.Y. Hsu. 1998. Involvement of de novo ceramide biosynthesis in tumor necrosis factor-alpha/cycloheximide-induced cerebral endothelial cell death. *J. Biol. Chem.* 273:16521–16526.
- Xu, J., S. Chen, S.H. Ahmed, H. Chen, G. Ku, M.P. Goldberg, and C.Y. Hsu. 2001. Amyloid-beta peptides are cytotoxic to oligodendrocytes. *J. Neurosci.* 21:RC118.
- Yaar, M., S. Zhai, P.F. Pilch, S.M. Doyle, P.B. Eisenhauer, R.E. Fine, and B.A. Gilchrist. 1997. Binding of beta-amyloid to the p75 neurotrophin receptor induces apoptosis. A possible mechanism for Alzheimer's disease. *J. Clin. Invest.* 100:2333–2340.
- Yankner, B.A., L.R. Dawes, S. Fisher, L. Villa-Komaroff, M.L. Oster-Granite, and R.L. Neve. 1989. Neurotoxicity of a fragment of the amyloid precursor associated with Alzheimer's disease. *Science.* 245:417–420.
- Yoshimura, S., Y. Banno, S. Nakashima, K. Takenaka, H. Sakai, Y. Nishimura, N. Sakai, S. Shimizu, Y. Eguchi, Y. Tsujimoto, and Y. Nozawa. 1998. Ceramide formation leads to caspase-3 activation during hypoxic PC12 cell death. Inhibitory effects of Bcl-2 on ceramide formation and caspase-3 activation. *J. Biol. Chem.* 273:6921–6927.
- Yoshimura, S., Y. Banno, S. Nakashima, K. Hayashi, H. Yamakawa, M. Sawada, N. Sakai, and Y. Nozawa. 1999. Inhibition of neutral sphingomyelinase acti-

- vation and ceramide formation by glutathione in hypoxic PC12 cell death. *J. Neurochem.* 73:675–683.
- Yu, S.P., Z.S. Farhangrazi, H.S. Ying, C.H. Yeh, and D.W. Choi. 1998. Enhancement of outward potassium current may participate in beta-amyloid peptide-induced cortical neuronal death. *Neurobiol. Dis.* 5:81–88.
- Zhang, P., B. Liu, G.M. Jenkins, Y.A. Hannun, and L.M. Obeid. 1997. Expression of neutral sphingomyelinase identifies a distinct pool of sphingomyelin involved in apoptosis. *J. Biol. Chem.* 272:9609–9612.
- Zhang, S.C., B. Ge, and I.D. Duncan. 1999. Adult brain retains the potential to generate oligodendroglial progenitors with extensive myelination capacity. *Proc. Natl. Acad. Sci. USA.* 96:4089–4094.