

2009

## G protein subunit dissociation and translocation regulate cellular response to receptor stimulation

Mariangela Chisari  
*Washington University School of Medicine in St. Louis*

Deepak Kumar Saini  
*Washington University School of Medicine in St. Louis*

Joon-Ho Cho  
*Washington University School of Medicine in St. Louis*

Vani Kalyanaraman  
*Washington University School of Medicine in St. Louis*

N. Gautam  
*Washington University School of Medicine in St. Louis*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)



Part of the [Medicine and Health Sciences Commons](#)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Chisari, Mariangela; Saini, Deepak Kumar; Cho, Joon-Ho; Kalyanaraman, Vani; and Gautam, N., "G protein subunit dissociation and translocation regulate cellular response to receptor stimulation." *PLoS One*. 4, 11. e7797. (2009).

[https://digitalcommons.wustl.edu/open\\_access\\_pubs/772](https://digitalcommons.wustl.edu/open_access_pubs/772)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

# G Protein Subunit Dissociation and Translocation Regulate Cellular Response to Receptor Stimulation

Mariangela Chisari<sup>1</sup>, Deepak Kumar Saini<sup>1</sup>, Joon-Ho Cho<sup>1</sup>, Vani Kalyanaraman<sup>1</sup>, N. Gautam<sup>1,2\*</sup>

**1** Department of Anesthesiology, Washington University School of Medicine, St. Louis, Missouri, United States of America, **2** Department of Genetics, Washington University School of Medicine, St. Louis, Missouri, United States of America

## Abstract

We examined the role of G proteins in modulating the response of living cells to receptor activation. The response of an effector, phospholipase C- $\beta$  to M3 muscarinic receptor activation was measured using sensors that detect the generation of inositol triphosphate or diacylglycerol. The recently discovered translocation of G $\beta\gamma$  from plasma membrane to endomembranes on receptor activation attenuated this response. A FRET based G protein sensor suggested that in contrast to translocating G $\beta\gamma$ , non-translocating G $\beta\gamma$  subunits do not dissociate from the  $\alpha_q$  subunit on receptor activation leading to prolonged retention of the heterotrimer state and an accentuated response. M3 receptors with tethered  $\alpha_q$  induced differential responses to receptor activation in cells with or without an endogenous translocation capable  $\gamma$  subunit. G protein heterotrimer dissociation and  $\beta\gamma$  translocation are thus unanticipated modulators of the intensity of a cell's response to an extracellular signal.

**Citation:** Chisari M, Saini DK, Cho J-H, Kalyanaraman V, Gautam N (2009) G Protein Subunit Dissociation and Translocation Regulate Cellular Response to Receptor Stimulation. PLoS ONE 4(11): e7797. doi:10.1371/journal.pone.0007797

**Editor:** Richard Steinhardt, University of California, Berkeley, United States of America

**Received:** April 8, 2009; **Accepted:** October 16, 2009; **Published:** November 11, 2009

**Copyright:** © 2009 Chisari et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This study was supported by National Institute of Health grants (GM69027 and GM080558) and American Heart Association postdoctoral fellowships (M.C. and D.K.S.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: gautam@wustl.edu

## Introduction

G proteins are the major modulators of cellular responses to external signals in mammalian cells [1,2]. There is limited information on the role that G proteins play in directly regulating the sensitivity of a cell's response to an external stimulus in living cells. Studies in intact yeast cells [3] and in a mammalian cell line [4] have attempted to address the quantitative relationship between G protein activation and activity downstream. The kinetics of the rod photoreceptor G protein, Gt, mediated phototransduction activity has also been examined in highly specialized rod photoreceptor cells which are amenable to such studies [5]. Overall however, little is known about such processes with regard to the large families of G protein subunits that are expressed widely in all mammalian cell types. Here we have used imaging methods to examine whether mechanisms at the level of the G protein subunits control the intensity of the response to receptor activation in intact living cells.

Recently we demonstrated that on receptor activation, a family of G protein  $\beta\gamma$  complexes translocates from the plasma membrane to endomembranes and reverse translocates when a receptor is deactivated [6,7]. The translocation reduces the concentration of G $\beta\gamma$  on the plasma membrane rapidly ( $t_{1/2} \sim 10$  s) [6,7]. We expected this reduced concentration of G $\beta\gamma$  on the plasma membrane would attenuate the response of an effector, phospholipase  $\beta$  (PLC $\beta$ ) to the activation of a receptor coupled to the G protein, G $q$ . We tested this hypothesis using live cell imaging experiments using the PLC $\beta$  mediated signaling pathway.

M3 muscarinic receptor activation of the G $\alpha_q$  activates phospholipases C  $\beta$  isozymes (PLCs), leading to phosphatidylinositol (4,5)-biphosphate (PIP2) hydrolysis on the plasma membrane

resulting in the production of inositol triphosphate (IP3) and diacylglycerol (DAG) [8]. PIP2 breakdown is detected quantitatively in living cells by imaging the translocation of a pleckstrin homology (PH) domain of PLC $\delta$  tagged with a fluorescent protein, mCherry (mCh) [9] or a DAG binding domain of protein kinase C  $\beta$  (DBD) tagged with yellow fluorescent protein (YFP) [10,11,12]. When M3 receptors are stimulated, PH-mCh translocates from the plasma membrane to the cytosol because it binds with higher affinity to IP3 in the cytosol compared to PIP2 in the plasma membrane [13]. YFP-DBD translocates in the opposite direction from the cytosol to the plasma membrane because it binds to DAG [11]. Using these sensors we examined the impact of activating G proteins containing translocating and non-translocating  $\beta\gamma$  complexes on downstream PLC  $\beta$  effector response. We then used a fluorescence resonance energy transfer (FRET) based G protein sensor similar to a previous sensor [14] containing different  $\gamma$  subunits to identify the mechanistic basis of response differences. Finally, to evaluate the relevance of our observation at physiological concentrations of G $\beta\gamma$ , we used an  $\alpha_q$  subunit tethered to the M3 receptor to evaluate the effect of endogenous translocating and non-translocating G $\gamma$  subunits.

## Results

### Functionality of fluorescently tagged fusion proteins


Fluorescently tagged G protein  $\gamma$  subunits used in this study have been previously described [6,7] and have been shown to be functionally active [14,15,16,17]. We also confirmed that the  $\alpha_q$ -CFP fusion protein was functional by examining PH-mCh translocation in M3-CHO cells which were transfected with  $\alpha_q$ -CFP and PH-mCh. PH-mCh translocates from the plasma

membrane to the cytosol on PLC  $\beta$  activation and the magnitude of PH-mCh translocation is indicative of extent of PIP2 breakdown on the PM. In cells containing  $\alpha$ q-CFP, the PH – domain translocation was significantly higher after receptor activation compared to untransfected cells (emission intensity in the cytosol was  $\sim 1.6$  fold higher (data not shown) indicating that the introduced  $\alpha$ q-CFP was active. We further confirmed functionality of  $\alpha$ q-CFP by evaluating its ability to support translocation of the  $\beta\gamma$  complex. Endogenous  $\alpha$ q does not support visually detectable  $\beta\gamma$  translocation (data not shown). This absence of background translocation of  $\beta\gamma$  induced by endogenous  $\alpha$ q allowed us to examine whether the tagged  $\alpha$ q-CFP was functional. We introduced  $\alpha$ q-CFP into M3-CHO cells along with YFP tagged  $\beta\gamma$ . On activation of M3 receptor in these cells,  $\beta\gamma$  translocated from the plasma membrane to the intracellular membranes (Fig. 1A, right panel). These results showed that  $\alpha$ q-CFP fusion protein is functionally active.

### PH domain translocation is reduced in the presence of a translocating G protein $\gamma$ subunit

To test the hypothesis that  $\gamma$  subunit translocation modifies the sensitivity of a cell's response to an agonist, PIP2 breakdown was measured at various concentrations of agonist. Gq containing different  $\gamma$  subunits tagged with fluorescent proteins were cotransfected into cells stably expressing M3 muscarinic receptors (M3-CHO) [14] along with the PH-mCh. Individual cells with

similar emission levels of CFP and YFP were chosen for analysis. Cells were exposed independently to concentrations of agonist ranging from 250 nM to 100  $\mu$ M. Buffer was introduced after agonist activation to deactivate the receptor. Emission intensities from the cytosol for PH-mCh and from intracellular membranes for YFP- $\gamma$  subunits were measured to quantify the extent of translocation of these proteins from the plasma membrane where they were initially localized (Fig. 1A). For measuring PH-mCh changes we selected regions of approximately equal size in the cytosol. We ensured that the changes within selected regions were reflected overall in the cell cytosol and not restricted to the region selected (Fig. 1A). Images (Fig. 1A) and bar diagram (Fig. 1B) show the extent of PH-mCh translocation in the presence of the  $\gamma$ 3 or  $\gamma$ 9 subunits when cells were exposed to individual agonist concentrations (as indicated). Cells expressing  $\gamma$ 3 at subsaturating agonist concentrations of 250 nM and 1  $\mu$ M showed a response that was significantly higher compared to cells expressing  $\gamma$ 9 (Fig. 1B). This difference is seen even at 5  $\mu$ M concentration (Fig. 1B). The PIP2 breakdown stimulated by the M3 receptor begins to saturate above 5  $\mu$ M (Fig. 1B). Fig. 1C shows the magnitude of YFP- $\gamma$ 9 translocation determined by measuring the fluorescence intensity in the plasma membrane before and after agonist addition. The YFP emission intensity on the plasma membrane was calculated as the average of selected regions as shown in the example in Fig. 1A. The magnitude of  $\beta\gamma$  translocation is directly correlated with increasing agonist concentration. These results together support



**Figure 1. Responses of cells expressing various  $\gamma$  subunits to different concentrations of agonist.** A. M3-CHO cells expressing  $\alpha$ q-CFP, YFP- $\gamma$  (as indicated) and PH-mCh were imaged as mentioned in Methods and Text S1. Representative images of cells exposed to 1  $\mu$ M carbachol (agonist) are shown. The decrease in YFP- $\gamma$ 3 intensity observed is due to bleaching during acquisition. B. Translocation of PH-mCh at specific agonist concentrations. Bars show changes for PH-mCh translocation in the presence of various  $\gamma$  subunits. No. of  $\gamma$ 3 cells examined at 250 nM–20; at 1  $\mu$ M–11; at 5  $\mu$ M–17 and at 100  $\mu$ M–17. No. of  $\gamma$ 9 cells examined at 250 nM–23; at 1  $\mu$ M–11; at 5  $\mu$ M–17 and at 100  $\mu$ M–14. Cells with comparable expression of PH-mCh were imaged. Regions in the cytosol were selected for analysis (as indicated by circles). Values were calculated as ratio of M-B/B, where M is the maximal intensity at a particular concentration and B, the intensity before exposure to that concentration of agonist. Results are the means  $\pm$  SEM. \* $p$ <0.01, \*\* $p$ <0.001. C. Translocation of YFP- $\gamma$ 9 subunits at specific agonist concentrations. Bar diagrams show changes in YFP- $\gamma$ 9 emission on the plasma membrane (N=4). Regions on the plasma membrane were selected for analysis (as indicated by arrows in image, Fig. 1A, right panel). Values are shown as percentile of means  $\pm$  SEM. \* $p$ <0.01, (response to 250 nM and 100  $\mu$ M to others). doi:10.1371/journal.pone.0007797.g001

the model that the magnitude of cellular response is decreased by the removal of the  $\beta\gamma$  complex from the plasma membrane on receptor stimulation (Fig. 2). Conversely, cells expressing non-translocating  $\gamma$  subunits show higher sensitivity to an agonist (Fig. 2).

### The cumulative response of cells expressing $\gamma 3$ and $\gamma 9$ to sequential addition of increasing concentrations of agonist are the same

We then measured the cumulative response of these cells to the sequential exposure of cells to 1  $\mu\text{M}$  and 100  $\mu\text{M}$ . While no detectable translocation of the  $\gamma 3$  subunit was observed as reported previously [7] (Fig. 3A),  $\gamma 9$  translocated in response to the initial stimulus with 1  $\mu\text{M}$  and then to the subsequent addition of 100  $\mu\text{M}$  agonist (Fig. 3A). In cells expressing YFP- $\gamma 3$  or endogenous  $\gamma$  (no introduced  $\gamma$  subunit), the amount of PH-mCh translocation at 1  $\mu\text{M}$  was very strong compared to those cells expressing YFP- $\gamma 9$ . With the sequential addition of higher concentration of carbachol (100  $\mu\text{M}$ ), the additional PH-mCh translocation in the presence of  $\gamma 3$  or endogenous  $\gamma$  showed a small increase. In contrast, the response to 100  $\mu\text{M}$  agonist was significantly stronger in cells co-expressing  $\gamma 9$  (Fig. 3A). In Fig. 3B representative plots of the time dependent responses of individual cells to the sequential addition of agonist are compared for cells expressing different  $\gamma$  subunits. This comparison shows that the cumulative response to the sequential addition of 1  $\mu\text{M}$  and 100  $\mu\text{M}$  agonist is similar in all cells. Overall these results show that the differential response of cells expressing different  $\gamma$  subunits to low agonist concentrations is not due to inhibition of PIP2 breakdown.

### Cells expressing a chimeric $\gamma 9-3$ subunit which is incapable of translocation shows heightened receptor sensitivity compared to $\gamma 9$


We then examined whether the differential impact of  $\gamma$  subunits on PIP2 breakdown is due to unidentified differential roles of the  $\gamma 3$  and  $\gamma 9$  subunit types or due to the translocation of the  $\beta\gamma$  complex away from the plasma membrane to internal membranes. To confirm the role of  $\beta\gamma$  translocation we constructed a chimeric  $\gamma 9$  subunit in which the C terminal 15 residues of the translocation proficient  $\gamma 9$  were replaced with the corresponding sequence of  $\gamma 3$  which does not translocate ( $\gamma 9-3$ , Fig. 4A). We have previously shown that the C terminal domain of a  $\gamma$  subunit controls its

translocation capability [7,18]. The chimeric molecule,  $\gamma 9-3$ , did not translocate in response to receptor activation thus acquiring the properties of  $\gamma 3$  while retaining most of the residues of  $\gamma 9$  upstream of the C terminal 15 residue domain (Fig. 4B). We analyzed the receptor stimulated translocation of the PH domain in the presence of  $\gamma 9-3$  in selected cells as described above. Responses to both 1  $\mu\text{M}$  and 100  $\mu\text{M}$  agonist, added sequentially, were measured. PH-mCh translocation in the presence of the  $\gamma 9-3$  subunit (Fig. 4B) was similar to that observed in the presence of  $\gamma 3$  or endogenous  $\gamma$  subunits in CHO cells (Fig. 3A and B). This result clearly shows that receptor stimulated downstream activity is controlled by the translocation of the  $\beta\gamma$  complex away from the plasma membrane and is not an unusual property of a particular  $\gamma$  subunit type.

The results of the experiments using sequential addition of agonists was examined further to compare the effects of the  $\gamma$  subunit constitution of the cells on effector response. A summary of these results is shown in Fig. 4C. The histograms show that the magnitude of the downstream signaling activity (PH domain translocation) is distinctly different in cells that contain a translocation capable  $\gamma$  subunit,  $\gamma 9$ , compared to others (no  $\gamma$ ,  $\gamma 3$  or  $\gamma 9-3$ ). The response to 1  $\mu\text{M}$  concentration of agonist in  $\gamma 3$  containing cells is 1.7 fold higher compared to cells expressing  $\gamma 9$ . The response to 100  $\mu\text{M}$  agonist in  $\gamma 9$  containing cells was 1.8 fold higher compared to cells expressing  $\gamma 3$ . These results show that the response is significantly more sensitive to the initial low agonist concentration when the cell contains a non-translocating  $\gamma$  subunit. At the higher concentration of agonist only the cells expressing the translocating  $\gamma$  subunit show a strong additional response (Fig. 4C and D). These results confirm that cells in which receptor stimulation leads to  $\beta\gamma$  translocation are more sensitive in terms of their effector responses compared to cells in which translocation does not occur.


### FRET-based G protein sensor shows differential dissociation properties between G proteins containing translocating and non-translocating $\gamma$ subunit

To examine the mechanism at the basis of the effect of different  $\gamma$  subunits on the sensitivity of a cell to activation, we used a FRET based sensor containing  $\alpha_q$ -CFP and YFP- $\gamma$ . The design of this sensor was based on a previous sensor we had developed using  $\alpha_o$  [14]. Briefly, the donor CFP is tagged to  $\alpha_q$  subunit and acceptor YFP to G $\gamma$ . In the basal state due to formation of G protein




**Figure 2. Model for the mechanistic basis of differential sensitivity of cells containing different  $\gamma$  subunits.** On activation, G proteins progress through two different paths: (i) when they contain a non-translocating  $\gamma$  subunit (left) or (ii) when they contain a translocating  $\gamma$  subunit (right). Left: the activated  $\alpha$  and  $\beta\gamma$  remain on the plasma membrane. Since receptor activation requires the heterotrimer, this results in continual activation of G $\alpha_q$  reflected in a high level of PLC $\beta$  activation (multiple arrows). Right: the translocation of  $\beta\gamma$  away from the plasma membrane reduces the heterotrimer concentration and subsequent PLC $\beta$  stimulation. Although, we have been unable to obtain evidence for  $\beta\gamma$  activation of PLC $\beta$  in these cells (Text S2), this model is consistent with the results even if  $\beta\gamma$  activates PLC $\beta$  because  $\beta\gamma$  concentration available for PLC $\beta$  activation is lowered by translocation.

doi:10.1371/journal.pone.0007797.g002



**Figure 3. Responses of cells expressing different  $\gamma$  subunits to sequential exposure of increasing concentrations of agonist.** **A.** M3-CHO cells expressing  $\alpha$ q-CFP, YFP- $\gamma$  (as indicated) and PH-mCh were imaged as mentioned in Methods and Text S1. Representative images of cells exposed sequentially to 1  $\mu$ M and 100  $\mu$ M carbachol (agonist) are shown. Changes were measured in selected regions (black circles) in cytosol (for mCh) and endomembranes (for YFP). In the corresponding plots on the right, arrows in the top plot and red dots in all plots, indicate time points at which agonist and buffer were introduced. **B.** Plots of PH-mCh emission intensity for cells expressing different  $\gamma$  subunit types (as indicated). Basal level of mCh intensity was treated as 100%. doi:10.1371/journal.pone.0007797.g003




**Figure 4. Response of a  $\gamma 9-3$  chimera.** A. C terminal sequences of relevant  $\gamma$  subtypes. B. Representative images and plots of  $\gamma 9-3$  chimera and PH-mCh translocation. Experiments and analysis were performed as described in Fig. 3A. C. Bar diagrams of the extent of PH-mCh translocation at different concentrations of agonist in cells expressing indicated  $\gamma$  subunits. Values were calculated as in Fig. 1B D. Ratio of the 100  $\mu\text{M}$  over 1  $\mu\text{M}$  response. Results are represented as mean  $\pm$  SEM (N = no  $\gamma$ -51,  $\gamma 3$ -52,  $\gamma 9$ -54,  $\gamma 9-3$  -40). \*\* $p < 0.001$  ( $\gamma 9$  versus others at the same concentration). doi:10.1371/journal.pone.0007797.g004

heterotrimer there is FRET from CFP to YFP leading to diminished emission from CFP. When the G protein heterotrimer is activated by an active GPCR, if there is dissociation of  $G\alpha$  and  $G\beta\gamma$  then the FRET is lost leading to restoration of CFP intensity which is marked by an increase in its fluorescence emission. We observed the FRET signal in live M3-CHO cells expressing  $\alpha\text{q}$ -CFP and YFP- $\gamma 3$  or YFP- $\gamma 9$ . Only cells expressing comparable intensities of CFP and YFP proteins were selected for observation. FRET experiments were performed as described before [14,19]. Essentially, FRET was determined by monitoring gain in CFP emission intensity in the plasma membrane by photobleaching YFP (acceptor photobleaching [20]) in both the basal and agonist activated states. We did not detect any FRET from  $\alpha\text{q}$ -CFP YFP- $\gamma 3$  in the basal state (data not shown) while  $\alpha\text{q}$ -CFP YFP- $\gamma 9$  did provide a clearly detectable FRET signal. To ensure that the

lack of detectable FRET between  $\alpha\text{q}$ -CFP and  $\beta\text{YFP-}\gamma 3$  is not due to the inability of these subunits to form a heterotrimer, we examined the ability of  $\alpha\text{q}$ -CFP to support the translocation of a  $\gamma 3$  mutant capable of translocation [7]. M3 activation of M3-CHO cells expressing  $\alpha\text{q}$ -CFP and the  $\gamma 3$  mutant resulted in its translocation from the plasma membrane to internal membranes (Fig. S1). This result indicated that  $\alpha\text{q}$ -CFP and  $\beta\text{YFP-}\gamma 3$  are capable of forming a heterotrimer. FRET does not occur between the tagged FPs likely due to five extra residues in the N terminal domain of  $\gamma 3$  compared to  $\gamma 9$  (Fig. 5A). YFP fused to the  $\gamma$  subunit N terminus maybe in a non-optimal orientation with reference to CFP resulting in lack of FRET between  $\alpha\text{q}$ -CFP and YFP- $\gamma 3$  [20].

To obtain an alternate FRET sensor containing a non-translocating  $\gamma$  subunit we examined the amino acid sequences



**Figure 5. Differential effector and FRET responses of cells expressing different  $\gamma$  subunits.** A. Primary structures of  $\gamma 3$ ,  $\gamma 2$  and  $\gamma 9$ . Comparison shows that  $\gamma 3$  has five additional residues at the N terminus compared to  $\gamma 9$  while  $\gamma 2$  has only one (highlighted residues). B. PH-mCh translocation induced by the sequential addition of agonist from cells expressing different  $\gamma$  subunit types (as indicated). N = 11. Results are the means  $\pm$  SEM. \* $p < 0.01$ , \*\* $p < 0.001$ . C. A FRET sensor based on  $\alpha q$ -CFP and  $\beta\gamma$ -YFP. Representation of FRET sensor (left). Bars (right) show increase in CFP emission on the plasma membrane as a result of FRET loss in presence of various  $\gamma$  subunits at basal (black bars) and activated (white bars) state determined as in Methods and Text S1. Results are the means  $\pm$  SEM; N = 9, \*\* $p < 0.001$ . doi:10.1371/journal.pone.0007797.g005

of  $\gamma$  subunits and found that the  $\gamma 2$  sequence has only one extra residue in the N terminal domain compared to  $\gamma 9$  (Fig. 5A). We first examined whether the properties of  $\gamma 2$  were similar to  $\gamma 3$  by measuring the extent of PH-mCh translocation as mentioned above. The profile for PH domain translocation with  $\gamma 2$  (Fig. 5B and Fig. S2) was similar to that obtained with other non-translocating  $\gamma$  subunits ( $\gamma 3$  and  $\gamma 9$ -3 chimera). We then examined the FRET signal in the basal and agonist activated states from  $\alpha q$ -CFP to YFP- $\gamma 2$  or YFP- $\gamma 9$  in M3-CHO cells (Fig. 5C). In the basal state, a FRET signal was detected as a CFP increase after YFP photobleaching with both of subunits (Fig. 5C, black bars). On receptor stimulation by 100  $\mu$ M carbachol, the FRET signal decreased significantly in  $\gamma 9$  cells but not in cells expressing  $\gamma 2$  (Fig. 5C, white bars). The dissociation of  $\beta\gamma 9$  from  $\alpha q$  and its translocation away from the plasma membrane explains the reduction in the FRET signal. In surprising contrast, in the presence of  $\gamma 2$  the FRET signal did not change after receptor activation suggesting that either there is no dissociation of the heterotrimer or that heterotrimer reassociation is very rapid after dissociation so that it is not detected under the conditions used here. The differences in the translocation capabilities of  $\gamma$  subunit types thus have strikingly different impacts on heterotrimer dissociation after receptor stimulation.

These results are consistent with our model (Fig. 2) that the translocation of the  $\beta\gamma$  complex results in slower activation (cycling) of the  $\alpha$  subunit while in the case of heterotrimers containing the non-translocating  $\gamma$  subunit, significant dissociation does not occur allowing much higher levels of activation by a receptor because the G protein in the heterotrimer state is available continually.

#### DAG responses in cells expressing translocating or non-translocating $\gamma$ subunits are similar to IP3

To ensure that the effects seen above by measuring IP3 levels using Ph-mCh was reflected in the generation of a different second


messenger, DAG similarly in M3-CHO cells containing expressing YFP-DBD and CFP- $\gamma 11$  or CFP- $\gamma 2$ . Cells were activated with 1  $\mu$ M carbachol as above.  $\gamma 11$  has identical translocation properties to  $\gamma 9$  (Fig. S3) and was used in this case because CFP- $\gamma 11$  was available, has been characterized previously and allowed us to examine whether translocation capable subunits had similar effects [6]. CFP- $\gamma 11$  translocated strongly on M3 receptor activation demonstrating that the cells contained  $\alpha q$  (Fig. S3). When M3 was activated, in cells expressing  $\gamma 2$ , YFP-DBD translocated to the plasma membrane on activation of the receptor (Fig. 6). In contrast, cells expressing  $\gamma 11$  showed a weak response (Fig. 6). These results showed that regardless of the second messenger measured, cells expressing different  $\gamma$  subunits have differential responses.

#### Endogenous $\gamma$ subunit constitution influences signaling activity

We then examined whether cells that expressed an endogenous translocation proficient  $\gamma$  subunit showed differential response to cells not expressing any translocation proficient  $\gamma$  subunit. To pursue these experiments we used an  $\alpha q$  subunit tethered to the M3 receptor to obtain equimolar concentrations of expressed receptor and  $\alpha q$ . The N terminal of the  $\alpha$  subunit was fused to the C terminus of the M3 receptor.

We then confirmed the functionality of M3- $\alpha q$ -CFP fusion protein. The fusion protein visibly supported PH - domain translocation on activation by a M3 receptor agonist indicative of its ability to activate downstream signaling pathways like the wild type M3 and G $\alpha q$  (Fig. 7).

We then evaluated PH-mCh translocation as above to determine the effector activation abilities of M3- $\alpha q$ -CFP in the presence of YFP- $\gamma 2$  or  $\gamma 9$  in CHO cells. We ensured that the CFP and YFP emission intensities on the plasma membrane were similar between the  $\gamma 2$  and  $\gamma 9$  coexpressing cells (Table S1). Fig. 7A shows that in the presence of  $\gamma 2$ , which does not effectively



**Figure 6. YFP-DBD translocation in the presence of various  $\gamma$  subunits.** Representative images and bar diagrams of YFP-DBD translocation are shown in the presence of 1  $\mu\text{M}$  carbachol. Values were calculated as in Fig. 1B. Images of cells expressing  $\alpha\text{q}$  with CFP- $\gamma 2$  (N=9) or CFP- $\gamma 11$  (N=10) (as indicated) and YFP-DBD were acquired as described in Methods and Text S1. Values were calculated as ratio of M-B/B as described in Fig. 1B legend. Results are the means  $\pm$  SEM. \*\* $p < 0.001$ . doi:10.1371/journal.pone.0007797.g006

dissociate on receptor activation, the effector activation is very limited compared to the strong response in cells expressing the translocation capable,  $\gamma 9$ . These results were consistent with the conformationally restricted  $\alpha\text{q}$  not activating an effector when bound to  $\beta\gamma 2$  complex. Translocation of  $\beta\gamma 9$  allows the tethered  $\alpha\text{q}$  to access PLC  $\beta$  as described in the model in Fig. 8.

These results allowed us to examine whether a cell line which contains no detectable levels of a translocating  $\gamma$  subunit (CHO) or one that contains a translocating  $\gamma$  subunit (A549) demonstrate properties that are consistent with a role for these subunits in the regulation of signaling sensitivity in a cell. We screened a variety of cell lines for the presence of endogenous translocating  $\gamma$  subunits by checking for the potential cotranslocation of YFP- $\beta 1$  introduced into these cells. We have previously shown that YFP- $\beta 1$  translocates only when bound to a translocation capable  $\gamma$  subunit [7]. Using this approach we identified a lung epithelial fibroblast cell line, A549 containing translocation proficient  $\gamma$  subunits (manuscript submitted). Consistent with the ability of YFP- $\beta 1$  to cotranslocate in A549 cells, qRT-PCR showed that these cells contain significant levels of translocation proficient  $\gamma 11$ ,  $\gamma 10$  and  $\gamma 5$  subunits and only one non-translocating  $\gamma$  subunit,  $\gamma 12$  (Fig. S4). In contrast to A549 cells, YFP- $\beta 1$  did not cotranslocate in CHO cells (data not shown) suggesting that they do not contain translocating  $\gamma$  subunits at concentrations that are likely to have an effect on downstream activity. We examined these two cell types in experiments similar to those above.

When M3- $\alpha\text{q}$ -CFP was activated in CHO cells with 100  $\mu\text{M}$  carbachol, almost no response was detected (Fig. 7B) consistent with a non-dissociating  $\beta\gamma$  preventing the conformationally restricted  $\alpha\text{q}$  from acting on an effector. However, in A549 cells a strong response was detected consistent with the dissociation and translocation of an endogenous translocating  $\beta\gamma$  complex allowing  $\alpha\text{q}$  to access PLC $\beta$  (Fig. 7B). These results were consistent with the model shown in Fig. 8. They also show that the endogenous  $\gamma$  subunit constitution can regulate the sensitivity of a cell's response to an external stimulus and the results here are not peculiar to the introduced  $\gamma$  subunits.


## Discussion

We have used live cell imaging of single mammalian cells to identify mechanisms at the basis of the sensitivity of cellular responses to G protein activation by an extracellular signal. We show here that the ability of G protein  $\gamma$  subunit types to translocate away from the plasma membrane on receptor activation controls sensitivity of response. Cells that express a translocating  $\gamma$  subunit demonstrate attenuated PIP2 breakdown in response to M3 receptor stimulation compared to cells that express a non translocating  $\gamma$  subunit. The lowered sensitivity is not due to a specific property of a  $\gamma$  subunit type but due to the translocation of the  $\beta\gamma$  complex because cells expressing a translocation capable  $\gamma$  subunit mutated to abolish translocation showed heightened sensitivity similar to non translocating  $\gamma$  subunits.

We examined the basis of this differential effect of translocating and non-translocating  $\gamma$  subunits using a FRET sensor based on  $\alpha\text{q}$ -CFP and YFP- $\gamma$  interaction. The Gq sensor behaved similar to a Go sensor when it contained  $\gamma 9$ , demonstrating a significant loss in FRET on receptor activation. But surprisingly with a non-translocating  $\gamma$  subunit,  $\gamma 2$ , there was no loss in FRET on receptor activation. The lack of  $\beta\gamma 2$  dissociation from  $\alpha\text{q}$  is consistent with a long standing suggestion that the G protein heterotrimer does not have to dissociate to act on effectors [21,22] and recent reports of the inability of receptor activation to dissociate  $\alpha\text{q}$ ,  $\alpha\text{i-1}$ ,  $\alpha\text{z}$ ,  $\alpha\text{-cone}$  and  $\alpha\text{s-q}$  chimera from  $\beta 1\gamma 2$  [23,24,25,26,27]. This result is also consistent with our previous finding that  $\beta\gamma$  translocation is influenced by the associated  $\alpha$  subunit type [28]. The lack of dissociation of  $\alpha\text{q}\beta\gamma 2$  (or very brief dissociation) allows continual access to a receptor, a higher level of activation of  $\alpha\text{q}$  and high level of PLC $\beta$  activation and PIP2 breakdown (Fig. 2). In contrast, in the case of  $\alpha\text{q}\beta\gamma 9$ , translocation of  $\beta\gamma$  results in reduction of heterotrimer available on the plasma membrane for PLC $\beta$  activation resulting in lowered PIP2 breakdown (Fig. 2).

To further examine the role of subunit dissociation and translocation on the sensitivity of the cell's response to an agonist






**Figure 7. Effect of different  $\gamma$  subunits on M3- $\alpha$ q-CFP induced response.** A. Representative images of cells expressing M3- $\alpha$ q-CFP with YFP- $\gamma$ 2 (N = 17) or YFP- $\gamma$ 9 (N = 9) (as indicated) and PH-mCh. Bar diagrams of PH-mCh translocation are shown in the presence of 100  $\mu$ M carbachol. Values were calculated as in Fig. 1B. Results are the means  $\pm$  SEM. \*\* $p < 0.001$ . B. PH-mCh translocation induced by M3- $\alpha$ q-CFP fusion in cell lines containing endogenous non-translocating and translocating  $\gamma$  subunits. CHO (N = 11) and A549 (N = 13) cells expressing M3- $\alpha$ q-CFP, PH-mCh and no  $\gamma$  subunits. Values were calculated as in Fig. 1B. Results are the means  $\pm$  SEM. \*\* $p < 0.001$ . doi:10.1371/journal.pone.0007797.g007

we tethered the  $\alpha$  subunit to the receptor so that it is conformationally constrained. Similar receptor  $\alpha$  subunit fusions have previously been shown to be capable of functioning effectively on receptor activation [29]. PIP2 breakdown was strong in cells containing the M3- $\alpha$ q-CFP protein and translocating  $\gamma$ 9 and weak in cells expressing non-translocating  $\gamma$ 2. The results support the model in Fig. 8. Thus tethered  $\alpha$ q is unable to access PLC $\beta$  when bound to a non dissociating  $\beta\gamma$  complex. In contrast, in the presence of the translocating  $\beta\gamma$  subunits, the tethered  $\alpha$ q is able to activate PLC $\beta$  because it is accessible to PLC $\beta$  after  $\beta\gamma$  dissociation and translocation.

While mechanisms that desensitize a cell's response to external stimuli are well characterized at the level of the receptor [30], few mechanisms that act at the level of the G protein to attenuate or accentuate the response to a signal have been identified. In the *Drosophila* compound eye, DGq $\alpha$ , translocates from the rhabdome, the membrane structure containing the phototransduction machinery, to the cell body (Golgi/ER) upon rhodopsin activation with a time course of a few minutes and reverse translocates on rhodopsin inactivation with a time course of tens of minutes [31]. This translocation of DGq $\alpha$  has been shown to regulate the

sensitivity of the phototransduction cascade by reducing the amount of DGq $\alpha$  following illumination [32]. In rod photoreceptors, translocation of Gt subunits from outer to inner segment on rhodopsin activation has been shown [5,33]. This translocation has been shown to help adapt to the light signal by attenuating the response [33,34]. It is of interest that the G protein in the rods contains  $\gamma$ 1 that we have previously shown to be capable of translocation when activated by a GPCR [7]. However, there are several striking distinctions between the properties noted for translocation in rod photoreceptors and the translocation of the G protein  $\beta\gamma$  complex examined here. Activated rhodopsin induces the translocation of both the  $\alpha$  and  $\beta\gamma$  subunits of Gt unlike the translocation examined here. The translocation of Gt subunits also occurs at a much slower rate than the selective  $\beta\gamma$  translocation studied here ( $t_{1/2}$ :  $\alpha$   $\sim$ 5 min and  $\beta\gamma$   $\sim$ 12 min [33] in contrast to  $\beta\gamma$ 9  $\sim$ 10 sec [7]). Additionally, the reversal of G protein subunit translocation in the rods occurs much slower than its forward translocation (200 min for  $\alpha$ ) [35] compared to  $\beta\gamma$ 9 which reverses at the same rate as forward translocation [7]. Finally, there is no equivalent in rod photoreceptors to the heightened response seen in the case of non-translocating  $\gamma$



**Figure 8. Model for the basis of differential responses to the activation of a tethered  $\alpha_q$  subunit.** The tethered  $\alpha_q$  subunit when bound to a non-translocating  $\gamma$  subunit is unable to activate PLC $\beta$  after receptor activation because of its continued association with  $\beta\gamma$  and the inflexibility of the  $\alpha$  subunit that is fused to the C terminus of the receptor. The effector activating surface of the  $\alpha$  subunit and the  $\beta\gamma$  complex do not become accessible to PLC  $\beta$  in this situation. In contrast, when the tethered  $\alpha_q$  subunit is bound to a  $\gamma$  subunit that is capable of translocation away from the plasma membrane on receptor activation, the effector activating surface of the  $\alpha$  subunit is exposed to PLC  $\beta$  allowing it to activate the effector molecule.

doi:10.1371/journal.pone.0007797.g008

subunits. It is however possible that light adaptation conferred by the translocation of G protein subunits in rod photoreceptors is evolutionarily related to the attenuation of effector response seen here.

An important question is whether the results we have seen are of significance to the sensitivity of response of mammalian cells into which  $\gamma$  subunits have not been introduced. In CHO cells that do not contain a translocating  $\gamma$  subunit and in A549 cells that contain translocating  $\gamma$  subunits, the M3 tethered  $\alpha_q$  induced distinctly different PIP<sub>2</sub> breakdown responses that were consistent with the responses seen in cells into which non-translocating  $\gamma_2$  or translocating  $\gamma_9$  subunits were introduced. These results showed that endogenous  $\gamma$  subunit type constitution of cells can regulate the sensitivity of a cell's response to an external stimulus.

Previously we have shown that a FRET based sensor made up of  $\alpha_o$  and  $\beta\gamma_2$  shows FRET loss on receptor activation [14] suggesting that non-translocating  $\beta\gamma$ s dissociate from some  $\alpha$  subunits. In contrast, the results here suggest that  $\alpha_q$  does not dissociate from  $\beta\gamma_2$  on activation. Thus two different factors appear to control the concentration of heterotrimer available to the activated receptor. The  $\alpha$  subunit type which affects dissociation and the  $\gamma$  subunit type which regulates translocation and consequently dissociation. Signaling responses can thus be controlled by the  $\alpha$  and  $\gamma$  subunit type constitution of a cell. Consistent with this prediction, we have previously reported that in a receptor reconstituted system, both the  $\alpha$  subunit type and the  $\gamma$  subunit type affected the receptor stimulated activity of G proteins [36].

G proteins control the vast majority of signaling pathways in mammalian cells and the existence of diverse evolutionarily conserved families of G protein  $\alpha$  and  $\gamma$  subunits has been known for many years [1]. But the roles that these subunit types play in the dynamic environment of living cells have not been fully explored [37]. While there has been evidence for the specificity of interaction between G protein  $\alpha$  and  $\gamma$  subunits with receptors [38,39], their roles in regulating the kinetics of signaling in

mammalian cells has been poorly defined. The results here identify mechanisms that allow the existent diversity of G protein subunits in a cell to control the intensity of response to varying levels of external stimuli. Overall, these results suggest that G protein  $\beta\gamma$  complex translocation and dissociation of the heterotrimer are novel mechanisms that can control the intensity of response of a cell to receptor stimulation.

## Materials and Methods

### Expression constructs and cell culture

$\gamma$  subunits were tagged with YFP or CFP as described previously [6,7]. The chimera  $\gamma_9$ -3 was made by substitution of the last 15 amino acids of the C terminus from  $\gamma_3$  in  $\gamma_9$  subunit.  $\alpha_q$ -GFP (from C. Berlot, Geisinger Clinic) was mutated to obtain  $\alpha_q$ -CFP and was transferred to pcDNA3.1. PH domain of PLC  $\delta$  was obtained as PH-EGFP [13] (from T. Balla, NIH) and GFP was substituted with mCh (from R. Tsien, UC San Diego). YFP-DBD was obtained from A. Newton, UC San Diego [11].

CHO cells stably expressing the M3 muscarinic receptor (M3-CHO) have been described previously [14]. CHO cells were grown in CHO IIIa medium (Invitrogen) containing 10% dialyzed fetal bovine serum (Atlanta Biologicals), methotrexate, penicillin, streptomycin, and glutamine. A549 cells were grown in Ham's F12 (Mediatech) with 10% dialyzed fetal bovine serum, penicillin and streptomycin. All the transfections were performed using Lipofectamine 2000 (Invitrogen) as described previously [19] (additional details are in Text S1).

### Live cell imaging

Briefly, transfected cells cultured on glass coverslips were imaged in a chamber with a 25  $\mu$ l volume using a Zeiss Axioskop microscope with a 63x objective (1.4 NA), 100 W mercury arc lamp and Hamamatsu CCD Orca-ER camera. There was programmed delivery of solutions at 0.5 ml/min flow rate using an automated delivery system (Automate Scientific). For FRET

experiments, cells with equal expression levels of CFP and YFP were selected and experiments were performed as previously described [14]. Chromatic filters and other details are in Text S1.

## Supporting Information

**Figure S1** M3-CHO cells were co-transfected with  $\alpha_q$ -CFP and YFP- $\gamma 3$  C-terminal mutant. M3 activation induces translocation of the  $\gamma 3$  mutant. Images from transfected cells were acquired with 10 sec intervals. Cells were exposed to 100  $\mu$ M carbachol (agonist) followed by a wash with buffer at the indicated time points (as shown by arrows in plot). YFP emission intensity changes over time in Golgi (white arrow) were plotted.  
Found at: doi:10.1371/journal.pone.0007797.s001 (0.02 MB PDF)

**Figure S2** M3-CHO cells were co-transfected with  $\alpha_q$ -CFP, YFP- $\gamma 2$  and PH-mCherry. Images from transfected cells were acquired with 10 sec interval. Cells were exposed sequentially to 1  $\mu$ M and 100  $\mu$ M carbachol (agonist) followed by a wash with buffer at the indicated time points (as shown by arrows in plot). YFP changes over time in Golgi and mCherry emission intensity in cytosol (as indicated by black circles) were plotted.  
Found at: doi:10.1371/journal.pone.0007797.s002 (0.03 MB PDF)

**Figure S3** M3-CHO cells were co-transfected with  $\alpha_q$ , CFP- $\gamma 11$  and YFP-DBD. Images from transfected cells were acquired with 10 sec interval. Cells were exposed to 1  $\mu$ M carbachol (agonist)

followed by a wash with buffer. As shown in these images, CFP- $\gamma 11$  translocates with the same efficiency as YFP- $\gamma 9$ . Images of YFP-DBD are shown in main text Fig. 6 (same cell).  
Found at: doi:10.1371/journal.pone.0007797.s003 (0.01 MB PDF)

**Figure S4** Real time PCR analysis of the relative expression levels of the G protein gamma subunit family  
Found at: doi:10.1371/journal.pone.0007797.s004 (0.11 MB PDF)

**Table S1** CFP emission intensities are not correlated with cell responses  
Found at: doi:10.1371/journal.pone.0007797.s005 (0.02 MB PDF)

**Text S1** Materials and Methods  
Found at: doi:10.1371/journal.pone.0007797.s006 (0.04 MB PDF)

**Text S2** Effect of G protein  $\beta\gamma$  on PLC $\beta$  activation  
Found at: doi:10.1371/journal.pone.0007797.s007 (0.01 MB PDF)

## Author Contributions

Conceived and designed the experiments: MC DKS NG. Performed the experiments: MC DKS JHC. Analyzed the data: MC DKS JHC NG. Contributed reagents/materials/analysis tools: VK. Wrote the paper: MC DKS NG.

## References

- Simon MI, Strathmann MP, Gautam N (1991) Diversity of G proteins in signal transduction. *Science* 252: 802–808.
- Oldham WM, Hamm HE (2008) Heterotrimeric G protein activation by G-protein-coupled receptors. *Nat Rev Mol Cell Biol* 9: 60–71.
- Yi TM, Kitano H, Simon MI (2003) A quantitative characterization of the yeast heterotrimeric G protein cycle. *Proc Natl Acad Sci U S A* 100: 10764–10769.
- Suh BC, Horowitz LF, Hirdes W, Mackie K, Hille B (2004) Regulation of KCNQ2/KCNQ3 current by G protein cycling: the kinetics of receptor-mediated signaling by Gq. *J Gen Physiol* 123: 663–683.
- Arshavsky VY, Lamb TD, Pugh EN Jr (2002) G proteins and phototransduction. *Annu Rev Physiol* 64: 153–187.
- Akgoz M, Kalyanaraman V, Gautam N (2004) Receptor-mediated reversible translocation of the G protein betagamma complex from the plasma membrane to the Golgi complex. *J Biol Chem* 279: 51541–51544.
- Saini DK, Kalyanaraman V, Chisari M, Gautam N (2007) A family of G protein betagamma subunits translocate reversibly from the plasma membrane to endomembranes on receptor activation. *J Biol Chem* 282: 24099–24108.
- Rhee SG (2001) Regulation of phosphoinositide-specific phospholipase C. *Annu Rev Biochem* 70: 281–312.
- Zhang J, Campbell RE, Ting AY, Tsien RY (2002) Creating new fluorescent probes for cell biology. *Nat Rev Mol Cell Biol* 3: 906–918.
- Sakai N, Sasaki K, Ikegaki N, Shirai Y, Ono Y, et al. (1997) Direct visualization of the translocation of the gamma-subspecies of protein kinase C in living cells using fusion proteins with green fluorescent protein. *J Cell Biol* 139: 1465–1476.
- Gallegos LL, Kunkel MT, Newton AC (2006) Targeting protein kinase C activity reporter to discrete intracellular regions reveals spatiotemporal differences in agonist-dependent signaling. *J Biol Chem* 281: 30947–30956.
- Horowitz LF, Hirdes W, Suh BC, Hilgemann DW, Mackie K, et al. (2005) Phospholipase C in living cells: activation, inhibition, Ca<sup>2+</sup> requirement, and regulation of M current. *J Gen Physiol* 126: 243–262.
- Varnai P, Balla T (1998) Visualization of phosphoinositides that bind pleckstrin homology domains: calcium- and agonist-induced dynamic changes and relationship to myo-[<sup>3</sup>H]inositol-labeled phosphoinositide pools. *J Cell Biol* 143: 501–510.
- Azpiuz I, Gautam N (2004) A Fluorescence Resonance Energy Transfer-based Sensor Indicates that Receptor Access to a G Protein Is Unrestricted in a Living Mammalian Cell. *J Biol Chem* 279: 27709–27718.
- Hein P, Frank M, Hoffmann C, Lohse MJ, Bunemann M (2005) Dynamics of receptor/G protein coupling in living cells. *Embo J* 24: 4106–4114.
- Rebois RV, Robitaille M, Gales C, Dupre DJ, Baragli A, et al. (2006) Heterotrimeric G proteins form stable complexes with adenylyl cyclase and Kir3.1 channels in living cells. *J Cell Sci* 119: 2807–2818.
- Riven I, Iwanir S, Reuveni E (2006) GIRK channel activation involves a local rearrangement of a preformed G protein channel complex. *Neuron* 51: 561–573.
- Akgoz M, Kalyanaraman V, Gautam N (2006) G protein betagamma complex translocation from plasma membrane to Golgi complex is influenced by receptor gamma subunit interaction. *Cell Signal* 18: 1758–1768.
- Chisari M, Saini DK, Kalyanaraman V, Gautam N (2007) Shuttling of G protein subunits between the plasma membrane and intracellular membranes. *J Biol Chem* 282: 24092–24098.
- Sekar RB, Periasamy A (2003) Fluorescence resonance energy transfer (FRET) microscopy imaging of live cell protein localizations. *J Cell Biol* 160: 629–633.
- Klein S, Reuveni H, Levitzki A (2000) Signal transduction by a nondissociable heterotrimeric yeast G protein. *Proc Natl Acad Sci U S A* 97: 3219–3223.
- Levitzki A (1986) Beta-adrenergic receptors and their mode of coupling to adenylate cyclase. *Physiol Rev* 66: 819–854.
- Evanko DS, Thiyagarajan MM, Takida S, Wedegaertner PB (2005) Loss of association between activated G $\alpha_q$  and Gbetagamma disrupts receptor-dependent and receptor-independent signaling. *Cell Signal* 17: 1218–1228.
- Digby GJ, Lober RM, Sethi PR, Lambert NA (2006) Some G protein heterotrimers physically dissociate in living cells. *Proc Natl Acad Sci U S A* 103: 17789–17794.
- Frank M, Thumer L, Lohse MJ, Bunemann M (2005) G Protein activation without subunit dissociation depends on a G $\{\alpha\}$ -specific region. *J Biol Chem* 280: 24584–24590.
- Gales C, Van Durm JJ, Schaak S, Pontier S, Percherancier Y, et al. (2006) Probing the activation-promoted structural rearrangements in preassembled receptor-G protein complexes. *Nat Struct Mol Biol* 13: 778–786.
- Rosenzweig DH, Nair KS, Wei J, Wang Q, Garwin G, et al. (2007) Subunit dissociation and diffusion determine the subcellular localization of rod and cone transducins. *J Neurosci* 27: 5484–5494.
- Azpiuz I, Akgoz M, Kalyanaraman V, Gautam N (2006) G protein betagamma11 complex translocation is induced by Gi, Gq and Gs coupling receptors and is regulated by the alpha subunit type. *Cell Signal* 18: 1190–1200.
- Milligan G (2002) The use of receptor G-protein fusion proteins for the study of ligand activity. *Receptors Channels* 8: 309–317.
- Gainetdinov RR, Premont RT, Bohn LM, Lefkowitz RJ, Caron MG (2004) Desensitization of G protein-coupled receptors and neuronal functions. *Annu Rev Neurosci* 27: 107–144.
- Kosloff M, Elia N, Joel-Almagor T, Timberg R, Zars TD, et al. (2003) Regulation of light-dependent G $\alpha_q$  translocation and morphological changes in fly photoreceptors. *EMBO J* 22: 459–468.
- Frechter S, Elia N, Tzarfaty V, Selinger Z, Minke B (2007) Translocation of Gq alpha mediates long-term adaptation in Drosophila photoreceptors. *J Neurosci* 27: 5571–5583.
- Calvert PD, Strissel KJ, Schiesser WE, Pugh EN Jr, Arshavsky VY (2006) Light-driven translocation of signaling proteins in vertebrate photoreceptors. *Trends Cell Biol* 16: 560–568.

34. Kassai H, Aiba A, Nakao K, Nakamura K, Katsuki M, et al. (2005) Farnesylation of retinal transducin underlies its translocation during light adaptation. *Neuron* 47: 529–539.
35. Elias RV, Sezate SS, Cao W, McGinnis JF (2004) Temporal kinetics of the light/dark translocation and compartmentation of arrestin and alpha-transducin in mouse photoreceptor cells. *Mol Vis* 10: 672–681.
36. Hou Y, Azpiazu I, Smrcka A, Gautam N (2000) Selective role of G protein gamma subunits in receptor interaction. *J Biol Chem* 275: 38961–38964.
37. Saini DK, Chisari M, Gautam N (2009) Shuttling and translocation of heterotrimeric G proteins and Ras. *Trends Pharmacol Sci* 30: 278–286.
38. Gautam N (2003) A conformational switch regulates receptor-g protein interaction. *Structure (Camb)* 11: 359–360.
39. Neves SR, Ram PT, Iyengar R (2002) G protein pathways. *Science* 296: 1636–1639.