

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2011

Prevalence of nasal carriage of mupirocin-resistant Staphylococcus aureus among hospitalized patients in Thailand

Anucha Apisarnthanarak
Thammasat University Hospital

David K. Warren
Washington University School of Medicine in St. Louis

Victoria J. Fraser
Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs



Part of the [Medicine and Health Sciences Commons](#)

Please let us know how this document benefits you.

Recommended Citation

Apisarnthanarak, Anucha; Warren, David K.; and Fraser, Victoria J., "Prevalence of nasal carriage of mupirocin-resistant *Staphylococcus aureus* among hospitalized patients in Thailand." *Infection Control and Hospital Epidemiology*. 32, 5. 522-523. (2011).

https://digitalcommons.wustl.edu/open_access_pubs/796

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.



CHICAGO JOURNALS



Prevalence of Nasal Carriage of Mupirocin-Resistant *Staphylococcus aureus* among Hospitalized Patients in Thailand •

Author(s): Anucha Apisarnthanarak, David K. Warren, Victoria J. Fraser

Reviewed work(s):

Source: *Infection Control and Hospital Epidemiology*, Vol. 32, No. 5 (May 2011), pp. 522-523

Published by: [The University of Chicago Press](#) on behalf of [The Society for Healthcare Epidemiology of America](#)

Stable URL: <http://www.jstor.org/stable/10.1086/659953>

Accessed: 06/03/2012 23:07

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



The University of Chicago Press and The Society for Healthcare Epidemiology of America are collaborating with JSTOR to digitize, preserve and extend access to Infection Control and Hospital Epidemiology.

<http://www.jstor.org>

- multiple-locus variable-number tandem-repeat analysis to determine clonal spread of toxin A–negative *Clostridium difficile* in a general hospital in Buenos Aires, Argentina. *Clin Microbiol Infect* 2009;15:1080–1086.
3. van den Berg RJ, Claas EC, Oyib DH, et al. Characterization of toxin A–negative, toxin B–positive *Clostridium difficile* isolates from outbreaks in different countries by amplified fragment length polymorphism and PCR ribotyping. *J Clin Microbiol* 2004; 42(3):1035–1041.
 4. Gerding DN. Global epidemiology of *Clostridium difficile* infection in 2010. *Infect Control Hosp Epidemiol* 2010;31(suppl 1):S32–S34.
 5. Barbut F, Lalande V, Burghoffer B, Thien HV, Grimprel E, Petit JC. Prevalence and genetic characterization of toxin A variant strains of *Clostridium difficile* among adults and children with diarrhea in France. *J Clin Microbiol* 2002;40:2079–2083.
 6. Kato H, Kato N, Watanabe K, et al. Identification of toxin A–negative, toxin B–positive *Clostridium difficile* by PCR. *J Clin Microbiol* 1998;36:2178–2182.
 7. Geric B, Rupnik M, Gerding DN, Grabnar M, Johnson S. Distribution of *Clostridium difficile* variant toxinotypes and strains with binary toxin genes among clinical isolates in an American hospital. *J Med Microbiol* 2004;53(9):887–894.
 8. Rupnik M, Kato N, Grabnar M, Kato H. New types of toxin A–negative, toxin B–positive strains among *Clostridium difficile* isolates from Asia. *J Clin Microbiol* 2003;41(3):1118–1125.
 9. Kato H, Kato N, Katow S, Maegawa T, Nakamura S, Lyerly DM. Deletions in the repeating sequences of the toxin A gene of toxin A–negative, toxin B–positive *Clostridium difficile* strains. *FEMS Microbiol Lett* 1999;175:197–203.

Prevalence of Nasal Carriage of Mupirocin-Resistant *Staphylococcus aureus* among Hospitalized Patients in Thailand

To the Editor—In various parts of the world, mupirocin has been used as a component of a “search and destroy” strategy to prevent methicillin-resistant *Staphylococcus aureus* (MRSA) infection prior to a surgical procedure.^{1–3} Increased use of mupirocin has been associated with an increased prevalence of mupirocin-resistant MRSA.^{4,5} In Thailand, there is neither routine surveillance for nasal carriage of MRSA nor attempts to eradicate carriage among hospitalized inpatients. Given that mupirocin and antibiotics can be purchased at outpatient pharmacies in Thailand without a prescription,⁶ it is expected that the prevalence of mupirocin-resistant MRSA nasal carriage is high. We performed a point prevalence study to evaluate the prevalence of MRSA nasal carriage and to estimate the proportion of individuals with mupirocin-resistant MRSA nasal carriage among admitted inpatients at Thammasat University Hospital.

From January 1 to January 7, 2010, we cultured anterior nares swab specimens to detect MRSA carriage. Consecutive hospitalized patients who consented to the study participation

on day 1 of hospitalization provided swab samples from both anterior nares that were then cultured for presence of MRSA. Information regarding demographics and clinical history of previous admission and exposure to antibiotics (in both inpatient and outpatient settings) was collected from each patient. Detection of MRSA from swab specimens was performed as previously described.⁴ Samples containing MRSA isolates from all patients who tested positive for MRSA carriage were then subcultured to BBL trypticase soy agar with 5% sheep blood (BD Diagnostics) and incubated at 35°C for 24 hours. Colonies isolated from the BBL trypticase soy agar were then inoculated on a Mueller Hinton II plate, and a mupirocin Etest (AB Biodisk) strip was applied. After 24 hours of incubation at 35°C, the minimum inhibitory concentration (MIC) was read. Isolates were classified as susceptible (MIC, <8 µg/mL), low-level resistant (MIC, 8–256 µg/mL), or high-level resistant (MIC, 512 µg/mL).

During the study period, 250 (86%) of 290 adult patients who were admitted to all units at Thammasat University Hospital provided consent and were enrolled in the study; 149 (60%) were male, and the median age was 45 years (range, 20–89 years). Nasal carriage of MRSA was detected in 9 patients (3.6%). Four (44%) of these 9 patients were admitted to medical units, 3 (33%) were admitted to surgical units, and 2 (23%) were admitted to an orthopedic unit. All patients with nasal carriage of MRSA had a history of hospital admission within the past 12 months. Low-level mupirocin-resistant MRSA was detected in 2 patients (22%). These 2 patients had a history of skin abrasions and had self-purchased mupirocin and fluoroquinolones from a local pharmacy within 3 months of admission to the hospital.

Our data suggest that MRSA nasal carriage occurred among patients who had a history of contact with the Thai health care system, and that injudicious use of mupirocin in the community may be associated with carriage of low-level mupirocin-resistant MRSA isolates. The relatively low prevalence of MRSA nasal carriage (3.6%) among hospitalized patients in our study suggests that a search and destroy strategy may not be cost-effective to implement in this middle-income country, while the relatively high prevalence of mupirocin-resistant MRSA (22%) among patients with nasal carriage of MRSA suggests the need to employ an antimicrobial stewardship program at the community level to help limit the unnecessary use of mupirocin and other antibiotics. Further studies are needed to evaluate the relationship between the unnecessary use of mupirocin in the community and the emergence of mupirocin-resistant MRSA among hospitalized inpatients and to evaluate the cost-effectiveness of a search and destroy strategy for MRSA in Thailand.

ACKNOWLEDGMENTS

Financial support. This study was supported by the National Research University Project of Thailand Office of Higher Education Commission (to A.A.).

Potential conflicts of interest. All authors report no conflicts of interest relevant to this article.

**Anucha Apisarnthanarak, MD;¹ David K. Warren, MD;²
Victoria J. Fraser, MD²**

Affiliations: 1. Division of Infectious Diseases, Thammasat University Hospital, Pratumthani, Thailand; 2. Washington University School of Medicine, Saint Louis, Missouri.

Address correspondence to Anucha Apisarnthanarak, MD, Division of Infectious Diseases, Thammasat University Hospital, Pratumthani, Thailand, 12120 (anapisarn@yahoo.com).

Infect Control Hosp Epidemiol 2011;32(5):522-523

© 2011 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2011/3205-0022\$15.00. DOI: 10.1086/659953

REFERENCES

1. Perl TM, Cullen JJ, Wenzel RP, et al. Intranasal mupirocin to prevent postoperative *Staphylococcus aureus* infections. *N Engl J Med* 2002;346:1871-1877.
2. Wilcox MH, Hall J, Pike H, et al. Use of perioperative mupirocin to prevent methicillin-resistant *Staphylococcus aureus* (MRSA) orthopaedic surgical site infections. *J Hosp Infect* 2003;54:196-201.
3. Jog S, Cunningham R, Cooper S, et al. Impact of preoperative screening for methicillin-resistant *Staphylococcus aureus* by real-time polymerase chain reaction in patients undergoing cardiac surgery. *J Hosp Infect* 2008;69:124-130.
4. Jones JC, Rogers TJ, Brookmeyer P, et al. Mupirocin resistance in patients colonized with methicillin-resistant *Staphylococcus aureus* in a surgical intensive care unit. *Clin Infect Dis* 2007;45:541-547.
5. Cookson BD. The emergence of mupirocin resistance: a challenge to infection control and antibiotic prescribing practice. *J Antimicrob Chemother* 1998;41:11-18.
6. Apisarnthanarak A, Tunpornchai J, Tanawitt K, Mundy LM. Non-judicious dispensing of antibiotics by drug stores in Pratumthani, Thailand. *Infect Control Hosp Epidemiol* 2008;29:572-575.

Pseudo-outbreak of *Pseudomonas putida* Respiratory Infection Caused by Laboratory Contamination

To the Editor—*Pseudomonas putida* is a gram-negative, aerobic bacterium that is a common inhabitant of soil, plants, and water. It has been found to cause septicemia in immunocompromised patients,^{1,2} and nosocomial transmission has been reported, associated with contaminated heparin or antiseptic solutions.³⁻⁵ Nevertheless, its isolation from clinical specimens is rare, and it is usually considered an environmental contaminant. *P. putida* has also been reported as a cause of pseudo-outbreaks in contaminated urine collection kits and contaminated commercial antifog solutions.^{6,7} We now report a pseudo-outbreak of *P. putida* respiratory infections, involving 5 patients, caused by an automated spiral plater.

The infection control unit was notified on October 2, 2009, of a cluster of *P. putida* isolations from blind distal bronchial samples from 3 patients undergoing ventilation in the intensive care unit (ICU). The first positive sample had been recorded 5 days before, and the two others the day of notification. In each sample, *P. putida* was isolated among other bacteria at a significant level (at least 10³ cfu/mL). Review of the previous 3 months of microbiology laboratory records found 2 other instances of *P. putida* isolation, one from a bronchial sample of another ICU patient and the other from sputum of a patient in the pulmonology department. All isolates presented an identical antibiotic susceptibility pattern. During the same period, *P. putida* was not isolated from other clinical specimens. A thorough ward-based investigation revealed no epidemiological link to suggest cross-infection between the patients. In particular, the pulmonology department patient had never been hospitalized in the ICU and did not share any device with the ICU patients. Therefore, the investigation focused on the microbiology laboratory, where the 5 samples were processed by the same device (a Whitley Automated Spiral Plater WASP 2; Don Whitley Scientific) dedicated to the clinical respiratory samples.

The WASP 2 was used for many years without any problem. It is a fully automated spiral plater, able to load a sample with a stylus, inoculate a plate, clean the stylus in a sanitizing solution (70% alcohol), and finally rinse the stylus with sterile water loaded from a 110-mL reusable container (Figure 1). The recommendation of the WASP 2 user manual⁸ is to sterilize the containers filled with sterile water by autoclaving.

Laboratory procedures for handling specimens and cleaning processes were reviewed with laboratory personnel. Aseptically collected samples of domestic water, demineralized rinse water, stylus, sanitizing solution, and each of the 12 reusable containers were obtained for bacterial culture. For the stylus, a 100- μ L aliquot of sterile water was loaded by the stylus and directly deposited on a plate. For domestic water, demineralized water, sanitizing solution, and reusable containers containing 110 mL of sterile water, 100 mL of liquid were filtered and inoculated on plates. Cultures of all specimens were obtained using conventional microbiologic methods. Restriction endonuclease DNA profiles were determined by pulsed-field gel electrophoresis for all available isolates, using the restriction enzyme *Spe*I.

There was no change in personnel, microbiological technique, or culture medium. The review of the procedures with the microbiology laboratory personnel revealed a change in the process of reusable-container disinfection a few weeks before the first case occurred. The autoclave usually used to sterilize the containers filled with sterile water had broken down, and an alternate procedure consisting of a chemical disinfection was performed until the autoclave was repaired. However, as the written procedure requested a rinse with sterile water after immersion in a bactericidal solution (DDN 250, Franklab Laboratory) for 60 minutes, the employee responsible for container disinfection had immersed the con-