Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

7-1-2005

Comprehensive treatment of late-onset tibia vara

J. Eric Gordon

St. Louis Shriner's Hospital for Children

Fred P. Heidenreich

Washington University School of Medicine in St. Louis

Chris J. Carpenter

St. Louis Shriner's Hospital for Children

Jill Kelly-Hahn

St. Louis Shriner's Hospital for Children

Perry L. Schoenecker

St. Louis Shriner's Hospital for Children

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Part of the [Medicine and Health Sciences Commons](#)

Please let us know how this document benefits you.

Recommended Citation

Gordon, J. Eric; Heidenreich, Fred P.; Carpenter, Chris J.; Kelly-Hahn, Jill; and Schoenecker, Perry L., "Comprehensive treatment of late-onset tibia vara." *The Journal of Bone and Joint Surgery*. 87, 7. 1561-1570. (2005).

https://digitalcommons.wustl.edu/open_access_pubs/848

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

COMPREHENSIVE TREATMENT OF LATE-ONSET TIBIA VARA

BY J. ERIC GORDON, MD, FRED P. HEIDENREICH, MD, CHRIS J. CARPENTER, RN,
JILL KELLY-HAHN, RN, AND PERRY L. SCHOENECKER, MD

*Investigation performed at the Washington University School of Medicine,
St. Louis Shriners Hospital for Children, and St. Louis Children's Hospital, St. Louis, Missouri*

Background: Late-onset tibia vara (Blount disease) can be difficult to treat because of frequent morbid obesity and associated deformities, including distal femoral varus, proximal tibial procurvatum, and distal tibial valgus, that contribute to lower extremity malalignment. We present a comprehensive approach that addresses all components of the deformity and allows restoration of the anatomic and mechanical axes.

Methods: Fifteen consecutive patients (nineteen lower extremities) with late-onset tibia vara were managed with this comprehensive approach. The mean age of the patients at the time of surgery was 14.9 years, and the mean weight was 113 kg. Standing anteroposterior and lateral radiographs were made preoperatively and at the time of the final follow-up. Preoperatively, the mean mechanical axis deviation was 108 mm, the mean lateral distal femoral angle was 95°, and the mean mechanical medial proximal tibial angle was 71°. In all nineteen extremities, the proximal tibial varus deformity was corrected by means of a valgus osteotomy and application of an Ilizarov ring external fixator. Distal femoral varus was corrected by means of either hemiepiphyseal stapling or valgus osteotomy with blade-plate fixation in thirteen of the nineteen extremities. Distal tibial valgus was treated either with hemiepiphyseal stapling or with varus osteotomy and gradual correction with use of the Ilizarov external fixator in eleven of the nineteen extremities.

Results: After a mean duration of follow-up of 5.0 years, the mean mechanical axis deviation had improved to 1 mm (range, 20 to -30 mm), the lateral distal femoral angle had improved to 87° (range, 83° to 98°), and the mechanical medial proximal tibial angle had improved to 88° (range, 83° to 98°). The mean time required for correction of the proximal tibial varus deformity was thirty-one days, and the external fixator was removed at a mean of 4.5 months postoperatively. All patients had development of one or more superficial pin-track infections (mean, 1.9 pin-site infections per patient). No wound infections, nonunions, or neurovascular complications occurred. Eighteen of the nineteen extremities were pain-free at the time of the final follow-up.

Conclusions: This comprehensive approach allowed restoration of the mechanical and anatomic axes of the lower extremity in patients with late-onset tibia vara, resulting in a resolution of symptoms as a result of normalization of the weight-bearing forces across the knee and ankle. We believe that this approach will decrease the risk of early degenerative arthritis of the knee.

Level of Evidence: Therapeutic Level IV. See Instructions to Authors for a complete description of levels of evidence.

Blount described a group of children who had an onset of varus deformity of the proximal part of the tibia in later childhood or adolescence, a condition that he described as adolescent tibia vara¹. Thompson et al. clarified this classification by categorizing patients who had had a juvenile onset of varus deformity (at the age of four to ten years) along with those who had had a true adolescent onset (at the age of eleven years or more) as having late-onset tibia vara^{2,3}. Early-onset or infantile tibia vara is characterized by a more severe proximal tibial deformity and substantial depression of the medial aspect of the proximal tibial epiphysis^{2,5}. Children with late-onset tibia vara present a challenging combination of marked genu varum deformity, usually accompanied by obesity⁶. The deformity usually develops in middle childhood or

early adolescence^{7,8} and is differentiated from infantile tibia vara on the basis of the less severe epiphyseal deformity and the history of normal lower extremity alignment prior to the development of a genu varum deformity^{1,4,5,9}.

Although the name of the disorder suggests that varus of the proximal part of the tibia is the only deformity present, this is not the case. The proximal tibial varus deformity is produced by a posteromedial growth suppression, which initially produces varus and then progressive procurvatum of the proximal part of the tibia^{3,9-12}. Associated toeing-in gait also worsens as the deformity progresses, initially as a functional maneuver to allow the foot to be placed as close to the line of progression as possible and subsequently as a result of progressive internal tibial torsion⁹. The combination of proximal

tibial varus and procurvatum as well as internal tibial torsion results in a complex three-dimensional deformity. The progressive proximal tibial varus deformity produces increased forces on the medial aspect of the distal femoral physis, leading to growth suppression and a distal femoral varus deformity as well. In severe cases, the varus deformity in the distal part of the femur can be nearly as severe as that in the proximal part of the tibia¹³. As the proximal tibial and distal femoral varus deformities increase, substantial strain is placed on the lateral collateral ligament of the knee, which can lead to laxity and instability of the knee joint⁹. Finally, in severe cases, compensatory distal tibial valgus may develop, allowing the patient to align the ankle joint parallel to the floor⁹.

Although the formation of an osseous bar across the medial aspect of the proximal tibial physis along with medial epiphyseal depression is not uncommon in patients with severe infantile tibia vara, similar bar formation in patients with late-onset tibia vara has not been reported, to our knowledge^{5,14,15}. Depression of the medial aspect of the proximal tibial epiphysis requiring medial plateau elevation has not been reported in patients with late-onset tibia vara⁴.

Although late-onset tibia vara occasionally may occur in thin or normal-sized patients, the majority of patients are obese^{2,6,16-21}. These patients typically are in the ninetieth percentile of weight for both age and height^{17,18,20-22}. During normal walking, nonobese individuals place the foot close to the midline of foot progression, thereby minimizing weight transfer and energy expenditure. Obese individuals with large thighs have a very difficult time adducting the hip adequately to place the foot for midline progression. Davids et al.²² speculated that this "fat thigh syndrome" produces a varus moment on the knee that leads to increased pressure on the medial aspect of the proximal tibial physis and inhibits growth in accordance with the Hueter-Volkman law²³.

Since 1994, we have addressed all of the deformities associated with late-onset tibia vara with a comprehensive approach. Proximal tibial varus is treated with a valgus osteotomy followed by gradual correction with use of an Ilizarov external fixator. If the patient has open physes and adequate growth remains for correction of the deformities of the distal part of the femur and distal part of the tibia, hemiepiphyseal stapling is performed. When the physes are closing, these deformities are corrected with osteotomy. Distal femoral varus is treated with a valgus osteotomy and stabilization with a blade-plate. Distal tibial valgus is treated with a varus osteotomy followed by gradual correction with use of an Ilizarov external fixator. All procedures are performed simultaneously. The purpose of the present study was to evaluate the results of this comprehensive approach to the deformities in patients with late-onset tibia vara.

Materials and Methods

Fifteen consecutive patients (nineteen lower extremities) with late-onset tibia vara were managed with our comprehensive approach between January 1, 1994, and December 31, 1998. Medical records, hospital charts, and radiographs were

retrospectively reviewed with regard to demographic characteristics, operative procedures, complications, and functional and radiographic data. All nineteen extremities underwent a proximal tibial valgus osteotomy followed by gradual correction of the proximal tibial varus deformity with use of an Ilizarov circular external fixator (Smith-Nephew-Richards, Memphis, Tennessee). This device was also used to correct associated proximal tibial procurvatum and internal tibial torsion. The distal femoral varus deformity was corrected with use of a valgus osteotomy and stabilization with a 95° adult condylar blade-plate (Synthes, Paoli, Pennsylvania) if insufficient growth remained to allow lateral hemiepiphyseal stapling. Distal tibial valgus deformity was treated either with medial hemiepiphyseal stapling or with varus osteotomy and gradual correction with use of an extension of the Ilizarov frame. All patients were followed for a minimum of two years postoperatively.

The lower extremities were evaluated preoperatively and at the time of the final follow-up with use of standard radiographs. These included a standing anteroposterior radiograph of both lower extremities; a standing anteroposterior radiograph of both ankles, showing the distal half of the tibia; and a lateral radiograph of the knee, showing the proximal half of the tibia. Line measurements were performed according to the system of Paley et al. (Figs. 1-A and 1-B)²⁴⁻²⁷. These measurements included the mechanical axis deviation, the mechanical lateral distal femoral angle (normal, 88°; range, 86° to 89°), the mechanical medial proximal tibial angle (normal, 87°; range, 85° to 89°), the mechanical lateral distal tibial angle (normal, 91°; range, 88° to 95°), and the joint-line congruency angle (normal, 2°; range, 1° to 3°). Lateral radiographs were evaluated to determine the posterior proximal tibial angle (normal, 80°; range, 77° to 84°).

All patients complained of pain in the involved knee or knees preoperatively and were observed to have a lateral knee thrust when walking.

The mean age at the time of surgery was 14.9 years (range, 10.6 to 18.1 years). The etiology was juvenile onset tibia vara for three patients and adolescent tibia vara for the remaining twelve patients. The mean weight was 113 kg (range, 43 to 178 kg). Thirteen patients (87%) were above the ninety-fifth percentile for weight according to their age, and fourteen patients (93%) were above the ninety-fifth percentile for weight according to their height. Four patients were so obese that they had sleep apnea that necessitated nighttime use of continuous positive airway pressure.

Preoperative anteroposterior radiographic measurements of the lower extremities demonstrated a mechanical axis deviation of 108 mm (range, 41 to 208 mm), a mean mechanical lateral distal femoral angle of 95° (range, 82° to 102°), and a mean mechanical medial proximal tibial angle of 71° (range, 61° to 77°)^{28,29}. These measurements all indicated substantial lower extremity malalignment. The mean joint-line congruency angle was 3° (range, -2° to 12°). Radiographs of the ankles revealed a mean mechanical lateral distal tibial angle of 85° (range, 79° to 93°). Lateral radiographs revealed a mean posterior proximal tibial angle of 71° (range, 58° to 88°). The

Fig. 1-A

Fig. 1-B

Fig. 1-A Coronal plane illustration of the lower extremity, demonstrating the mechanical axis deviation (MAD), the lateral distal femoral angle (LDFA), the joint-line congruency angle (JLCA), the medial proximal tibial angle (MPTA), and the lateral distal tibial angle (LDTA). **Fig. 1-B** Sagittal plane illustration of the proximal part of the tibia and the distal part of the femur, demonstrating the posterior proximal tibial angle (PPTA).

mean preoperative lower limb-length discrepancy in the eleven patients with unilateral disease was 14 mm (range, 0 to 30 mm) as assessed either with scanography or on the standing anteroposterior radiograph of both lower extremities. None of the four patients with bilateral disease had a substantial limb-length discrepancy.

All nineteen extremities had a proximal tibial osteotomy with placement of an Ilizarov circular external fixator. Correction of the distal femoral varus was indicated if the lateral distal femoral angle was $>95^\circ$ (representing $>5^\circ$ of mechanical varus). According to this criterion, thirteen of the nineteen femora had a distal varus deformity that required correction. Eleven of these thirteen extremities had a distal femoral valgus

osteotomy with blade-plate fixation, whereas the other two had enough remaining growth to undergo lateral distal femoral hemiepiphyseal stapling.

Correction of the distal tibial valgus deformity was performed if the mechanical lateral distal tibial angle was $\leq 86^\circ$ (representing $\geq 5^\circ$ of mechanical valgus). According to this criterion, eleven of the nineteen tibiae had a distal valgus deformity that required correction. Eight of these extremities had a distal tibial varus osteotomy with gradual correction, and the other three were corrected by means of medial hemiepiphyseal stapling. Two extremities had lateral instability of the knee secondary to lateral collateral ligament laxity and were treated with concomitant distal transport of the proximal part of the

Fig. 2-A

Fig. 2-B

Fig. 2-C

Figs. 2-A through 2-H Radiographs of the right lower extremity of a 13.4-year-old boy who presented with right adolescent tibia vara with proximal tibial varus, distal tibial valgus, and joint-line incongruity with lateral knee-joint laxity. **Fig. 2-A** Preoperative standing anteroposterior radiograph. The mechanical axis deviation is 98 mm, the lateral distal femoral angle is 92°, the medial proximal tibial angle is 72°, and the joint-line congruency angle is 5°. **Fig. 2-B** Lateral radiograph of the right knee, demonstrating proximal tibial procurvatum with a posterior proximal tibial angle of 73°. **Fig. 2-C** Standing anteroposterior radiograph of the right ankle, showing mild distal tibial valgus with a lateral distal tibial angle of 86°.

fibula to tighten the ligament. Four extremities underwent only a proximal tibial osteotomy, six extremities underwent a proximal tibial osteotomy with correction of either the distal part of the femur or the distal part of the tibia, and seven extremities underwent deformity correction at the proximal part of the tibia, the distal part of the femur, and the distal part of the tibia. Two extremities required deformity correction at the proximal part of the tibia, the distal part of the femur, and the distal part of the tibia as well as fibular transport to tighten the lateral collateral ligament.

Surgical Technique

Preoperatively, radiographs were analyzed and planning was performed according to the techniques described by Paley and

Tetsworth^{24,25}. An Ilizarov circular external fixator was preconstructed to lengthen the tibia as needed and to simultaneously correct varus and procurvatum through an oblique-plane placement of the hinges. After correction of the angulatory deformity and restoration of length, the fixator was modified in situ to correct rotational malalignment. In two patients (three tibiae) with greater rotational deformity, an inclined oblique-plane hinge initially was constructed to provide simultaneous correction of alignment in the frontal, coronal, and rotational planes, with restoration of length. If the patient was skeletally mature, a proximal two-ring block, with the rings located approximately 2 cm apart, was utilized to stabilize the proximal part of the tibia. If the patient was skeletally immature, a single proximal ring was used to avoid performing the osteotomy

Fig. 2-D

Anteroposterior radiograph of the right tibia, made after application of the Ilizarov external fixator, showing the proximal tibial osteotomy site and the placement of pins into the fibula for distal fibular transport and proximal tibial osteotomy.

distally in the diaphysis of the tibia. This proximal ring block was connected to the distal ring segments with appropriately placed hinges or struts. Two distal rings were used; the rings were locked with rods if a distal tibial osteotomy was not required, and they were connected with appropriately placed hinges or struts if a distal osteotomy was to be performed. If clinically important lateral collateral ligament laxity was present, a separate block of pins was used to secure the distal end of the proximal fibular fragment and to transport it distally, thereby tightening the lateral collateral ligament.

After appropriate general endotracheal anesthesia had been induced and the patient had been carefully positioned on a radiolucent surgical table (Orthopedic Systems, Hayward, California), the entire lower extremity was prepared and draped and a sterile tourniquet was placed about the proximal part of the thigh. The distal femoral varus deformity was corrected first, when necessary. A supracondylar distal femoral opening-wedge valgus osteotomy was performed through a lateral approach³⁰⁻³³. A large Kirschner wire was first placed

perpendicular to the lateral cortex of the femur to serve as a reference wire. A second Kirschner wire was then placed into the distal metaphysis at or just proximal to the physis. It was oriented such that the subtended angle between it and the initial Kirschner wire was 5° less than the preoperatively measured deformity (to allow for the placement of the 95° condylar blade-plate). The initial reference Kirschner wire was then removed. The chisel was then placed into the distal femoral metaphysis, just proximal and parallel to the second Kirschner wire and the blade length. An osteotomy was then created 2 cm proximal to the chisel with use of an oscillating saw, and the chisel was withdrawn. The five-hole 95° blade-

Fig. 2-E

Standing anteroposterior radiograph, made seven months postoperatively, demonstrating a neutral mechanical axis.

plate was then inserted into the slot created by the chisel and was impacted into place. The second Kirschner wire was then removed. The distal part of the femur and the attached blade-plate were then reduced to the femoral shaft, creating a stable opening-wedge osteotomy.

After the femoral osteotomy but prior to closure of the femoral wound, while the tourniquet was still inflated, a posterolateral incision was made over the fibula at the junction of its middle and distal thirds. Subperiosteal exposure of the fibula was performed to avoid injury of the branches of the deep peroneal nerve to the extensor hallucis longus or the peroneal artery and vein. A 1-cm section of the fibula was removed with use of an oscillating saw and, if necessary, was cut into pieces and placed into the femoral osteotomy site as bone graft. Both fem-

oral and fibular osteotomy wounds were then thoroughly irrigated and closed. The femoral wound was closed over a drain.

If hemiepiphyseal stapling of the distal part of the femur or the distal part of the tibia was planned, the staples were placed extraperiosteally under fluoroscopic guidance as described by Zuege et al.³⁴ prior to application of the external fixator.

The tourniquet was deflated, the preconstructed circular fixator was placed onto the lower extremity, and suction tubing was used to suspend the leg within the external fixator. This tubing was placed under the leg and over the external fixator and was clamped into place with use of hemostats to secure the leg appropriately within the fixator. Transverse and transfibular wires initially were placed both proximally and

Fig. 2-F

Fig. 2-G

Fig. 2-H

Fig. 2-F Lateral radiograph of the knee, showing partial correction of the proximal tibial procurvatum with a healed osteotomy site. The posterior proximal tibial angle is 76°. **Fig. 2-G** Standing anteroposterior radiograph of the right ankle, made seven months postoperatively, showing partial correction of the distal tibial valgus with a lateral distal tibial angle of 88°. **Fig. 2-H** Standing anteroposterior long-cassette radiograph, made four years postoperatively, showing maintenance of the correction and a neutral mechanical axis. The mechanical axis deviation is 7 mm, the lateral distal femoral angle is 90°, the medial proximal tibial angle is 88°, and the joint-line congruency angle is 2°. At the time of the most recent follow-up, the patient was asymptomatic.

TABLE I Radiographic Measurements

	Preoperative*	Postoperative*
Mechanical axis deviation (<i>mm</i>)	108 (41 to 208)	1 (20 to -30)
Mechanical lateral distal femoral angle (<i>deg</i>)	95 (82 to 102)	87 (83 to 98)
Joint-line congruency angle (<i>deg</i>)	3 (-2 to 12)	2 (0 to 5)
Mechanical medial proximal tibial angle (<i>deg</i>)	71 (61 to 77)	88 (83 to 98)
Mechanical lateral distal tibial angle (<i>deg</i>)	85 (79 to 93)	89 (82 to 93)
Posterior proximal tibial angle (<i>deg</i>)	71 (58 to 88)	77 (57 to 89)

*The values are given as the mean, with the range in parentheses.

distally, with care being taken to place the closest wires approximately 5 mm away from the physis. If a fibular transport to tighten the lateral collateral ligament had been planned, no transfibular wire was placed. The transfibular wire, if placed, was positioned within the metaphysis of the fibula as the surgeon carefully palpated to identify and avoid the peroneal nerve. If a distal tibial osteotomy was necessary, a second distal transverse wire was placed. Anterolateral and anteromedial half-pins were then placed into the proximal part of the tibia. Three or four half-pins were then placed into the tibial diaphysis, in locations ranging from anterior to medial.

The tourniquet was reinflated and the proximal tibial osteotomy was performed after a careful subperiosteal exposure distal to the patellar tendon insertion and approximately 1 cm distal to the proximal tibial fixation pins. A drill was used in a plane parallel to the proximal ring to outline the osteotomy site according to the technique described by De Bastiani et al.³⁵. An Ilizarov osteotome was used to connect the drill-holes and to cut the medial and lateral tibial cortices. The osteotome was then driven into the osteotomy site and was twisted to complete the osteotomy and to fracture the posterior cortex. If required, a distal tibial osteotomy was created in an identical fashion. The wounds were then closed, and the tourniquet was deflated. Generous pin releases were performed at the conclusion of the procedure.

Gradual correction was begun on the second postoperative day at a rate of 0.25 mm every six hours. Serial radiographs were made weekly and as needed to monitor correction and bone formation. The patient was encouraged to bear weight as early as possible, and vigorous physical therapy was instituted to maintain mobility and range of motion of the knee and ankle. Daily showers were instituted as pin care and soft-tissue releases were carried out as needed³⁶. The fixator was left in place and was progressively dynamized until complete consolidation of the osteotomy site had occurred.

Results

The fifteen patients were followed postoperatively for a mean of 5.0 years (range, 2.3 to 9.8 years). All patients were skeletally mature at the time of the last follow-up. The mean time required for correction of the proximal tibial varus

deformity was thirty-one days (range, eleven to sixty-eight days). The external fixator was removed after a mean of 4.5 months (range, 2.9 to 6.4 months).

All patients had clinical improvement. At the time of the last follow-up, fourteen patients (eighteen extremities) were asymptomatic and one patient (one extremity) continued to complain of knee pain, the etiology of which could not be determined. The latter patient had a final mechanical axis deviation of 7 mm lateral to the joint center, with good alignment of the distal part of the femur and the proximal part of the tibia. Magnetic resonance imaging demonstrated no evidence of intra-articular abnormality. The pain did not limit the patient's activities, and the patient declined additional intervention.

The lateral knee thrust was eliminated in all patients. All patients had regained full ranges of knee and ankle motion at the time of the last follow-up. No patient had clinically important knee laxity to varus or valgus stress at the time of the last follow-up.

Radiographs that were made at the time of the last follow-up revealed improvement in lower extremity alignment. The mechanical axis deviation improved to a mean of 1 mm (range, 20 to -30 mm) (Table I). The lower extremities were essentially equal in length, with a mean discrepancy of 1 mm (range, -22 mm to 6 mm) at the time of the final follow-up (Figs. 2-A through 2-H).

The mean mechanical lateral distal femoral angle improved to 87° (range, 83° to 98°). In the two patients who underwent hemiepiphyseal stapling, the mechanical lateral distal femoral angle improved from 92° to 87° in one patient and from 96° to 87° in the other. The mechanical lateral distal femoral angle in the eleven patients who underwent distal femoral osteotomy improved from a mean of 97° (range, 94° to 102°) to a mean of 88° (range, 84° to 96°).

The mean mechanical medial proximal tibial angle improved to 88° (range, 83° to 98°). The mean joint-line congruency angle improved to 2° (range, 0° to 5°). The mean posterior proximal tibial angle improved mildly from 71° (range, 58° to 88°) to 77° (range, 57° to 89°). The mean mechanical lateral distal tibial angle improved slightly from 85° (range, 79° to 93°) to 89° (range, 82° to 93°). The mean mechanical lateral distal tibial

angle in the three patients who underwent hemiepiphyseal stapling of the distal part of the tibia improved from 84° (range, 83°, 86°, and 83°) to 89° (86°, 88°, and 92°). The mean mechanical lateral distal tibial angle in the eight patients who underwent osteotomy of the distal part of the tibia improved from 83° (range, 79° to 86°) to 90° (range, 86° to 92°).

All patients had development of one or more superficial pin-site infections (mean, 1.9 infections per patient), which resolved with oral antibiotics³⁶. No patient required intravenous antibiotic therapy. No neurovascular complications were encountered. No patient had development of a deep infection, osteomyelitis, or nonunion.

Postoperatively, the need for additional procedures required fourteen of the fifteen patients to return to the operating room between one and three times per involved extremity. Removal of the external fixator under a general anesthetic was requested by the patient for eighteen of the nineteen extremities. Other indications for additional procedures included wire removal (thirteen patients), fixator modification (two patients), and staple removal (two patients). In all, only one extremity required no additional procedures, six required one additional procedure, nine required two additional procedures, and three required three additional procedures.

Discussion

Patients with late-onset tibia vara typically are obese and can present with complaints of genu varum, knee pain, toeing-in, or knee instability^{5,16-19,22}. The natural history of this condition is unclear, but residual infantile tibia vara leads to chronic knee pain and degenerative arthritis³⁷⁻⁴⁰. Varying amounts of respiratory distress may be present, depending on the patient's obesity and activity level⁴¹⁻⁴³. Occasionally, even the walk from the waiting room to the examination area can be exhausting for these patients. Patients with knee pain often describe medial pain and symptoms of increased pressure in the medial compartment of the knee as well as symptoms of anterior knee pain, probably secondary to maintenance of the knee in a flexed position during gait.

A standing long-cassette anteroposterior radiograph of both lower extremities is essential for the evaluation of these patients. If the patient appears to have unilateral bowing, careful attention should be paid to the contralateral limb because the patient's obesity frequently can mask mild contralateral genu varum. Because of the obesity of these patients, we have found it necessary to make preoperative preparations for special equipment, including a large tourniquet and a radiolucent table that can support the patient's weight.

Although some authors have recommended physeal distraction as a means of achieving correction, this is not possible in skeletally mature adolescents^{44,45}. Laurencin et al.⁴⁶ effectively utilized an oblique, laterally based closing-wedge osteotomy that hinges at the intact cortex just distal to the proximal tibial physis for acute correction of mild deformities. Most patients with late-onset tibia vara require at least proximal tibial osteotomy to correct the deformity. Osteotomy with limited internal fixation, the mainstay of treatment

of infantile tibia vara^{5,15} has little application in the treatment of late-onset tibia vara¹⁸.

The obesity of the patient makes the application of a well-fitting above-the-knee cast difficult, if not impossible, and prevents the adequate control of the position of the osteotomy site. In addition, the older age and size of the patient make patient mobility highly desirable. The problems associated with non-weight-bearing on the affected extremity make walking extremely difficult for these patients. Furthermore, the difficulties associated with assessing limb alignment either intraoperatively or postoperatively make residual proximal tibial varus or undercorrection a possibility. External fixation of the tibia is indicated for the correction of more severe deformities^{21,47-49}. This allows either acute or gradual correction with a monolateral or circular fixator. Price et al. demonstrated that monolateral fixation with either acute or gradual correction can effectively correct the tibial varus deformity in the frontal plane⁴⁸. Alignment in the sagittal plane and rotational correction are more difficult to obtain with use of the monolateral fixator^{21,50}.

A detailed preoperative plan is essential for any surgical procedure. The magnitude and location of the various osseous deformities, laxity of the lateral collateral ligament, the presence of joint contractures, and lower limb-length discrepancy should all be assessed and incorporated into the preoperative plan. The overall plan should correct each of the constituent deformities, normalizing the mechanical and anatomic axes in order to avoid increased stresses resulting from chronic malalignment as well as increased shear forces on the knee or ankle resulting from joint-line obliquity. Although the long-term results of our comprehensive technique are not known, this approach has provided pain relief, a stable correction, and restoration of normal lower extremity alignment in this difficult patient population.

During the planning and execution of the varus correction, particular attention should be paid to normalizing distal femoral and proximal tibial alignment. Because of the obesity of these patients, joint-contact forces are excessive even when normal mechanical alignment is present. Although the mechanical alignment in these patients could be corrected through a single tibial osteotomy that compensates for existing femoral varus, this would leave the knee joint oblique to the mechanical axis and would introduce shear forces during weight-bearing. Because of these concerns, we believe that it is essential to correct all of the deformities separately in order to provide optimal mechanical alignment and to delay the onset of degenerative arthritis.

Circular external fixation with gradual correction of the proximal part of the tibia allows for maximal adjustability of the alignment in all planes and is indicated for more severe deformities and the most obese patients^{17,51}. Other advantages include stable fixation with improved patient mobility; the ability to evaluate alignment in a functional, standing position; and the ability to accurately correct all deformities of the tibia, including proximal tibial varus and procurvatum, internal tibial torsion, and distal tibial valgus.

After correction of the genu varum deformity, the distal

tibial valgus, if uncorrected, can produce a pes planus deformity with hindfoot valgus⁹. This persistent distal tibial valgus leaves the ankle in a position in which it is oblique to forces involved in walking, leading to sheer forces at the joint surface. This can become painful, particularly in obese patients with late-onset tibia vara. Wagner et al.⁵² found decreased tibiotalar contact areas at the ankle in patients with distal tibial deformities of $\geq 10^\circ$. Ting et al.⁵³ found that this effect was exacerbated in patients with decreased subtalar motion who were unable to compensate for the valgus. This has led us to correct deformities of $>5^\circ$ at the distal part of the tibia.

In conclusion, this comprehensive approach to late-onset tibia vara is effective, but careful preoperative planning and an analysis of all of the deformities are essential. In particular, correction of each of the deformities in the lower extremity allows the mechanical alignment of the lower extremity to be optimized with the restoration of anatomic and mechanical axes. This approach has led to relief of pain through adolescence, and we are hopeful that it will eliminate the onset of degenerative arthritis in the knee and ankle, although the long-term results remain to be seen. ■

J. Eric Gordon, MD
Chris J. Carpenter, RN
Jill Kelly-Hahn, RN
Perry L. Schoenecker, MD
St. Louis Shriners Hospital for Children, 2001 South Lindbergh Boulevard, St. Louis, MO 63131. E-mail address for J.E. Gordon: gordone@msnotes.wustl.edu

Fred P. Heidenreich, MD
Department of Orthopaedic Surgery, Washington University School of Medicine, One Barnes-Jewish Hospital Plaza, Suite 11300 West Pavilion, St. Louis, MO 63110

The authors did not receive grants or outside funding in support of their research or preparation of this manuscript. They did not receive payments or other benefits or a commitment or agreement to provide such benefits from a commercial entity. No commercial entity paid or directed, or agreed to pay or direct, any benefits to any research fund, foundation, educational institution, or other charitable or nonprofit organization with which the authors are affiliated or associated.

doi:10.2106/JBJS.02276

References

- Blount WP. Tibia vara. Osteochondrosis deformans tibiae. *J Bone Joint Surg.* 1937;19:1-29.
- Thompson GH, Carter JR, Smith CW. Late-onset tibia vara: a comparative analysis. *J Pediatr Orthop.* 1984;4:185-94.
- Thompson GH, Carter JR. Late-onset tibia vara (Blount's disease). Current concepts. *Clin Orthop Relat Res.* 1990;255:24-35.
- Schoenecker PL, Johnston R, Rich MM, Capelli AM. Elevation of the medial plateau of the tibia in the treatment of Blount disease. *J Bone Joint Surg Am.* 1992;74:351-8.
- Schoenecker PL, Meade WC, Pierron RL, Sheridan JJ, Capelli AM. Blount's disease: a retrospective review and recommendations for treatment. *J Pediatr Orthop.* 1985;5:181-6.
- Cook SD, Lavernia CJ, Burke SW, Skinner HB, Haddad RJ Jr. A biomechanical analysis of the etiology of tibia vara. *J Pediatr Orthop.* 1983;3:449-54.
- Feldman MD, Schoenecker PL. Use of the metaphyseal-diaphyseal angle in the evaluation of bowed legs. *J Bone Joint Surg Am.* 1993;75:1602-9.
- Kling TF Jr. Angular deformities of the lower limbs in children. *Orthop Clin North Am.* 1987;18:513-27.
- Schoenecker PL, Rich MM. The lower extremity. In: Morrissy RT, Weinstein SL, editors. *Lovell and Winter's pediatric orthopaedics*. 5th ed. Philadelphia: Lippincott, Williams and Wilkins; 2001. p 1059-104.
- Carter JR, Leeson MC, Thompson GH, Kalamchi A, Kelly CM, Makley JT. Late-onset tibia vara: a histopathologic analysis. A comparative evaluation with infantile tibia vara and slipped capital femoral epiphysis. *J Pediatr Orthop.* 1988;8:187-95.
- Beskin JL, Burke SW, Johnston CE 2nd, Roberts JM. Clinical basis for a mechanical etiology in adolescent Blount's disease. *Orthopedics.* 1986;9:365-70.
- Wenger DR, Mickelson M, Maynard JA. The evolution and histopathology of adolescent tibia vara. *J Pediatr Orthop.* 1984;4:78-88.
- Kline SC, Bostrum M, Griffin PR. Femoral varus: an important component in late-onset Blount's disease. *J Pediatr Orthop.* 1992;12:197-206.
- Loder RT, Schaffer JJ, Bardenstein MB. Late-onset tibia vara. *J Pediatr Orthop.* 1991;11:162-7.
- Rab GT. Oblique tibial osteotomy for Blount's disease (tibia vara). *J Pediatr Orthop.* 1988;8:715-20.
- Langenskiold A. Tibia vara. A critical review. *Clin Orthop Relat Res.* 1989;246:195-207.
- Stanitski DF, Dahl M, Louie K, Grayhack J. Management of late-onset tibia vara in the obese patient by using circular external fixation. *J Pediatr Orthop.* 1997;17:691-4.
- Buckley SL, Carlson JR, Robertson WW Jr. Osteotomy of the proximal tibia in the treatment of adolescent tibia vara. *Orthop Rev.* 1993;22:1319-27.
- Henderson RC, Kemp GJ Jr, Greene WB. Adolescent tibia vara: alternatives for operative treatment. *J Bone Joint Surg Am.* 1992;74:342-50.
- Dietz WH Jr, Gross WL, Kirkpatrick JA Jr. Blount disease (tibia vara): another skeletal disorder associated with childhood obesity. *J Pediatr.* 1982;101:735-7.
- Stanitski DF, Srivastava P, Stanitski CL. Correction of proximal tibial deformities in adolescents with the T-Garches external fixator. *J Pediatr Orthop.* 1998;18:512-7.
- Davids JR, Huskamp M, Bagley AM. A dynamic biomechanical analysis of the etiology of adolescent tibia vara. *J Pediatr Orthop.* 1996;16:461-8.
- Hueter C. Anatomische Studien an den Extremitätengelenken Neugeborener und Erwachsener. *Arch Pathol Anat.* 1862;25:572-99.
- Paley D, Tetsworth K. Mechanical axis deviation of the lower limbs. Preoperative planning of uniapical angular deformities of the tibia or femur. *Clin Orthop Relat Res.* 1992;280:48-64.
- Paley D, Tetsworth K. Mechanical axis deviation of the lower limbs. Preoperative planning of multiapical frontal plane angular and bowing deformities of the femur and tibia. *Clin Orthop Relat Res.* 1992;280:65-71.
- Henderson RC, Kemp GJ. Assessment of the mechanical axis in adolescent tibia vara. *Orthopedics.* 1991;14:313-6.
- Paley D, Chaudray M, Pirone AM, Lentz P, Kautz D. Treatment of malunions and mal-nonunions of the femur and tibia by detailed preoperative planning and the Ilizarov techniques. *Orthop Clin North Am.* 1990;21:667-91.
- Moreland JR, Bassett LW, Hanker GJ. Radiographic analysis of the axial alignment of the lower extremity. *J Bone Joint Surg Am.* 1987;69:745-9.
- Bar HF, Breitfuss H. Analysis of angular deformities on radiographs. *J Bone Joint Surg Br.* 1989;71:710-1.
- Healy WL, Anglen JO, Wasilewski SA, Krackow KA. Distal femoral varus osteotomy. *J Bone Joint Surg Am.* 1988;70:102-9.
- Muller ME, Allgower M, Schneider R, Willenegger H, editors. *Manual of internal fixation: techniques recommended by the AO-ASIF group*. 3rd ed. New York: Springer; 1991.
- Phillips MJ, Krackow KA. High tibial osteotomy and distal femoral osteotomy for valgus or varus deformity around the knee. *Instr Course Lect.* 1998;47:429-36.
- Morrey BF, Edgerton BC. Distal femoral osteotomy for lateral gonarthrosis.

Instr Course Lect. 1992;41:77-85.

34. Zuege RC, Kempken TG, Blount WP Epiphyseal stapling for angular deformity at the knee. *J Bone Joint Surg Am.* 1979;61:320-9.

35. De Bastiani G, Aldegheri R, Renzi-Brivio L, Trivella G. Limb lengthening by callus distraction (callotaxis). *J Pediatr Orthop.* 1987;7:129-34.

36. Gordon JE, Kelly-Hahn J, Carpenter CJ, Schoenecker PL. Pin site care during external fixation in children: results of a nihilistic approach. *J Pediatr Orthop.* 2000;20:163-5.

37. Hofmann A, Jones RE, Herring JA. Blount's disease after skeletal maturity. *J Bone Joint Surg Am.* 1982;64:1004-9.

38. Ingvarsson T, Hagglund G, Ramgren B, Jonsson K, Zayer M. Long-term results after adolescent Blount's disease. *J Pediatr Orthop B.* 1997;6:153-6.

39. Ippolito E, Cauti D. Tibia vara or Blount's disease: pathogenesis, clinical features and treatment. *Ital J Orthop Traumatol.* 1983;9:521-8.

40. Zayer M. Osteoarthritis following Blount's disease. *Int Orthop.* 1980;4:63-6.

41. Barness L. Nutrition and nutritional disorders: obesity. In: Behrman R, Kliegman R, editors. *Nelson textbook of pediatrics.* Philadelphia: Saunders; 1992. p 133-4.

42. Barlow SE, Dietz WH. Obesity evaluation and treatment: Expert Committee recommendations. The Maternal and Child Health Bureau, Health Resources and Services Administration and the Department of Health and Human Services. *Pediatrics.* 1998;102:E29.

43. Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents:

the Bogalusa Heart Study. *Pediatrics.* 1999;103:1175-82.

44. Ganel A, de Pablos J, Alfaro J, Barrios C. Treatment of adolescent Blount disease by asymmetric physeal distraction. *J Pediatr Orthop.* 1997;17:54-8.

45. Ganel A, Heim M, Farine I. Asymmetric epiphyseal distraction in treatment of Blount's disease. *Orthop Rev.* 1986;15:237-40.

46. Laurencin CT, Ferriter PJ, Millis MB. Oblique proximal tibial osteotomy for the correction of tibia vara in the young. *Clin Orthop Relat Res.* 1996;327:218-24.

47. Laville JM, Chau E, Willemen L, Kohler R, Garin C. Blount's disease: classification and treatment. *J Pediatr Orthop B.* 1999;8:19-25.

48. Price CT, Scott DS, Greenberg DA. Dynamic axial external fixation in the surgical treatment of tibia vara. *J Pediatr Orthop.* 1995;15:236-43.

49. Rajacich N, Bell DF, Armstrong PF. Pediatric applications of the Ilizarov method. *Clin Orthop Relat Res.* 1992;280:72-80.

50. Gaudinez R, Adar U. Use of Orthofix T-Garche fixator in late-onset tibia vara. *J Pediatr Orthop.* 1996;16:455-60.

51. Coogan PG, Fox JA, Fitch RD. Treatment of adolescent Blount disease with the circular external fixation device and distraction osteogenesis. *J Pediatr Orthop.* 1996;16:450-4.

52. Wagner KS, Tarr RR, Resnick C, Sarmiento A. The effect of simulated tibial deformities on the ankle joint during the gait cycle. *Foot Ankle.* 1984;5:131-41.

53. Ting AJ, Tarr RR, Sarmiento A, Wagner K, Resnick C. The role of subtalar motion and ankle contact pressure changes from angular deformities of the tibia. *Foot Ankle.* 1987;7:290-9.