

9-1-2022

The transition to parenthood in obstetrics: Enhancing prenatal care for 2-generation impact

Catherine Monk
Columbia University

Sona Dimidjian
University of Colorado, Boulder

Ellen Galinsky
Bezos Family Foundation

Kimberly D. Gregory
Cedars-Sinai

M. Camille Hoffman
University of Colorado, Denver

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/oa_4



Part of the [Medicine and Health Sciences Commons](#)

Please let us know how this document benefits you.

Recommended Citation

Monk, Catherine; Dimidjian, Sona; Galinsky, Ellen; Gregory, Kimberly D.; Hoffman, M. Camille; Howell, Elizabeth A.; Miller, Emily S.; Osborne, Cynthia; Rogers, Cynthia E.; Saxbe, Darby E.; and D'Alton, Mary E., "The transition to parenthood in obstetrics: Enhancing prenatal care for 2-generation impact." *American Journal of Obstetrics & Gynecology MFM*. 4, 5. 100678 (2022).
https://digitalcommons.wustl.edu/oa_4/1582

This Open Access Publication is brought to you for free and open access by the Open Access Publications at Digital Commons@Becker. It has been accepted for inclusion in 2020-Current year OA Pubs by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Catherine Monk, Sona Dimidjian, Ellen Galinsky, Kimberly D. Gregory, M. Camille Hoffman, Elizabeth A. Howell, Emily S. Miller, Cynthia Osborne, Cynthia E. Rogers, Darby E. Saxbe, and Mary E. D'Alton

The transition to parenthood in obstetrics: enhancing prenatal care for 2-generation impact



Catherine Monk, PhD; Sona Dimidjian, PhD; Ellen Galinsky, MS; Kimberly D. Gregory, MD, MPH; M. Camille Hoffman, MD, MSc; Elizabeth A. Howell, MD, MPP; Emily S. Miller, MD, MPH; Cynthia Osborne, PhD, MPP; Cynthia E. Rogers, MD; Darby E. Saxbe, PhD; Mary E. D'Alton, MD

Obstetrics, the specialty overseeing infant and parent health before birth, could be expanded to address the interrelated areas of parents' prenatal impact on children's brain development and their own psychosocial needs during a time of immense change and neuroplasticity. Obstetrics is primed for the shift that is happening in pediatrics, which is moving from its traditional focus on physical health to a coordinated, whole-child, 2- or multigeneration approach. Pediatric care now includes developmental screening, parenting education, parent coaching, access to developmental specialists, brain-building caregiving skills, linkages to community resources, and tiered interventions with psychologists. Drawing on decades of developmental origins of health and disease research highlighting the prenatal beginnings of future health and new studies on the transition to parenthood describing adult development from pregnancy to early postpartum, we have proposed that, similar to pediatrics, the integration of education and intervention strategies into the prenatal care ecosystem should be tested for its potential to improve child cognitive and social-emotional development and parental mental health. Pediatric care programs can serve as models of change for the systematic development, testing and, incorporation of new content into prenatal care as universal, first-tier treatment and evidenced-based, triaged interventions according to the level of need. To promote optimal beginnings for the whole family, we have proposed an augmented prenatal care ecosystem that aligns with, and could build on, current major efforts to enhance perinatal care individualization through consideration of medical, social, and structural determinants of health.

Keywords: brain development, care individualization, cognitive development, expectant parent, fetal development, mental health, neurodevelopment, neuroplasticity, prenatal programming, prenatal stress, reproductive justice, social support, social-emotional development, whole person

Cite this article as: Monk C, Dimidjian S, Galinsky E, et al. The transition to parenthood in obstetrics: enhancing prenatal care for 2-generation impact. *Am J Obstet Gynecol MFM* 2022;4:100678.

From the Departments of Obstetrics (Drs Monk and D'Alton) and Gynecology (Dr Monk), Columbia University Irving Medical Center, New York, NY; Psychiatry, Columbia University Irving Medical Center, New York, NY; Department of Psychology and Neuroscience, University of Colorado Boulder, Boulder, CO (Dr Dimidjian); Bezos Family Foundation, Seattle, WA (Ms Galinsky); Department of Obstetrics and Gynecology, Cedars-Sinai, Los Angeles, CA (Dr Gregory); Departments of Obstetrics and Gynecology (Dr Hoffman) and Psychiatry (Dr Hoffman), University of Colorado School of Medicine, Aurora, CO; Psychiatry, University of Colorado School of Medicine, Aurora, CO; Department of Obstetrics and Gynecology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; Departments of Obstetrics and Gynecology (Dr Miller) and Psychiatry (Dr Miller), Northwestern University, Chicago, IL; Psychiatry, Northwestern University, Chicago, IL; Department of Leadership, Policy, and

Organizations, Peabody College, Vanderbilt University, Nashville, TN (Dr Osborne); Department of Psychiatry, Washington University School of Medicine, St. Louis, MO (Dr Rogers); Department of Psychology, University of Southern California, Los Angeles, CA (Dr Saxbe).

Received June 13, 2022; accepted June 15, 2022.

C.M. sits on the board of ZERO TO THREE and has received honoraria from that organization, which has received funding from Blue Meridian Partners and other sponsors to support HealthySteps. E.G. is the chief science officer at the Bezos Family Foundation, which provided a grant to support C.M.'s time on this project. S.D. receives royalties from Guilford Press for work related to mindfulness-based cognitive therapy books and is also a cofounder of and receives revenue from MindfulNoggin, Inc, which supports the online dissemination of mindfulness programs and therapist training tools. S.D., C.E.R., and D.E.S. received honoraria from the Bezos Family Foundation. C.O. directs the Prenatal-to-3 Policy Impact Center, which receives funding from the Pritzker's Children Initiative, the Buffet Early

Childhood Fund, and the Valhalla Foundation, which also provides funding for HealthySteps. M.E.D.A. has had a leadership role in the American College of Obstetricians and Gynecologists II's Safe Motherhood Initiative, which has received unrestricted funding from Merck for Mothers. The remaining authors report no disclosure. The authors report no conflict of interest.

This research was funded, in part, by the Bezos Family Foundation grant and grant number R01HD092062 (C.M.), grant number R01MH117251 (S.D.), The Helping Hand of Los Angeles (K.D.G.), grant number R01MD016029 (E.A.H.), grant number R01HD105499 (E.S.M.), and grant numbers R01MH113570, R01DA046224, and R01MH121877 (C.E.R.). The funding sources had no direct involvement in the writing of this report.

Corresponding author: Catherine Monk, PhD. cem31@columbia.edu

2589-9333/\$36.00

© 2022 Elsevier Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.ajogmf.2022.100678>

Introduction

The US prevalence of neurobehavioral problems is staggering¹—6.1 million children have been diagnosed with attention deficit hyperactivity disorder (ADHD), 4.5 million children have been diagnosed with behavior problems, 4.4 million children have been diagnosed with anxiety, and nearly 7 million children are classified as disabled based on learning difficulties, emotional disturbance, and speech or language impairment.^{2,3} To work toward preventing these conditions, 21st-century pediatrics has pivoted to go beyond its traditional focus on physical health; as such, pediatrics increasingly includes science-informed innovations that optimize 0- to 3-year-olds' sensitivity to early psychosocial influences to shape positive development and mitigate the impact of toxic stress. Obstetrics, the specialty overseeing child development before birth and expectant parent health, similarly could be augmented to address the interrelated areas of parents' prenatal impact on future children's development and their own psychosocial needs. Drawing on decades of developmental origins of health and disease (DOHaD) research highlighting the prenatal beginnings of future health through gene-environment interactions and new studies on the transition to parenthood (TtP) describing adult development from pregnancy to early postpartum, we propose that the integration of education and intervention strategies into the prenatal care ecosystem should be tested through rigorous trials and demonstration projects as we predict that results will show returns to society through improved child cognitive and social-emotional development and parental mental health, similar to those accruing from the shift in pediatrics to a coordinated, whole-child, 2-generation philosophy.^{1,4,5} Our vision has aligned with, and could build on, current major efforts to enhance perinatal care individualization through consideration of medical, social, and structural determinants of health.^{6,7}

Pediatrics enhanced, why not obstetrics?

At the molecular, cellular, organ, system, and behavioral levels, neurodevelopment is shaped by ongoing adaptations to the psychosocial environment, with heightened plasticity in children ages 0 to 3 years underscoring the public health benefits of improving early childhood experiences. Accordingly, pediatrics pivoted, declaring relational health (the promotion of safe, stable, and nurturing relationships) as necessarily in their purview⁸ as well as the child's social-emotional context—including social determinants of health⁹ and how these influences translate into cognitive and emotional outcomes. Pediatric care has expanded to include developmental screening, parenting education, parent coaching, access to developmental specialists, brain-building caregiving skills, and linkages to community resources, leveraging new interdisciplinary teams that may include tiered interventions with psychologists. Key targets include supporting children's language and social development, managing challenging child behaviors, and promoting parent mental health—since absent their own mental well-being, parents are much less likely to have a thriving baby and future child.¹ The outcomes include reductions in developmental delays and behavioral problems and improved cognitive development.¹⁰

During fetal life, one of the most rapid periods of brain development takes place. From gestational weeks 7 to 18, corticogenesis occurs—the cerebral cortex of the brain is formed. Throughout gestation, the interplay of genetic programming along with genetic variation interacts with characteristics of the fetal environment to organize neural elements into complex structural and functional networks known as the connectome. This crucial period of brain development, similar to ages 0 to 3 years, is characterized by tremendous plasticity. The fetal brain is shaped, in part, by adaptation to biological cues related to the pregnant parent's well-being, lifestyle, nutritional status, social factors, and mental health (each of which a co-parent can affect).^{11–15} As DOHaD

data demonstrate, parents' psycho-social experiences are embedded in the future child's neurobiology. Parents affect children's neurodevelopment before birth.

During pregnancy and the postpartum period, parents also experience dramatic psychological, behavioral, and neurobiological adjustments as the foundations of parenting are established. How they navigate this period impacts their future parenting, well-being, and, consequently, child outcomes. Typically, practitioners have bracketed these concerns into a narrow focus on perinatal mental health (PMH) and, specifically, psychiatric diagnoses. However, there is a need to move beyond deficit models to address more holistically the totality of the TtP experience, including the social, cultural, and financial contexts in which this transition occurs.¹⁶

Using a lens of 2-generation development and neuroplasticity, enhanced obstetrical care could help interrupt the intergenerational transmission of disparities and trauma and promote optimal beginnings for the whole family.^{17,18} Pediatric care programs can serve as models of change for the systematic development, testing, and incorporation of new content into prenatal care as universal, first-tier treatment and evidenced-based, triaged interventions according to the level of need. As in pediatrics, hybrid approaches to sharing universal, new content with expectant parents could be created and tested that include residents attending trainings to deliver it; social workers, nurses, and paraprofessionals in obstetrics offices taking on the educating role to provide it; links to community groups who provide it; and novel approaches using the Internet and digital health platforms to teach it (with the necessary attention to access issues for marginalized and less literate populations). As we review below, examples of key content areas to develop include brain development before birth and expectations of parenthood. Provider training and system improvements could include actionable changes to ensure equity and address the mental health of expectant parents—since absent their own mental well-being before birth, parents are less likely to have a thriving

baby and future child.^{1,19} Potential outcomes include reductions in developmental delays and risk of psychiatric problems and improved cognitive development and, for parents, improved mental health and better adjustment to their new role, which, in turn, promotes positive child development.

The womb is an influential first home¹ Brain on board before birth

Third-trimester fetuses register, recognize, and differentially respond to varied stimuli measured by heart rate changes during habituation paradigms.^{20–24} Of note, 3-day-old infants have shown that they can identify their mother's voice vs another woman's voice,² sucking more vigorously on a nonnutritive nipple to elicit hearing it.²⁵ Newborn recognition of maternal voices vs a stranger's voice has been shown in electrophysiological signals as well.²⁶ Newborns demonstrate memory from the in utero period for a spoken passage or melody even if produced by an unfamiliar speaker. The preference for the maternal voice vs another woman's voice likely promotes bonding, while transnatal memory (prebirth learning that carries across the birth divide) for language-specific properties may undergird language development.^{27,28} New research has suggested that maternal speech shapes the frontotemporal networks associated with language and voice processing.²⁹ As in the postnatal period, the fetus' social context shapes its brain development. Parents' relationship with their child develops before birth: this relationship and their emerging caregiving role could be substantiated and supported. Prenatal care could be designed to engage future parents in learning about in utero brain-behavior development and the demonstrable impact parents can have on their child in the prenatal period, for example, in relation to maternal mental health.

The impact of mental health conditions

For almost all parents, pregnancy can bring increased anxiety and uncertainty. In studies of pregnant women, an estimated 30% report psychosocial stress in their daily lives, including job strain and depressive or anxiety symptoms.³⁰ Approximately 10% to 12% of pregnant women meet the criteria for a PMH condition, most often anxiety or depression. Maternal psychiatric morbidity is the most frequent child-birth complication.^{31,32} In the context of structural racism and discrimination, the rates are nearly double for women of color and those experiencing poverty.^{33,34} Furthermore, it is estimated that more than 10% of new fathers experience anxiety and/or depression during the perinatal period.³⁵

Although many fetuses seem resilient to the potential moderating effects of maternal stress and PMH conditions, children of pregnant women in the top 15% of self-reported prenatal anxiety and depression have an estimated 2-fold increased risk of a mental disorder (eg, ADHD or anxiety). This effect extends from childhood to adolescence.³⁶ Furthermore, greater maternal prenatal psychosocial stress is associated with small but statistically significant reductions in intelligence quotient (IQ) scores and less advanced language development.³⁷ A recent task-based functional magnetic resonance imaging (MRI) study showed that newborns of more anxious mothers have greater responses to aberrant stimuli in brain regions related to anxiety disorders in adults (right anterior insula, ventrolateral prefrontal cortex, and cingulate cortex area).³⁸ Importantly, experimental animal studies that control for shared genes have demonstrated that maternal prenatal stress predicts anxiety like behavior in the offspring related to impaired hypothalamic-pituitary-adrenal (HPA) axis regulation and decreased expression of a brain-derived neurotrophic factor that mediates neural plasticity.^{39–41} Excessive maternal stress is transduced to the fetus through maternal HPA axis hormones, immune activation,

and alterations in the maternal-infant microbiome and placental gene expression.^{13,31,32,42,43} Finally, untreated mental health problems during pregnancy often continue into the postpartum period, with untoward effects on child development, such as compromised parent-child bonding, decrements in IQ, and risk of depression in adolescence.^{19,44} In contrast, health-promoting experiences during pregnancy, such as exercise and social support, are positively associated with levels of maternal, fetal, and infant cardiac control (an early marker of emotion regulation)^{45–47} and more adaptive infant stress responses, respectively.⁴⁸ Pre-clinical studies with fertilized chick eggs have shown that auditory enrichment (music and noise exposure) reduces a chick's stimulus-based fear response and associated cortisol reactivity.⁴⁹

Alerting all prospective parents to the fact that their well-being and behaviors may impact their future baby could contribute to the prioritization of health that would benefit both generations as the TtP often is a time of great motivation for change.^{50,51} Sharing the associations between mental health and prenatal brain development in nonthreatening and supportive ways could facilitate the identification with the parenting role: parents matter before birth. Importantly, there is strong evidence that PMH conditions can be prevented^{52–54} and effectively treated through psychotherapy^{54–57} and/or pharmacotherapy.^{58,59} However, the process of connecting perinatal women and their partners with mental healthcare is often fragmented and inefficient, serving neither patients nor providers well. Obstacles to treatment exist at the systems, provider, and patient levels.^{60,61} It is estimated that <50% of women who need it get care⁶⁰ and the rate is even lower among women with fewer resources and of color.⁶² To support healthy parents and children, access to mental healthcare during the TtP should be dramatically and systematically improved via integrated care approaches, such as Compass at Northwestern, Women's Behavioral Health Service at the University of Rochester, and Women's

¹ David Barker is credited with originating this phrase and using it in DOHaD lectures.

² We used gender terms consistent with the research reports we have cited.

Mental Health @Ob/Gyn at Columbia^{60,63–67} and federally funded psychiatry access programs enabling frontline providers to screen, assess, refer, and treat pregnant and postpartum women for depression and other behavioral health conditions—including in rural and medically underserved areas that originated as Massachusetts Child Psychiatry Access Program for Moms (<https://www.mcpapformoms.org/>; <https://www.hrsa.gov/about/news/press-releases/hrsa-awards-12-million-maternal-child-mental-health-programs>).

Unequal beginnings

Annually, 10% of US births are preterm birth (PTB; <37 weeks of gestation), with nearly 400,000 births affected⁶⁸ and recent data showing rates rising.⁶⁹ PTB is associated with a future risk of poor mental health outcomes.⁷⁰ It is estimated that 40% of children born prematurely will experience compromised neurodevelopment, with effects on social, emotional, and physical health.⁷¹ Compared with babies born at or near 37 weeks of gestation, very premature (<32 weeks of gestation) and extremely premature (<28 weeks of gestation) children have 2.25 and 4.04 times the odds, respectively, of receiving an ADHD diagnosis.⁷² However, crucially, those who do have PTB should not be subject to negative bias about their child's future especially as not all children are negatively affected.

PTB is not distributed equally across the population.⁷³ The racial and ethnic inequities are dramatic and independent of socioeconomic status. Overall, approximately 14% of Non-Hispanic Black women give birth before term compared with 9% of Non-Hispanic White women.⁶⁹ In a systematic review of 30 studies, Non-Hispanic Black women were found to have a 2-fold increased risk of PTB compared with Non-Hispanic White women.⁷⁴ The PTB rates for American Indian or Alaska Native women (11.5%) are higher than for White women.⁶⁸

Our previous work,⁷⁵ along with other reports^{76–78} but not all,⁷⁹ has suggested that psychosocial stress is a factor

associated with the risk of PTB. This association seems particularly strong when looking at first- and second-trimester stress^{78,80}; in these studies, women with increased life event stress during the first trimester of pregnancy had 2.40 times the risk of PTB compared with women with average or below-average stress.⁸⁰ Other studies show significant associations between experiences of racial discrimination and PTB.^{81–83} More specifically, women endorsing experiences of racism vs those without had 1.29 times the odds of experiencing PTB.⁸⁴ There is a growing recognition that exposure to racism rather than race itself is a risk factor for adverse maternal health outcomes.⁸⁵

As articulated in 2 recent articles on reproductive justice,^{86,87} prenatal care must become equitable and antiracist to interrupt the long arc of discrimination-based stress experiences, which manifest in adverse health, pregnancy, and birth outcomes.⁷³ Specific recommendations that should be tested and scaled include educating all providers about the history and systems that contribute to the social determinants of health, the early origins of obstetrical research based on the exploitation of the Black female body, how power dynamics impact care, cultural competency and humility, and the new approach of decolonized care that considers individual experiences and social-historic systems of oppression as affecting health.^{6,7,87} With training in cultural competency and humility, active listening skills, and shared decision-making, providers can be expected to commit to lifelong self-evaluation and critique to readdress new perspectives on the delivery of prenatal care as it evolves.⁸⁶ Although many factors contribute to the racial and ethnic disparities in the PTB rate, the problem will not be solved without promoting an antiracist culture and making equitable, respectful, supportive care routine. Expanding access to group prenatal care can foster provider-patient engagement and thereby reduce clinician bias, build patient trust with providers, lower patient stress, and increase patient social support across diverse populations, positive changes

that can contribute to healthier birth outcomes.^{73,86,88}

The psychological birth of the parent Opportunities and risk during the transition to parenthood

The TtP typically does not garner anywhere near the research and clinical attention as adolescence.⁸⁹ However, new parenthood is increasingly compared with our current understanding of the teenage years, even at the brain level. Both periods are transitional, marked by disequilibrium and iterative developmental adaptation, learning, and resiliency.^{17,90,91} Recent brain MRI research following female adolescents, first-time pregnant-to-postpartum women, and never-pregnant women for 2 years found that both the adolescents and pregnant-to-postpartum women show comparable reductions in cerebral gray matter and decreases in cortical thickness compared with minimal difference in the never-pregnant group.⁹¹ Elicited by hormonal fluctuations and the evolving role of parenting, these changes reflect neurocognitive adaptations and the brain's plasticity and also indicate increased vulnerability to pathologic alterations. Both adolescence and the perinatal period are times of increased risk of the emergence of mental health problems.⁹² In studies comparing men who are fathers and not fathers, the preconception-to-postpartum period also is marked by reductions in cortical volume and thickness.⁹³ Much less is known about the TtP for sexual and gender-diverse childbearing parents (and partners), single parents, and other nontraditional parenting structures; there is significant urgency to include these other parents in perinatal research to better understand their experiences and better support them.⁹⁴ Overall, calling attention to the TtP as comparable to other sensitive periods in brain development and similar to adolescence in its dramatic changes and openness to experience—contributing to both vulnerability and considerable opportunity—could validate expectant parents' experiences and underscore the significance of their emerging parenting roles.

Managing expectations

The demands on new parents are numerous and often overwhelming, with financial hardship exacerbating these challenges. Challenges include role and time management conflicts; diminished autonomy; shifting identity; increased financial concerns; lower self-esteem, particularly for the pregnant parent facing an altered physical appearance and abilities; and decreased relationship satisfaction as adult relationships become secondary to child-care (as summarized in a 2018 review by Saxbe et al¹⁷). However, not all new parents respond to the demands in the same way. Studies show that parents' prenatal expectations of these postpartum experiences impact intrapersonal adjustment to parenting and interpersonal couple functioning.^{95,96} In a recent qualitative study with heterosexual couples, individuals whose anticipations included "expecting the worst" (often an adaptive cognitive strategy to feel protected from disappointment) reported an easier adjustment once the baby arrived compared with those who treated the impending arrival of their baby as "unknowable" (something for which one could not plan or even envision). Embracing the unknown or tolerating uncertainty as a future parent also is associated with healthy adjustment.⁹⁷ Couples who envisioned the changed future with a child necessitating a team effort reported greater marital satisfaction compared with those who assumed the couple relationship would buffer child-related stressors.⁹⁶ In another report with heterosexual couples, having higher prenatal expectations relative to experiences once the baby was born (eg, about being a parent, competency as a parent, outside help and emotional support, and satisfactory division of family responsibilities) was associated with greater declines in levels of marital satisfaction after birth.⁹⁵

To promote family well-being and associated healthy infant development, the prenatal care setting potentially offers the unique opportunity to help prepare parents for the social-emotional changes, challenges, and expectations associated with impending parenthood.

New content could include authentic information on the parenting roles and life with a newborn because knowing what to expect and developing constructive mindsets promote a sense of agency and overall adjustment; specific and already-existing evidenced-based approaches to managing and adjusting expectations also could be taught.^{95,96,98}

Planning for daily life

Parenting an infant alters daily routines and complicates the establishment of new ones, particularly in the first months. Again, the context of poverty amplifies these difficulties. Key health behaviors, such as having adequate sleep, eating a healthy diet, and engaging in physical activity, can be frustratingly out of reach.¹⁷ Planning during the TtP for this level of disruption in daily habits and ideally problem-solving in advance can ease the challenges of early caregiving and potentially prevent depression, which is linked to sleep deprivation. Solutions to maintain consistent self-care can include calling on people in one's support network or accessing augmented care, such as a doula (a trained professional who can provide physical, emotional, and informational support before, during, and shortly after childbirth, facilitating opportunities for sleep, exercise, and decompression). Programs to broaden access to doula or similar care are garnering attention in states nationwide (<https://healthlaw.org/medical-aid-coverage-for-doula-care-2021-state-roundup/>). Additional research suggests that educating and preparing birthing people for the postpartum experience and promoting coping strategies can prevent depression among underserved mothers.⁹⁹ While feeding and/or pumping schedules can disrupt sleep, for some, anxiety and worrying can add to sleeping difficulties. To help with sleep disruption and insomnia, cognitive behavioral sleep techniques could be taught to expectant parents in prenatal care settings,^{100,101} and digital health may be a tool to promote broad dissemination (although outcome data must include attention to access issues for marginalized and less literate populations).

Intentional reflection on the parenting role

People often parent as they were parented or try to counter these influences. Envisioning and caring for a child are deeply influenced by one's own childhood experiences, even those earlier than explicit memory that nonetheless establish social-cognitive models of, and expectations for, caregiving relationships (as summarized in a 2020 article by Slade et al¹⁰²). In the extreme, children of maltreating parents are 2 times more likely to become maltreating parents themselves, although most children who were abused do not go on to abuse their children.¹⁰³ Even before the baby is born, elicited statements about caregiving reflect "internal working models of attachment" and predict qualities of future caregiving.^{104,105} For some parents to be, interventions that promote mentalization and perspective-taking—the ability to attend to another's mental states and subjective experiences—concerning their child are associated with more sensitive caregiving.¹⁰² As new content for the prenatal care setting, parents could be asked to reflect on how they were parented, what aspects of care they wish to repeat or change, and what strategies they can use to help them specify and realize their envisioned futures.

How to make change: designing and testing augmented prenatal care

Obstetrical visits typically last under 20 minutes; however, those with nurse-midwives are slightly longer.¹⁰⁶ Until recently, these sessions have been frequent (12–14 prenatal visits), with a call for added fourth-trimester sessions. Pandemic modifications may build on already existing initiatives to reduce the number of in-person prenatal visits for healthy pregnancies (<https://www.acog.org/education-and-events/creog/curriculum-resources/cases-in-high-value-care/expanding-prenatal-care-options-for-low-risk-patients>). Nonetheless, the TtP is a period of unprecedented frequent contact with health providers and the move to convenient, remote monitoring and text-based communication for some visits could align with stress-

reducing plans and other added information modules to help future parents and positively impact child outcomes during pregnancy and the fourth trimester of pregnancy.

The information we reviewed could form the basis for the development and testing of new content to be incorporated as routine, first-tier treatment for all birthing people using varied approaches. For example, the Mount Sinai Parenting Program (<https://parenting.mountsinai.org>) leverages web-based, parent-directed content, classes, and pediatric resident training to provide parenting information during routine well-child visits. A community resource, the Program for Early Parent Support (PEPS; <https://www.peps.org/about>), in Washington State, facilitates parent groups focused on practical information related to feeding and sleeping and parent wellness. To address social determinants of health, structural racism, mental health conditions, the prevention of postpartum depression, or the effects of chronic stress and trauma on perinatal individuals and their children would necessitate new provider training, interdisciplinary collaborations, and the development, testing, and incorporation of evidenced-based, triaged interventions based on level of need as exemplified by HealthySteps in pediatrics (<https://www.healthysteps.org>), which engages different behavioral health experts to provide psychosocial care for children and parents, or efforts to scale up existing evidenced-based approaches, such as Mothers & Babies (<https://www.motherandbabiesprogram.org/providers-interventionists/>); Reach Out, Stay Strong, Essentials for mothers of newborns⁵²; and Practical Resources for Effective Postpartum Parenting.^{107,108}

There is not yet an initiative to coordinate these types of programs, or to include the education and intervention strategies we described here based on DOHaD and TtP research, into a demonstration project testing augmented prenatal care models.¹⁰⁹ But, there should be.

Based on decades-long program development, implementation, and testing,¹⁰ healthcare systems are evolving to

support whole-person, 2-generation efforts in pediatrics. California recently became the first state to pay for dyadic care, dedicating \$800 million to cover HealthySteps for the 5.4 million children who have Medicaid insurance (http://www.physiciansnewsnetwork.com/la_county/california-could-be-first-state-to-implement-dyadic-care/article_7cf46842-e34a-11eb-8d68-1f0d8edb9934.html; https://cachildrenstrust.org/wp-content/uploads/2020/05/Dyadic_final_May2020.pdf). Pediatric care programs can serve as models for change in the perinatal care ecosystem, illustrating potential solutions to the myriad implementation challenges, such as provider time constraints, prioritization of patients' needs, budgetary challenges, and translation of knowledge into realistic practice guidelines.¹¹⁰ Pragmatic clinical trials (<https://rethinkingclinicaltrials.org>) and demonstration projects could be established to develop and test approaches to expanded education, intervention, prevention, and interdisciplinary services in prenatal care and generate evidence-based outcomes relevant to funding and scaling up these initiatives. To promote healthy families, we need to enhance healthcare at its origins—in obstetrics during the TtP. ■

REFERENCES

- Shonkoff JP, Boyce WT, Levitt P, Martinez FD, McEwen B. Leveraging the biology of adversity and resilience to transform pediatric practice. *Pediatrics* 2021;147:e20193845.
- Centers for Disease Control and Prevention. Data and statistics on children's mental health. 2021. Available at: www.cdc.gov/childrensmentalhealth/data.html. Accessed XXX.
- Schaeffer K. As schools shift to online learning amid pandemic, here's what we know about disabled students in the U.S. Pew Research Center. 2020. Available at: <https://www.pewresearch.org/fact-tank/2020/04/23/as-schools-shift-to-online-learning-amid-pandemic-heres-what-we-know-about-disabled-students-in-the-u-s/>. Accessed XXX.
- Heckman JJ, García JL. Social Policy: targeting programs effectively. *Nat Hum Behav* 2017;1:19.
- Knudsen EI, Heckman JJ, Cameron JL, Shonkoff JP. Economic, neurobiological, and behavioral perspectives on building America's future workforce. *Proc Natl Acad Sci U S A* 2006;103:10155–62.
- Barrera CM, Powell AR, Biermann CR, et al. A review of prenatal care delivery to inform the Michigan plan for appropriate tailored health-care in pregnancy panel. *Obstet Gynecol* 2021;138:603–15.
- Peahl AF, Zahn CM, Turrentine M, et al. The Michigan plan for appropriate tailored health-care in pregnancy prenatal care recommendations. *Obstet Gynecol* 2021;138:593–602.
- Garner A, Yogman M. Committee on Psychosocial Aspects of Child and Family Health, Section on Developmental and Behavioral Pediatrics, Council on Early Childhood. Preventing childhood toxic stress: partnering with families and communities to promote relational health. *Pediatrics* 2021;148:e2021052582.
- Beck AF, Edwards EM, Horbar JD, Howell EA, McCormick MC, Pursley DM. The color of health: how racism, segregation, and inequality affect the health and well-being of preterm infants and their families. *Pediatr Res* 2020; 87:227–34.
- Peacock-Chambers E, Ivy K, Bair-Merritt M. Primary care interventions for early childhood development: a systematic review. *Pediatrics* 2017;140:e20171661.
- Falk D. Evolution of brain and culture: the neurological and cognitive journey from Australopithecus to Albert Einstein. *J Anthropol Sci* 2016;94:99–111.
- Matthews LG, Walsh BH, Knutsen C, et al. Brain growth in the NICU: critical periods of tissue-specific expansion. *Pediatr Res* 2018;83: 976–81.
- Monk C, Lugo-Candelas C, Trumpff C. Prenatal developmental origins of future psychopathology: mechanisms and pathways. *Annu Rev Clin Psychol* 2019;15:317–44.
- Cardenas SI, Morris AR, Marshall N, et al. Fathers matter from the start: the role of expectant fathers in child development. *Child Dev Perspectives* 2022;16:54–9.
- Marshall NE, Abrams B, Barbour LA, et al. The importance of nutrition in pregnancy and lactation: lifelong consequences. *Am J Obstet Gynecol* 2022;226:607–32.
- Foster VA, Harrison JM, Williams CR, et al. Reimagining perinatal mental health: an expansive vision for structural change. *Health Aff (Millwood)* 2021;40:1592–6.
- Saxbe D, Rossin-Slater M, Goldenberg D. The transition to parenthood as a critical window for adult health. *Am Psychol* 2018;73: 1190–200.
- Scorza P, Duarte CS, Lee S, et al. Epigenetic intergenerational transmission: mothers' adverse childhood experiences and DNA methylation. *J Am Acad Child Adolesc Psychiatry* 2020;59:900–1.
- Rajyaguru P, Kwong ASF, Braithwaite E, Pearson RM. Maternal and paternal depression and child mental health trajectories: evidence from the Avon Longitudinal Study of Parents and Children. *BJPsych Open* 2021;7:e166.
- Dirix CE, Nijhuis JG, Jongsma HW, Hornstra G. Aspects of fetal learning and memory. *Child Dev* 2009;80:1251–8.

- 21.** Hepper PG. Fetal memory: does it exist? What does it do? *Acta Paediatr Suppl* 1996; 416:16–20.
- 22.** Kisilevsky BS, Hains SM, Lee K, et al. Effects of experience on fetal voice recognition. *Psychol Sci* 2003;14:220–4.
- 23.** Sandman CA, Wadhwa P, Hetrick W, Porto M, Peeke HV. Human fetal heart rate dis-habituation between thirty and thirty-two weeks gestation. *Child Dev* 1997;68:1031–40.
- 24.** Sandman CA, Wadhwa PD, Chicx-DeMet A, Porto M, Garite TJ. Maternal corticotropin-releasing hormone and habituation in the human fetus. *Dev Psychobiol* 1999;34:163–73.
- 25.** Moon C, Zernzach RC, Kuhl PK. Mothers say “baby” and their newborns do not choose to listen: a behavioral preference study to compare with ERP results. *Front Hum Neurosci* 2015;9:153.
- 26.** Siddappa AM, Georgieff MK, Wewerka S, Worwa C, Nelson CA, Deregnier RA. Iron deficiency alters auditory recognition memory in newborn infants of diabetic mothers. *Pediatr Res* 2004;55:1034–41.
- 27.** Busnel MC, Fifer W, Granier-Deferre C, et al. Tony DeCasper, the man who changed contemporary views on human fetal cognitive abilities. *Dev Psychobiol* 2017;59:135–9.
- 28.** DeCasper AJ, Fifer WP. Of human bonding: newborns prefer their mothers’ voices. *Science* 1980;208:1174–6.
- 29.** Uchida-Ota M, Arimitsu T, Tsuzuki D, et al. Maternal speech shapes the cerebral fronto-temporal network in neonates: a hemodynamic functional connectivity study. *Dev Cogn Neurosci* 2019;39:100701.
- 30.** Loomans EM, van Dijk AE, Vrijkotte TG, et al. Psychosocial stress during pregnancy is related to adverse birth outcomes: results from a large multi-ethnic community-based birth cohort. *Eur J Public Health* 2013;23:485–91.
- 31.** Long MM, Cramer RJ, Jenkins J, Bennington L, Paulson JF. A systematic review of interventions for healthcare professionals to improve screening and referral for perinatal mood and anxiety disorders. *Arch Womens Ment Health* 2019;22:25–36.
- 32.** Gavin NI, Gaynes BN, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T. Perinatal depression: a systematic review of prevalence and incidence. *Obstet Gynecol* 2005; 106:1071–83.
- 33.** Robinson AM, Benzie KM, Cairns SL, Fung T, Tough SC. Who is distressed? A comparison of psychosocial stress in pregnancy across seven ethnicities. *BMC Pregnancy Childbirth* 2016;16:215.
- 34.** Lomonaco-Haycraft KC, Hyer J, Tibbits B, et al. Integrated perinatal mental health care: a national model of perinatal primary care in vulnerable populations. *Prim Health Care Res Dev* 2018;20:e77.
- 35.** O’Brien AP, McNeil KA, Fletcher R, et al. New fathers’ perinatal depression and anxiety-treatment options: an integrative review. *Am J Mens Health* 2017;11:863–76.
- 36.** O’Donnell KJ, Glover V, Barker ED, O’Connor TG. The persisting effect of maternal mood in pregnancy on childhood psychopathology. *Dev Psychopathol* 2014;26: 393–403.
- 37.** Laplante DP, Brunet A, Schmitz N, Ciampi A, King S. Project Ice Storm: prenatal maternal stress affects cognitive and linguistic functioning in 5 1/2-year-old children. *J Am Acad Child Adolesc Psychiatry* 2008;47:1063–72.
- 38.** Sylvester CM, Myers MJ, Perino MT, et al. Neonatal brain response to deviant auditory stimuli and relation to maternal trait anxiety. *Am J Psychiatry* 2021;178:771–8.
- 39.** Weinstock M. Prenatal stressors in rodents: effects on behavior. *Neurobiol Stress* 2017;6:3–13.
- 40.** Darnaudéry M, Maccari S. Epigenetic programming of the stress response in male and female rats by prenatal restraint stress. *Brain Res Rev* 2008;57:571–85.
- 41.** Reynolds RM. Glucocorticoid excess and the developmental origins of disease: two decades of testing the hypothesis—2012 Curt Richter Award Winner. *Psychoneuroendocrinology* 2013;38:1–11.
- 42.** Matthey S, Barnett B, Howie P, Kavanagh DJ. Diagnosing postpartum depression in mothers and fathers: whatever happened to anxiety? *J Affect Disord* 2003;74:139–47.
- 43.** Milgrom J, Schembri C, Ericksen J, Ross J, Gemmill AW. Towards parenthood: an antenatal intervention to reduce depression, anxiety and parenting difficulties. *J Affect Disord* 2011;130:385–94.
- 44.** Goodman SH, Gotlib IH. Risk for psychopathology in the children of depressed mothers: a developmental model for understanding mechanisms of transmission. *Psychol Rev* 1999;106:458–90.
- 45.** May LE, Glaros A, Yeh HW, Clapp 3rd JF, Gustafson KM. Aerobic exercise during pregnancy influences fetal cardiac autonomic control of heart rate and heart rate variability. *Early Hum Dev* 2010;86:213–7.
- 46.** May LE, Knowlton J, Hanson J, et al. Effects of exercise during pregnancy on maternal heart rate and heart rate variability. *PM R* 2016;8:611–7.
- 47.** May LE, Scholtz SA, Suminski R, Gustafson KM. Aerobic exercise during pregnancy influences infant heart rate variability at one month of age. *Early Hum Dev* 2014;90:33–8.
- 48.** Thomas JC, Letourneau N, Bryce CI, Campbell TS, Giesbrecht GF. APRON Study Team. Biological embedding of perinatal social relationships in infant stress reactivity. *Dev Psychobiol* 2017;59:425–35.
- 49.** Zhao S, Xu C, Zhang R, Li X, Li J, Bao J. Effect of prenatal different auditory environment on learning ability and fearfulness in chicks. *Anim Biosci* 2022. [Epub ahead of print].
- 50.** Edvardsson K, Ivarsson A, Eurenus E, et al. Giving offspring a healthy start: parents’ experiences of health promotion and lifestyle change during pregnancy and early parenthood. *BMC Public Health* 2011;11:936.
- 51.** Dencker A, Premberg Å, Olander EK, et al. Adopting a healthy lifestyle when pregnant and obese - an interview study three years after childbirth. *BMC Pregnancy Childbirth* 2016;16: 201.
- 52.** O’Connor E, Senger CA, Henninger ML, Coppola E, Gaynes BN. Interventions to prevent perinatal depression: evidence report and systematic review for the US Preventive Services Task Force. *JAMA* 2019;321:588–601.
- 53.** Goodman SH, Cullum KA, Dimidjian S, River LM, Kim CY. Opening windows of opportunities: evidence for interventions to prevent or treat depression in pregnant women being associated with changes in offspring’s developmental trajectories of psychopathology risk. *Dev Psychopathol* 2018;30:1179–96.
- 54.** Dimidjian S, Goodman SH, Felder JN, Gallop R, Brown AP, Beck A. Staying well during pregnancy and the postpartum: a pilot randomized trial of mindfulness-based cognitive therapy for the prevention of depressive relapse/recurrence. *J Consult Clin Psychol* 2016;84:134–45.
- 55.** Sockol LE, Epperson CN, Barber JP. A meta-analysis of treatments for perinatal depression. *Clin Psychol Rev* 2011;31:839–49.
- 56.** Cooper PJ, Murray L, Wilson A, Romaniuk H. Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. *Br J Psychiatry* 2003;182:412–9.
- 57.** Nilini YI, Mehrizade A, Mayer L, Milanovic S. Treatment of depression, anxiety, and trauma-related disorders during the perinatal period: a systematic review. *Clin Psychol Rev* 2018;66:136–48.
- 58.** Leader LD, O’Connell M, VandenBerg A. Brexanolone for postpartum depression: clinical evidence and practical considerations. *Pharmacotherapy* 2019;39:1105–12.
- 59.** Milgrom J, Gemmill AW, Ericksen J, Burrows G, Buist A, Reece J. Treatment of postnatal depression with cognitive behavioural therapy, sertraline and combination therapy: a randomised controlled trial. *Aust N Z J Psychiatry* 2015;49:236–45.
- 60.** Moore Simas TA, Flynn MP, Kroll-Desrosiers AR, et al. A systematic review of integrated care interventions addressing perinatal depression care in ambulatory obstetric care settings. *Clin Obstet Gynecol* 2018;61:573–90.
- 61.** Byatt N, Biebel K, Lundquist RS, et al. Patient, provider, and system-level barriers and facilitators to addressing perinatal depression. *J Reprod Infant Psychol* 2012;30:436–49.
- 62.** Salameh TN, Hall LA, Crawford TN, Staten RR, Hall MT. Racial/ethnic differences in mental health treatment among a national sample of pregnant women with mental health and/or substance use disorders in the United States. *J Psychosom Res* 2019;121:74–80.
- 63.** Miller ES, Grobman WA, Ciolino JD, et al. Increased depression screening and treatment recommendations after implementation of a

perinatal collaborative care program. *Psychiatr Serv* 2021;72:1268–75.

64. Miller ES, Jensen R, Hoffman MC, et al. Implementation of perinatal collaborative care: a health services approach to perinatal depression care. *Prim Health Care Res Dev* 2020;21:e30.

65. Byatt N, Xu W, Levin LL, Moore Simas TA. Perinatal depression care pathway for obstetric settings. *Int Rev Psychiatry* 2019;31:210–28.

66. Byatt N, Straus J, Stopa A, Biebel K, Mittal L, Moore Simas TA. Massachusetts Child Psychiatry Access Program for Moms: utilization and quality assessment. *Obstet Gynecol* 2018;132:345–53.

67. Poleshuck EL, Woods J. Psychologists partnering with obstetricians and gynecologists: meeting the need for patient-centered models of women's health care delivery. *Am Psychol* 2014;69:344–54.

68. Martin JA, Hamilton BE, Osterman MJK, Driscoll AK. Births: final data for 2018. *Natl Vital Stat Rep* 2019;68:1–47.

69. Martin JA, Hamilton BE, Osterman MJK, Driscoll AK. Births: final data for 2019. *Natl Vital Stat Rep* 2021;70:1–51.

70. Luyckx VA. Preterm birth and its impact on renal health. *Semin Nephrol* 2017;37:311–9.

71. Hodyl NA, Aboustate N, Bianco-Miotto T, Roberts CT, Clifton VL, Stark MJ. Child neurodevelopmental outcomes following preterm and term birth: what can the placenta tell us? *Placenta* 2017;57:79–86.

72. Franz AP, Bolat GU, Bolat H, et al. Attention-deficit/hyperactivity disorder and very preterm/very low birth weight: a meta-analysis. *Pediatrics* 2018;141:e20171645.

73. Braveman P, Dominguez TP, Burke W, et al. Explaining the black-white disparity in preterm birth: a consensus statement from a multidisciplinary scientific work group convened by the March of Dimes. *Front Reprod Health* 2021;3:1–24.

74. Manuck TA. Racial and ethnic differences in preterm birth: a complex, multifactorial problem. *Semin Perinatol* 2017;41:511–8.

75. Walsh K, McCormack CA, Webster R, et al. Maternal prenatal stress phenotypes associate with fetal neurodevelopment and birth outcomes. *Proc Natl Acad Sci U S A* 2019;116(23):996–4005.

76. Szegda K, Bertone-Johnson ER, Pekow P, et al. Prenatal perceived stress and adverse birth outcomes among Puerto Rican women. *J Womens Health (Larchmt)* 2018;27:699–708.

77. Gillespie SL, Christian LM, Alston AD, Salisbury PJ. Childhood stress and birth timing among African American women: cortisol as biological mediator. *Psychoneuroendocrinology* 2017;84:32–41.

78. Hoffman MC, Mazzoni SE, Wagner BD, Laudenslager ML, Ross RG. Measures of maternal stress and mood in relation to preterm birth. *Obstet Gynecol* 2016;127:545–52.

79. Lima SAM, El Dib RP, Rodrigues MRK, et al. Is the risk of low birth weight or preterm labor greater when maternal stress is experienced during pregnancy? A systematic review and meta-analysis of cohort studies. *PLoS One* 2018;13:e0200594.

80. Zhu P, Tao F, Hao J, Sun Y, Jiang X. Prenatal life events stress: implications for preterm birth and infant birthweight. *Am J Obstet Gynecol* 2010;203:34.e1–8.

81. Giurgescu C, McFarlin BL, Lomax J, Craddock C, Albrecht A. Racial discrimination and the black-white gap in adverse birth outcomes: a review. *J Midwifery Womens Health* 2011;56:362–70.

82. Alhusen JL, Bower KM, Epstein E, Sharps P. Racial discrimination and adverse birth outcomes: an integrative review. *J Midwifery Womens Health* 2016;61:707–20.

83. Dominguez TP, Dunkel-Schetter C, Glynn LM, Hobel C, Sandman CA. Racial differences in birth outcomes: the role of general, pregnancy, and racism stress. *Health Psychol* 2008;27:194–203.

84. Bower KM, Geller RJ, Perrin NA, Alhusen J. Experiences of racism and preterm birth: findings from a pregnancy risk assessment monitoring system, 2004 through 2012. *Womens Health Issues* 2018;28:495–501.

85. Glazer KB, Howell EA. A way forward in the maternal mortality crisis: addressing maternal health disparities and mental health. *Arch Womens Ment Health* 2021;24:823–30.

86. Capp AI. They Make You Feel Less of a Human Being™: understanding and responding to Milwaukee's racial disparity in infant mortality. *Matern Child Health J* 2022;26:736–46.

87. Matthews K, Morgan I, Davis K, Estriplett T, Perez S, Crear-Perry JA. Pathways to equitable and antiracist maternal mental health care: insights from black women stakeholders. *Health Aff (Millwood)* 2021;40:1597–604.

88. Carter EB, EleVATE Women Collaborative-Mazzoni SE. A paradigm shift to address racial inequities in perinatal healthcare. *Am J Obstet Gynecol* 2021;224:359–61.

89. National Academies of Sciences, Engineering, and Medicine. The promise of adolescence: realizing opportunity for all youth. Washington, DC: National Academies Press (US); 2019.

90. Athan AM. Reproductive identity: an emerging concept. *Am Psychol* 2020;75:445–56.

91. Carmona S, Martínez-García M, Paternina-Die M, et al. Pregnancy and adolescence entail similar neuroanatomical adaptations: a comparative analysis of cerebral morphometric changes. *Hum Brain Mapp* 2019;40:2143–52.

92. Bale TL, Epperson CN. Sex differences and stress across the lifespan. *Nat Neurosci* 2015;18:1413–20.

93. Paternina-Die M, Martínez-García M, Prentus C, et al. The paternal transition entails neuroanatomic adaptations that are associated

with the father's brain response to his infant cues. *Cereb Cortex Commun* 2020;1:tgaa082.

94. Johansson M, Wirén AA, Ssempera D, Wells M. A qualitative study of midwives' challenges to support transmen during childbirth: a short report. *Eur J Midwifery* 2020;4:3.

95. Lawrence E, Nysten K, Cobb RJ. Prenatal expectations and marital satisfaction over the transition to parenthood. *J Fam Psychol* 2007;21:155–64.

96. Riggs DW, Worth A, Bartholomaeus C. The transition to parenthood for Australian heterosexual couples: expectations, experiences and the partner relationship. *BMC Pregnancy Childbirth* 2018;18:342.

97. Staneva A, Wittkowski A. Exploring beliefs and expectations about motherhood in Bulgarian mothers: a qualitative study. *Midwifery* 2013;29:260–7.

98. Welch T, Rouleau-Mitchell E, Farero A, Lachmar EM, Wittenborn AK. Maintaining relationship quality during the transition to parenthood: the need for next generation interventions. *Contemp Fam Ther* 2019;41:211–8.

99. Howell EA, Balbierz A, Wang J, Parides M, Zlotnick C, Leventhal H. Reducing postpartum depressive symptoms among Black and Latina mothers: a randomized controlled trial. *Obstet Gynecol* 2012;119:942–9.

100. Morgenthaler T, Kramer M, Alessi C, et al. Practice parameters for the psychological and behavioral treatment of insomnia: an update. An American Academy of Sleep Medicine report. *Sleep* 2006;29:1415–9.

101. Manber R, Bei B, Simpson N, et al. Cognitive behavioral therapy for prenatal insomnia: a randomized controlled trial. *Obstet Gynecol* 2019;133:911–9.

102. Slade A, Holland ML, Ordway MR, et al. Minding the Baby®: enhancing parental reflective functioning and infant attachment in an attachment-based, interdisciplinary home visiting program. *Dev Psychopathol* 2020;32:123–37.

103. Madigan S, Cyr C, Eirich R, et al. Testing the cycle of maltreatment hypothesis: meta-analytic evidence of the intergenerational transmission of child maltreatment. *Dev Psychopathol* 2019;31:23–51.

104. Berlin LJ, Dodge KA, Reznick JS. Examining pregnant women's hostile attributions about infants as a predictor of offspring maltreatment. *JAMA Pediatr* 2013;167:549–53.

105. Terry M, Finger B, Lyons-Ruth K, Sadler LS, Slade A. Hostile/helpless maternal representations in pregnancy and later child removal: a pilot study. *Infant Ment Health J* 2021;42:60–73.

106. Tai-Seale M, McGuire TG, Zhang W. Time allocation in primary care office visits. *Health Serv Res* 2007;42:1871–94.

107. Werner EA, Gustafsson HC, Lee S, et al. PREPP: postpartum depression prevention through the mother-infant dyad. *Arch Womens Ment Health* 2016;19:229–42.

108. Scorza P, Monk C, Lee S, Feng T, Berry OO, Werner E. Preventing maternal mental health disorders in the context of poverty: pilot efficacy of a dyadic intervention. *Am J Obstet Gynecol MFM* 2020;2:100230.

109. Schulz MS, Cowan CP, Cowan PA. Promoting healthy beginnings: a randomized controlled trial of a preventive intervention to preserve marital quality during the transition to parenthood. *J Consult Clin Psychol* 2006;74:20–31.

110. Molinaro ML, Evans M, Regnault TRH, de Vrijer B. Translating developmental origins of health and disease in practice: health care providers' perspectives. *J Dev Orig Health Dis* 2021;12:404–10.