

2005

Replication protein A-directed unloading of PCNA by the Ctf18 cohesion establishment complex

Goran O. Bylund

Washington University School of Medicine in St. Louis

Peter M. J. Burgers

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Bylund, Goran O. and Burgers, Peter M. J., "Replication protein A-directed unloading of PCNA by the Ctf18 cohesion establishment complex." *Molecular and Cellular Biology*.25,13. 5445-5455. (2005).
https://digitalcommons.wustl.edu/open_access_pubs/2248

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact engeszer@wustl.edu.

**Replication Protein A-Directed Unloading of
PCNA by the Ctf18 Cohesion Establishment
Complex**

Göran O. Bylund and Peter M. J. Burgers
Mol. Cell. Biol. 2005, 25(13):5445. DOI:
10.1128/MCB.25.13.5445-5455.2005.

Updated information and services can be found at:
<http://mcb.asm.org/content/25/13/5445>

REFERENCES

These include:

This article cites 51 articles, 37 of which can be accessed free
at: <http://mcb.asm.org/content/25/13/5445#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Replication Protein A-Directed Unloading of PCNA by the Ctf18 Cohesion Establishment Complex

Göran O. Bylund and Peter M. J. Burgers*

*Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine,
660 S. Euclid, St. Louis, Missouri 63110*

Received 3 March 2005/Returned for modification 29 March 2005/Accepted 31 March 2005

The replication clamp PCNA is loaded around DNA by replication factor C (RFC) and functions in DNA replication and repair. Regulated unloading of PCNA during the progression and termination of DNA replication may require additional factors. Here we show that a *Saccharomyces cerevisiae* complex required for the establishment of sister chromatid cohesion functions as an efficient unloader of PCNA. Unloading requires ATP hydrolysis. This seven-subunit Ctf18-RFC complex consists of the four small subunits of RFC, together with Ctf18, Dcc1, and Ctf8. Ctf18-RFC was also a weak loader of PCNA onto naked template-primer DNA. However, when the single-stranded DNA template was coated by the yeast single-stranded DNA binding protein replication protein A (RPA) but not by a mutant form of RPA or a heterologous single-stranded DNA binding protein, both binding of Ctf18-RFC to substrate DNA and loading of PCNA were strongly inhibited, and unloading predominated. Neither yeast RFC itself nor two other related clamp loaders, containing either Rad24 or Elg1, catalyzed significant unloading of PCNA. The Dcc1 and Ctf8 subunits of Ctf18-RFC, while required for establishing sister chromatid cohesion *in vivo*, did not function specifically in PCNA unloading *in vitro*, thereby separating the functionality of the Ctf18-RFC complex into two distinct paths.

The process of sister chromatid cohesion ensures that replicated chromosomes are distributed equally to progeny cells during cell division. Cohesion is mediated through a large ring-like structure, cohesin, which is deposited on the chromosomes in the late G₁ phase of the cell cycle. Sister chromatid cohesion is established during S phase, presumably by the actual passage of the replication fork. Several models exist by which cohesin is proposed to interact with the duplicated chromosomes in order to mediate cohesion. These models are based either on the idea that one cohesin ring encircles both sister chromosomes or on the idea that cohesion is mediated by the interlocking of two cohesin rings, with each ring more peripherally associated with a daughter chromosome (reviewed in references 33, 34, and 48). If the first type of model is correct, the mere passage of the replication fork through a cohesin ring would invariably ensure that the two sister chromatids stay attached until mitosis. However, the disadvantage of this elegant solution to the chromosome sorting problem is that it may be problematic for the replication fork to pass through the estimated 30- to 40-nm hole of the cohesin ring. Recent studies have shown that cohesin is specifically redistributed along the chromosome to transcription termination sites in a transcription-dependent manner, suggesting that the actual transcription machinery may push the cohesin rings ahead of the transcription bubble (26). If this is caused by steric problems because of the size of the transcription apparatus, similar steric problems may also occur with passage of the replication fork. Failure to establish sister chromatid cohesion would result if fork passage were sterically restricted.

Therefore, it is reasonable to assume that a dedicated apparatus exists to mediate passage of the replication fork through the cohesin ring, and the successful establishment of cohesion depends on the intimate coordination between the establishment and replication machineries.

There are several links between DNA replication and the establishment of sister chromatid cohesion, although a model describing the mechanistic rationale for these links is still lacking. First, the Ctf7 acetyltransferase is required for the establishment of chromatid cohesion (20, 44, 46). Temperature-sensitive mutations in the essential *CTF7* gene are synthetic lethal with conditional mutations in *POL30* (PCNA), while *POL30* overexpression suppresses the temperature sensitivity of *CTF7* mutants (44). Second, mutations in the *POL2* gene for the leading-strand DNA polymerase ϵ show cohesion establishment defects (9). Third, the cohesion establishment protein Ctf4 has been identified as a DNA polymerase α -interacting protein and *CTF4* mutants show replication-related defects (12, 19). Fourth, cohesion establishment factors Ctf18, Dcc1, and Ctf8 form a complex with the four small subunits (Rfc2-5) of replication factor C (RFC) (30). A temperature-sensitive mutation in *RFC4* showed defects in the establishment of chromatid cohesion and resulted in premature sister chromatid separation (30).

RFC loads PCNA onto DNA in an ATP-dependent manner. This RFC-dependent loading is essential for all PCNA-related processes during DNA replication and repair (reviewed in reference 27). It is not entirely clear whether RFC can also efficiently unload PCNA from DNA, either during DNA replication or after completion of DNA replication or repair. During lagging-strand DNA replication, unloading of PCNA by RFC from one completed Okazaki fragment may provide the required PCNA for the next Okazaki fragment to be synthesized. However, proof for this model is still lacking. Re-

* Corresponding author. Mailing address: Washington University School of Medicine, Department of Biochemistry, 660 S. Euclid, St. Louis, MO 63110. Phone: (314) 362-3872. Fax: (314) 362-7183. E-mail: burgers@biochem.wustl.edu.

modeling of the replication fork during cellular stress, DNA damage, or as a more regular aspect of replication progression may also depend on the temporary unloading of PCNA. Unloading of PCNA by RFC has been demonstrated in the human system. RFC mediates unloading of PCNA from fully replicated simian virus 40 chromatin (40). In model loading/unloading systems, human RFC has been shown to unload PCNA from template-primer DNA in an ATP-dependent reaction (6, 50). However, unloading efficiencies and rates, and therefore, physiological relevance of this type of unloading, have not been addressed. Curiously, in several unpublished studies, we have never been able to detect any significant PCNA unloading catalyzed by *Saccharomyces cerevisiae* RFC.

Several RFC-like complexes have recently been identified, all of which retain the four small subunits of RFC: the Rfc2-5 core. A complex of Rfc2-5 with *S. cerevisiae* Rad24 (Rad17 in human and *Schizosaccharomyces pombe*) functions in the DNA damage checkpoint, whereas the chromosome maintenance function of a third complex, consisting of Elg1 and Rfc2-5, has been less clearly defined (reviewed in reference 27). Finally, a complex consisting of the Rfc2-5 core, Ctf18, Dcc1, and Ctf8 (Ctf18-RFC) is required for the establishment of sister chromatid cohesion in yeast (19, 30). This seven-subunit complex is conserved from yeast to mammals. In an effort to understand the function of this cohesion-related clamp loader, we have purified this complex and studied its biochemical properties. Surprisingly, we find that Ctf18-RFC efficiently unloads PCNA from DNA. Unloading requires the energy of ATP hydrolysis. We discuss our results in light of two recent reports that conclude that the human Ctf18-RFC is primarily a loader of PCNA (2, 41). Furthermore, we propose that regulated unloading of PCNA may serve an important function during the establishment of sister chromatid cohesion.

MATERIALS AND METHODS

Strains, plasmids, DNA, and proteins. *S. cerevisiae* strain BJ2168 (*MATa ura3-52 trp1-289 leu2-3,112 prb1-1122 prc1-407 pep4-3*) was used for overexpression purposes. Complementation studies were carried out in strain Y06212 (*mata his3-Δ1 leu2-Δ0 met15-Δ0 ura3-Δ0 ctf18-Δ::kanMX4*) from Euroscarf. All genes to be expressed and purified were cloned into multicopy yeast shuttle vectors under the control of the galactose-inducible bidirectional *GALI-10* promoter (4). Plasmid pBL438 (Bluescript, 2 μ m ori, *URA3*, M13 ori, *GALI-10 GST-CTF18*) was constructed by PCR amplifying the *CTF18* gene from chromosomal yeast DNA and cloning it into pRS426-GAL, followed by N-terminal fusion of the glutathione *S*-transferase (*GST*) gene from pGEX-6P-1 containing a PreScission protease site (Amersham Pharmacia). PreScission treatment of the *GST*-Ctf18 polypeptide resulted in a Ctf18 protein with the two first amino acids (MV) replaced by an N-terminal four-amino-acid extension (GPKL). Plasmid pBL441 was derived from pBL438 by additionally inserting the *CTF8* gene under *GALI-10* control, counterclockwise to *GST-CTF18*. The *ctf18-K189E* mutation was introduced in this plasmid (pBL441-KE) by site-directed mutagenesis. Plasmid pBL441-ΔC82 contained a frameshift mutation resulting in an 82-amino-acid C-terminal truncation of the *CTF18* gene. The *DCC1* gene was similarly cloned into pRS425-GAL, yielding pBL440 (Bluescript, 2 μ m ori, *LEU2*, M13 ori, *GALI-10 DCC1*). The *ELG1* gene was cloned as a *GST-ELG1* fusion exactly analogously to *CTF18* into plasmid pRS426-GAL, yielding pBL448 (Bluescript, 2 μ m ori, *URA3*, M13 ori, *GALI-10 GST-ELG1*). PreScission treatment of the *GST*-Elg1 polypeptide resulted in an N-terminal four-amino-acid extension (GPEF). An *Escherichia coli* expression plasmid, pBL446, contained both the *DCC1* gene and the His₇-tagged *CTF8* gene cloned under control of the bacteriophage T7 promoter. DNA sequencing of all constructed plasmids was performed to verify the clones.

Bluescript SKII⁺ single-stranded DNA (ssDNA), prepared from *E. coli* using phagemid technology, was primed with 10 30-mer primers, roughly equally spaced along the circle (deca-primed DNA). A 2-kb plasmid containing a BbvC1

restriction endonuclease site (pUC19B1) was treated with either the BbvC1A or BbvC1B nickase (New England Biolabs) and reisolated by phenol extraction and ethanol precipitation to give nicked DNA.

Replication protein A (RPA), *E. coli* SSB, DNA polymerase δ (Pol δ), Rad24-RFC, Rfc2-5, and PCNA were purified as described previously (5, 16, 28). Most biochemical experiments in this paper were carried out with a form of RFC lacking the N-terminal 272-amino-acid domain that has been shown to be dispensable for clamp loading both in vitro and in vivo (16). Where indicated, full-length wild-type RFC isolated from yeast was used in unloading assays (14). PCNA containing an N-terminal phosphorylatable tag (RRASVGS) was ³²P labeled (200 to 500 cpm/fmol) with the catalytic subunit of cAMP-dependent protein kinase (NE Biolabs) as described previously (10). RPA-1ΔN lacking the N-terminal 180-amino-acid domain of Rpa1 was overproduced in *E. coli* and purified as described previously (1). T4 protein gp32 was purchased from USB (Cleveland, OH).

Overexpression and purification of clamp loaders. Strain BJ2168 was transformed with plasmid pBL422 (2 μ M ori *TRP1 RFC2 RFC3 RFC4 RFC5*) with the four small-subunit genes under control of the *GALI-10* promoter (14). This strain was transformed with pBL438 to produce the five-subunit Ctf18-RFC, with pBL441 and pBL440 to produce the seven-subunit Ctf18-RFC, and with pBL448 to produce Elg1-RFC. Cell growth, induction with galactose, and crude extract preparation by dry ice breakage of 150 g of yeast cell paste were carried out as described previously (4). The crude lysate was adjusted to 150 mM ammonium sulfate and stirred with 45 μ l/ml of 10% Polymin P for 15 min at 0°C to precipitate nucleic acids. After centrifugation at 39,000 \times g for 45 min, protein in the supernatant was precipitated with 0.35 g/ml of solid ammonium sulfate, followed by centrifugation at 39,000 \times g for 45 min. Protein pellets were resuspended and diluted in buffer A₀ (50 mM HEPES-NaOH [pH = 7.5], 10% glycerol, 1 mM EDTA, 1 mM EGTA, 0.05% Tween 20, 2.5 mM sodium pyrophosphate, 1 mM β -glycerophosphate, 3 mM dithiothreitol, 10 mM NaHSO₃, 5 μ M pepstatin A, 10 μ M leupeptin, 2.5 mM benzamidine, 5 μ g/ml chymostatin, 0.5 mM phenylmethylsulfonyl fluoride) to a conductivity equal to that of A₁₅₀ (buffer with 150 mM NaCl).

Glutathione-Sepharose (5 ml; Amersham Pharmacia) equilibrated in A₁₅₀ was added to the protein extract, and proteins were batch bound at 4°C for 3 h on a rotator. The Sepharose was washed with 20 volumes of A₁₅₀, and protein was eluted with buffer B₁₅₀ (buffer A at pH 8.0 with 150 mM NaCl, 20 mM glutathione, and 0.05% ampholytes [pH = 3 to 10] [Amersham Pharmacia] and no benzamidine or phenylmethylsulfonyl fluoride). Clamp loader containing fractions were identified by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and treated with PreScission protease on ice overnight (Amersham Pharmacia).

Protein fractions from the affinity column were diluted with C₀ (buffer B, pH 7.5, without glutathione) to a conductivity equivalent to that of C₁₀₀ before further purification. Ctf18-RFC(7) and Elg1-RFC were further purified on a 1-ml monoS column, while Ctf18-RFC(5) was further purified on a 5-ml heparin agarose column.

The His₇-Ctf8-Dcc1 heterodimer was overproduced from plasmid pBL446 in *E. coli* strain BL21(DE3)-RIL using a standard induction protocol at 20°C to improve complex solubility and purified using Ni-agarose chromatography. The protein was eluted with a 100 mM imidazole buffer and further purified by MonoS fast-performance liquid chromatography.

Reconstitution of the seven-subunit Ctf18-RFC complex. His₇-Ctf8-Dcc1 (50 pmol) was incubated with 20 pmol of either Ctf18-RFC(5), Ctf18-ΔC82(5), or Rfc2-5 in 100 μ l of buffer C₂₀₀ containing 7 mM MgAc₂ and 1 mM ATP where indicated for 1 h at 0°C. The reaction mixture was further incubated for 1 h at +4°C on a shaker with 100 μ l Ni²⁺-nitrilotriacetic acid (NTA) agarose (QIAGEN), previously equilibrated in buffer C₃₀₀ containing 20 mM imidazole. The Ni²⁺-NTA agarose, with bound protein, was washed three times with 200 μ l of buffer C₃₀₀ containing 20 mM imidazole before protein was eluted with 200 μ l of buffer C₃₀₀ containing 100 mM imidazole. The protein composition of the eluates was precipitated with 10% trichloroacetic acid and analyzed by 15% SDS-PAGE and Coomassie staining. No binding of clamp loader complexes to the Ni²⁺-NTA agarose was observed when the procedure was repeated in the absence of His₇-Ctf8-Dcc1.

ATPase assays. Assays (25 μ l) in buffer H₇₅ (30 mM HEPES-NaOH [pH 7.5], 0.5 mM EDTA, 10% glycerol, 10 mM magnesium acetate, 5 mM dithiothreitol, 0.1% ampholytes pH3 to 10], 0.01% Nonidet P-40, 10 μ g/ml bovine serum albumin, 75 mM NaCl) contained 100 μ M [α -³²P]ATP, 60 fmol of deca-primed bluescript SKII⁺ ssDNA (600 fmol primer termini), 7 pmol RPA or RPA-1ΔN when present, 600 fmol PCNA, and 300 fmol of Ctf18-RFC. Reactions were incubated at 30°C, and after 2, 4, and 6 min, 5- μ l aliquots were removed, quenched with 2.5 μ l of 50 mM EDTA, 1% SDS, and 25 mM cold ATP and

ADP, and processed for polyethyleneimine cellulose thin-layer chromatography and phosphorimager analysis. Linear initial rates were calculated from each time course.

Isolation of DNA-PCNA complexes. The standard 100- μ l reaction mixture contained in buffer H₇₅ was as follows: 500 fmol of bluescript SKII⁺ ssDNA primed by 10 evenly spaced 30-mer primers (5 pmol primer termini), 60 pmol of RPA if present, 6 pmol of PCNA (containing when present 0.5 pmol of ³²P-PCNA as a radioactive tracer), 3 pmol of RFC or Ctf18-RFC, and 1 mM ATP, 5 min at room temperature. The mixture was filtered through a 2-ml Biogel A5m column equilibrated in buffer H₇₅ at room temperature. Four-drop fractions were collected; the void volume containing DNA-protein complexes eluted in fractions 9 to 12, and the unbound protein in fractions 16 to 22. After counting of the fractions, complexes were precipitated with an equal volume of 20% trichloroacetic acid and analyzed by SDS-PAGE and Coomassie staining and, after drying of the gel, by phosphorimaging analysis (see Fig. 3).

PCNA-nicked DNA complexes and PCNA-gapped DNA complexes used for our unloading studies were isolated similarly after loading ³²P-PCNA (2.5 pmol) onto 500 fmol of nicked pUC19B1 DNA by 200 fmol of RFC and 1 mM ATP at 30°C for 5 min. Complex-containing fractions were identified with a Geiger counter and immediately used further in unloading assays.

PCNA unloading assay using magnetic beads. Streptavidin-coated magnetic beads were charged with a 286-mer ssDNA via a 5'-terminal biotin linkage and primed with a 30-mer approximately in the middle of the template sequence (15). The standard 50- μ l loading assay contained in buffer UB (as buffer H, but 5 mM magnesium acetate and 200 μ g/ml bovine serum albumin) was as follows: 100 fmol DNA (~3 μ l beads), 2 pmol of RPA, 200 μ M ATP, 200 fmol ³²P-PCNA, and 400 fmol of RFC, 2 min at room temperature. The beads were pulled down to the magnet, the supernatant taken off, and the beads washed once with 150 μ l of UB by resuspension and pull-down. The beads were resuspended in 45 μ l UB and used in the unloading assay. For time courses, the loading reactions were scaled up accordingly.

The particular clamp loader (500 fmol) and either no nucleotide, 1 mM ATP, or 100 μ M ATP γ S in a total volume of 5 μ l was added to the beads, and after the indicated times at room temperature, the beads were pulled down for 15 s and the radioactivity in the supernatant and the beads was determined by scintillation counting.

PCNA unloading assay using glutaraldehyde cross-linking. ³²P-PCNA was loaded by RFC onto either RPA-coated deca-primed bluescript ssDNA, gapped bluescript DNA (90-nucleotide [nt] ssDNA gap), or nicked DNA, and the PCNA-DNA complexes were purified by BioGel-A5m gel filtration as described above. Complexes in buffer H₇₅ were incubated with RFC or Ctf18-RFC and the indicated nucleotide, and after the indicated time, the reaction was stopped by adding 1% final glutaraldehyde. After 30 to 60 min on ice, loading buffer containing 0.2% SDS and 10 mM EDTA (final concentrations) was added, and the samples were electrophoresed at 1 V/cm for 16 h on a 1 to 1.5% agarose gel in 50 mM Tris-borate-EDTA buffer. The gel was dried and radioactivity quantitated on a phosphorimager (see Fig. 5 and 6). Migration positions of glutaraldehyde-cross-linked PCNA and Ctf18-RFC-PCNA were determined independently.

Rates of disappearance of the DNA-PCNA complex were calculated using standard first-order kinetics, and these rates were taken to represent rates of PCNA unloading.

Quantitation and visualization techniques. All SDS-PAGE gels were stained with colloidal Coomassie and photographed with a charge-coupled-device camera under linear response conditions. Bands were quantitated using NIH-image software. Gels stained with SYPRO orange were quantitated using a STORM phosphorimager, using the blue fluorescence mode, and ImageQuant software (Molecular Dynamics). Dried gels containing ³²P-PCNA and thin-layer chromatography plates used in the ATPase assays were subjected to phosphorimaging analysis, and bands were quantitated using ImageQuant software. After quantitation, all gel files were contrast enhanced for visualization purposes using NIH-image and Canvas software.

RESULTS

Yeast clamp loaders have the Rfc2-5 core assembly in common. We overproduced the five-subunit Ctf18-RFC(5) complex in yeast from multicopy plasmids with each gene placed downstream of the galactose-inducible *GAL1-10* promoter. A cleavable GST tag was added to the *CTF18* gene to aid in purification (see Materials and Methods). Following the glu-




FIG. 1. Yeast RFC-like clamp loaders. The indicated complexes (2 μ g each) were separated by 10% SDS-PAGE (lanes 1 to 6) or 15% SDS-PAGE (in order to visualize Ctf8; lane 7), and the gel was stained with colloidal Coomassie. Migration positions of marker proteins are indicated between lanes 6 and 7. RFC-1 Δ N (lane 1) lacks a dispensable N-terminal 272-amino-acid domain and was used in most of the studies in this paper (see Materials and Methods).

tathione affinity column, the GST tag was proteolytically cleaved and the remaining complex further purified by HPLC ion exchange chromatography. The seven-subunit Ctf18-RFC(7) complex was similarly purified from a system in which in addition the *CTF8* and *DCC1* genes were overexpressed. We also overexpressed and purified the five-subunit Elg1-RFC complex using the same methodology as was used to obtain Ctf18-RFC(5). A comparative electrophoretic analysis of these purified clamp loaders with those previously isolated, i.e., RFC and Rad24-RFC, showed that all clamp loaders have the Rfc2-5 core in common. All subunits unique to the individual clamp loaders are present at equal stoichiometry, within an error of 20%, with the Rfc2-5 core (Fig. 1). The Ctf8 subunit of Ctf18-RFC was the only exception. Based on staining of the SDS-polyacrylamide gel, this subunit appeared to be overrepresented in the complex at a ratio of 1.6:1 (Coomassie staining) or 1.4:1 (SYPRO orange staining; data not shown) compared to Rfc2-5.

The seven-subunit complex could also be reconstituted by mixing the Ctf18(5) complex, purified from yeast, with Ctf8-Dcc1 overproduced in *E. coli*. Because only Ctf8 contained a heptahistidine tag, complex formation was simply assessed by determining retention of Ctf18-RFC(5) onto a Ni-agarose column. Retention of Ctf18-RFC(5) on this column did not require ATP (Fig. 2, lane 5 versus lane 6). The four-subunit Rfc2-5 core complex failed to interact with Dcc1-Ctf8 (lane 8). Successful reconstitution required the C-terminal 82-amino-acid tail of Ctf8. A Ctf8 truncation mutant lacking this tail was not retained on the column (lane 7). In addition, we were unable to purify a seven-subunit complex from yeast when *CTF18- Δ C82* replaced wild-type *CTF18* in our overexpression system (data not shown). This lack of interaction was unlikely caused by misfolding of Ctf18- Δ C82, because the truncated complex was fully active in PCNA loading and unloading (data not shown).




FIG. 2. The C-terminal tail of Ctf18 is required for binding the Ctf8 and Dcc1 subunits. Lanes 1 to 4 are the input complexes: lane 1, Ctf18-RFC(5); lane 2, Ctf18- Δ C82-RFC(5); lane 3, Rfc2-5; lane 4, His₇-Ctf8-Dcc1. Lanes 5 to 8 are the 100 mM imidazole eluates from the Ni²⁺-agarose column. Lane 5, Ctf18-RFC(5) incubated with His₇-Ctf8-Dcc1; lane 6, Ctf18-RFC(5) incubated with His₇-Ctf8-Dcc1 and 1 mM ATP; lane 7, Ctf18- Δ C82-RFC(5) incubated with His₇-Ctf8-Dcc1 and 1 mM ATP; lane 8, Rfc2-5 incubated with His₇-Ctf8-Dcc1 and 1 mM ATP. See Materials and Methods for details.

Since the two Ctf18-containing complexes, Ctf18-RFC(5) and Ctf18-RFC(7), showed very similar biochemical properties with regard to PCNA loading and unloading, primarily those studies carried out with Ctf18-RFC(7) will be reported. This seven-subunit complex will be referred to as Ctf18-RFC.

RPA inhibits PCNA loading by Ctf18-RFC. Recently a human Ctf18-RFC complex was identified, and two studies showed that this complex can function, albeit inefficiently, as a loader of PCNA (2, 32, 41). Moreover, this loaded PCNA acts as a processivity factor for DNA Pol δ , indicating that it is loaded in the proper orientation at the 3'-primer-template junction. This was surprising to us because we had consistently failed to observe evidence for a processive PCNA-Pol δ complex in a coupled replication assay with yeast Ctf18-RFC as the putative loader (data not shown). In this coupled assay, PCNA was loaded by the clamp loader of interest onto RPA-coated primed single-stranded DNA, and loading was assessed indirectly by measuring processive DNA synthesis by Pol δ . However, such a coupled assay, because of its indirectness, did not allow us to determine whether PCNA had actually been loaded or not, since it could be present on the DNA in a form unable to interact with Pol δ .

In order to determine PCNA loading directly, we separated the products of the loading reaction on a gel filtration column in order to separate DNA-bound PCNA from other forms of PCNA, either free or in a complex with the respective clamp loader (Fig. 3A). A radiolabeled form of PCNA was used to facilitate detection (10). We observed virtually no loading of PCNA by Ctf18-RFC onto multiply primed ssDNA, which was coated with RPA, the eukaryotic ssDNA binding protein. Surprisingly, a reasonable loading efficiency was observed when we carried out the reaction on naked DNA substrate. This loading of PCNA was ATP dependent. The observation that Ctf18-RFC loaded PCNA only onto naked template-primer DNA was very surprising, because in contrast, PCNA loading by its normal partner, RFC, proceeded with much higher efficiency when the ssDNA was coated with RPA, as shown before in many studies (Fig. 3A) (11, 25, 47, 51). The isolated DNA-




FIG. 3. RPA inhibits PCNA loading by Ctf18-RFC. (A) Deca-primed ssDNA was coated with RPA where indicated. Loading of ³²P-PCNA by RFC (open symbols) or Ctf18-RFC (filled symbols) and ATP where indicated and BioGel-A5m gel filtration were as described in Materials and Methods (see inset for a scheme of the assay). The amount of PCNA (fmol) in each fraction is indicated. The dashed line shows background loading of PCNA by RFC or by Ctf18-RFC without ATP present (results were identical). (B) DNA was incubated with increasing levels of RPA at 30°C for 10 min prior to loading, and gel filtration was done as for (A). The void volume fractions (DNA-PCNA) were acid precipitated and analyzed by SDS-PAGE as described in Materials and Methods. The gel was stained with colloidal Coomassie (upper panel), dried, and subjected to phosphorimager analysis (lower panel). Saturation binding of the ssDNA was achieved with ~45 pmol of RPA (22). The open arrow to the right indicates a proteolytic fragment of RPA. It migrates close to, but is distinct from, PCNA.

PCNA complex formed by Ctf18-RFC action was active for processive DNA replication upon addition of RPA and Pol δ , indicating that PCNA had been loaded properly for interaction with Pol δ (data not shown).

One possible explanation for these results may be that the PCNA-Ctf18-RFC complex is incapable of binding RPA-coated primed DNA. The deca-primed ssDNA was coated with increasing concentrations of RPA and subsequently subjected to a PCNA loading reaction by Ctf18-RFC. Proteins coeluting with the DNA were analyzed by SDS-PAGE and phosphorimager analysis (³²P-PCNA). Partial coating of the ssDNA actually resulted in an increased loading of PCNA even though binding of Ctf18-RFC to the DNA was slightly decreased (Fig.

TABLE 1. Specificity parameters for PCNA loading^a

Condition	SSB	Loader	Amt of PCNA loaded (fmol)
Deca-primed SKII+ ssDNA	None	RFC	400
	RPA	RFC	1,240
	RPA(1-ΔN)	RFC	1,820
	<i>E. coli</i> SSB	RFC	800
	T4 gp32	RFC	980
No ATP	None	Ctf18-RFC	10
	None	Ctf18-RFC	460
	None	Ctf18-RFC	410 ^b
	RPA	Ctf18-RFC	40
	RPA	Ctf18-RFC	30 ^b
	RPA(1-ΔN)	Ctf18-RFC	500
	<i>E. coli</i> SSB	Ctf18-RFC	300
	T4 gp32	Ctf18-RFC	320
Nicked pUC19B1 DNA	None	RFC	360
	None	Ctf18-RFC	7

^a All loading reactions were as described in Materials and Methods with 5 pmol template-primer termini, 6 pmol of PCNA, and 3 pmol of RFC or Ctf18-RFC, except where indicated.

^b Reaction mixtures contained 2.5 pmol of PCNA and 5 pmol of Ctf18-RFC.

3B). However, further coating of the ssDNA led to a dramatic decrease in Ctf18-RFC binding and an even more dramatic decrease in PCNA loading. At saturated RPA levels, PCNA loading was inhibited 82 to 96% compared to naked DNA (six determinations). In contrast, RFC-mediated loading of PCNA was strongly stimulated at all levels of RPA (Fig. 3A, data not shown).

We wondered whether the observed inhibition was specific for yeast RPA or whether any ssDNA binding protein would inhibit loading by Ctf18-RFC. The latter scenario would suggest that Ctf18-RFC preferred binding to a template-primer junction with adjacent naked ssDNA. Therefore, the loading reaction was carried out with several different ssDNA binding proteins, and the PCNA loaded was quantitated by gel filtration analysis, similarly to loading shown in Fig. 3A (Table 1). For each ssDNA binding protein, a plateau value of PCNA loading efficiency was reached upon saturation binding of the ssDNA, and addition of up to a twofold excess was without further effect. Surprisingly, while wild-type RPA strongly inhibited PCNA loading, a mutant RPA (RPA-1ΔN) lacking the N-terminal regulatory domain of Rpa1 (Δ1-180) allowed Ctf18-RFC binding to the DNA and completely supported PCNA loading even though this mutant coated the ssDNA as efficiently as wild-type RPA (Table 1, data not shown) (1). The

N-terminal 180-amino-acid domain of Rpa1 mediates many of the known interactions with other replication proteins including a direct interaction with Rfc4, a component of Ctf18-RFC (23). The implication of these experiments is that the interactions between the N terminus of Rpa1 and one or more subunits of Ctf18-RFC may serve to inhibit the PCNA loading function of this complex. Consistent with this hypothesis, we found that compared to RPA, two heterologous ssDNA binding proteins, *E. coli* SSB and bacteriophage T4 gp32, did not cause a strong inhibition of PCNA loading by Ctf18-RFC (Table 1). Control experiments showed that compared to naked DNA, all four ssDNA binding proteins stimulated PCNA loading by RFC, with RPA-1ΔN being the most stimulatory (Table 1).

We also determined whether Ctf18-RFC was able to load PCNA onto a nicked DNA substrate. Loading at a nick may be important for the proper function of several DNA metabolic pathways including base excision repair and nucleotide excision repair, and RFC was shown to be a proficient PCNA loader onto a nicked substrate (Table 1) (6, 37). In contrast, Ctf18-RFC completely failed to load PCNA onto a nick.

ATPase activities of clamp loaders. Table 2 summarizes the ATPase activities of all known clamp loaders, including the four-subunit Rfc2-5 core. The ATPase activity of RFC was stimulated about 20-fold when both PCNA and effector DNA were present in the assay, indicating efficient loading of PCNA. Similarly, the ATPase of Ctf18-RFC was also strongly stimulated by PCNA and DNA, but only when the ssDNA was naked or coated with RPA-1ΔN. In contrast, when the ssDNA was coated with RPA, the ATPase in the presence of cofactors was not significantly different from that without (Table 2). This lack of ATPase stimulation strongly suggests that Ctf18-RFC does not bind RPA-coated DNA and therefore does not load PCNA onto such a substrate.

Although different in quantitative detail, the ATPase activities of the other two clamp loaders, Rad24-RFC and Elg1-RFC, showed similar responses to the presence of PCNA and DNA as observed for the Rfc2-5 core. The ATPase activity of each of these three complexes was stimulated severalfold by addition of PCNA to the assay, indicating that these complexes do interact with PCNA (17). However, this ATPase was not further stimulated in the additional presence of template-primer DNA, suggesting that the bound PCNA was not loaded onto the DNA. Indeed, in PCNA-loading assays as described in the legend to Fig. 3A, neither Rad24-RFC nor Elg1-RFC showed evidence for stable loading of PCNA (data not shown) (28).

TABLE 2. ATPase activities of clamp loaders

Cofactor	Turnover ^a										
	RFC		Ctf18-RFC			Elg1-RFC		Rad24-RFC ^b		Rfc2-5	
	-RPA	+RPA	-RPA	+RPA	+RPA-1ΔN	-RPA	+RPA	-RPA	-RPA	+RPA	
None	6	4	32	35	33	3	4	13	15	22	
PCNA	12	10	30	34	29	21	23	42	35	41	
DNA	90	30	89	46	92	3	9	18	33	25	
DNA+PCNA	95	102	138	37	142	19	25	43	42	44	

^a Turnover numbers (ADP formed/ min/RFC) are given. Standard errors are 5%. See Materials and Methods for details.

^b Data for Rad24-RFC from reference 28.




FIG. 4. Unloading of PCNA from DNA by Ctf18-RFC. (A) Scheme of the assay. See Materials and Methods for details. (B) Unloading by Ctf18-RFC and Ctf18-RFC(5). Beads were incubated with Ctf18-RFC without nucleotide, with 1 mM ATP, or with 100 μ M ATP γ S. Bead-associated radioactivity was determined at the indicated times as described in Materials and Methods. (C) Mutant Ctf18-K189E replaced the wild type in Ctf18-RFC. (D) Successive 10-fold serial dilutions of a *ctf18*- Δ strain containing either empty vector (-), plasmid pBL441 (*CTF18*), pBL441-KE (*ctf18-K189E*), or pBL441- Δ C82 [*ctf18*- Δ (82)] were grown for 3 days at 30°C on yeast-peptone-dextrose plates or yeast-peptone-dextrose plates containing 10 μ g/ml of camptothecin. (E) Unloading of PCNA by other clamp loaders as described in Materials and Methods. The assays were carried out in duplicate in two or three independent assays, and the average is given. Standard errors ranged from 5 to 10% for all assays.

Unloading of PCNA by Ctf18-RFC. Rather than being a loader, we reasoned that Ctf18-RFC might actually be a PCNA unloader. To test this hypothesis, we developed a rapid and convenient assay for PCNA unloading. Radiolabeled PCNA was loaded by RFC onto RPA-coated template-primer DNA attached to magnetic beads, and RFC was removed by washing of the beads (17). PCNA unloading was determined in a subsequent assay as loss of bead-associated radioactivity (Fig. 4A).

PCNA unloading by Ctf18-RFC was observed in a reaction that required hydrolysis of ATP, since ATP γ S was inactive (Fig. 4B). Unloading did not require the Dcc1 and Ctf8 subunits of the seven-subunit complex, because a five-subunit clamp loader consisting of Ctf18 together with Rfc2-5, Ctf18-RFC(5), also unloaded PCNA in an ATP-dependent reaction (Fig. 4B). Furthermore, a C-terminal truncation mutant of Ctf18 abolished all interaction with Dcc1-Ctf8 (Fig. 2). The five-subunit Ctf18- Δ C82-RFC complex was also proficient in PCNA unloading, as well as in PCNA loading (data not shown).

However, an intact ATP-binding domain in Ctf18 was required for activity. We mutated the invariant lysine in the Walker A motif of this ATP-binding domain to glutamic acid. The purified complex failed to unload PCNA (Fig. 4C), nor was PCNA loading by this mutant complex observed (data not shown).

Defects in the establishment of sister chromatid cohesion are associated with sensitivity to the topoisomerase I drug camptothecin (35). When assayed in a yeast *ctf18* deletion strain, *ctf18-K189E* failed to rescue the camptothecin sensitivity of the deletion mutant (Fig. 4D). The C-terminal truncation *ctf18*- Δ C82 mutant also failed to rescue camptothecin sensitivity of the deletion strain, suggesting that the chromatid establishment function of the Dcc1 and Ctf8 subunits requires their participation in the seven-subunit Ctf18-RFC complex.

None of the other clamp loaders was able to unload PCNA. Addition of RFC and ATP or ATP γ S in the assay merely resulted in a reloading of the 10 to 20% PCNA that freely slid off the DNA during the assay period (Fig. 4E). Neither the four-subunit Rfc2-5 core complex nor the checkpoint clamp loader Rad24-RFC was able to unload PCNA. A fourth putative clamp loader, Elg1-RFC, was also inactive in unloading PCNA. In addition, in an experiment identical to that shown in Fig. 3A, loading of PCNA by Elg1-RFC was also not observed (data not shown). This complex also failed to load the DNA damage checkpoint clamp Rad17/Mec3/Dcc1 onto primed DNA, under conditions that gave efficient loading by Rad24-RFC, leaving it currently as the only clamp loader with no specific biochemical function (data not shown) (29).

In order to test the DNA substrate preference for PCNA unloading by Ctf18-RFC, we used an assay in which radiolabeled PCNA encircling DNA was quantitated by glutaraldehyde cross-linking, followed by agarose gel electrophoresis and phosphorimager analysis (Fig. 5A). This assay was carried out at 20°C, because initial assays showed that unloading proceeded too rapidly at 30°C to be measured reliably by this assay. Control experiments showed that addition of glutaraldehyde promptly inactivated the loader, with the reaction being complete within 10 s. First, we assessed unloading from circular double-stranded DNA containing a 90-nt ssDNA gap. PCNA was loaded onto the gapped DNA by RFC, and the DNA-bound PCNA was purified by gel filtration and used in a subsequent unloading reaction. Ctf18-RFC unloaded PCNA from the gapped DNA, and this unloading required a hydrolyzable form of ATP, since ATP γ S-mediated unloading was inefficient (Fig. 5B, lane 5 versus lane 6).

The unloaded PCNA was released in a complex with Ctf18-RFC; this complex was efficiently cross-linked by glutaraldehyde and migrated at a unique position in the agarose gel.




FIG. 5. Rates of PCNA unloading by Ctf18-RFC. (A) Scheme of the assay. Multiple ³²P-PCNA rings were loaded onto SKII⁺ DNA containing a 90-nt ssDNA gap. (B) ³²P-PCNA was loaded onto gapped DNA by RFC and isolated by BioGel-A5m filtration as described in Materials and Methods. Isolated complex containing 120 fmol ³²P-PCNA was incubated for 5 min at 20°C with 250 fmol of Ctf18-RFC where indicated and either without nucleotide or with 1 mM ATP or 100 μM ATP_γS. After glutaraldehyde cross-linking and gel electrophoresis, the gel was dried and subjected to phosphorimager analysis. Migration positions of glutaraldehyde-cross-linked PCNA and Ctf18-RFC-PCNA were determined independently. (C) Unloading reactions were identical to those analyzed in (B), except that increasing amounts of Ctf18-RFC were used, and aliquots were removed at the indicated times and cross-linked prior to electrophoresis. The DNA-PCNA band shown in (B) was quantitated.

Unloading by Ctf18-RFC was rapid, being complete after 1 min (Fig. 5C). Unloading also appeared to be stoichiometric rather than catalytic; an excess of loader over PCNA was required to completely unload the clamp (Fig. 5C). Although one explanation of these results could be that the act of PCNA unloading inactivates Ctf18-RFC, we think that based upon

inhibition data presented below, lack of turnover is caused by the extremely slow PCNA product dissociation.

Surprisingly, under these same reaction conditions, yeast RFC failed to significantly unload PCNA even though several groups have reported that human RFC can unload PCNA in an ATP-dependent reaction (6, 40, 50). Since most of the unloading assays reported by these groups were carried out with nicked plasmid DNA, we repeated our unloading studies with nicked DNA plasmids. During prolonged incubation with substoichiometric concentrations of RFC, multiple PCNA trimers are loaded onto nicked plasmids because rapid turnover of RFC occurs. Consequently, the PCNA-DNA complexes isolated by gel filtration are virtually free of RFC (data not shown). When this isolated complex was incubated with ATP and an excess of Ctf18-RFC, rapid unloading was detected (Fig. 6A). However, when the complex was incubated with ATP and RFC, we merely observed a reloading of the small fraction of PCNA that had spontaneously dissociated from the DNA during the complex isolation procedure. Mere incubation of the isolated complex with or without ATP resulted in a very slow spontaneous dissociation of PCNA with an estimated half-life of 40 to 60 min at 20°C (Fig. 6A and data not shown, respectively). Similar unloading assays with RFC were also carried out at very low DNA-PCNA concentrations, since unloading might be favored under these conditions (see Discussion). Even at the lowest concentration practical (0.3 nM DNA-PCNA), RFC still was more proficient in reloading spontaneously dissociated PCNA than in active unloading (Fig. 6B, experiment 2). Unloading by Ctf18-RFC was still proficient at these low concentrations (experiment 1).

The rates of PCNA unloading from different DNA substrates by Ctf18-RFC were determined. At 3 nM DNA-PCNA, unloading from a DNA nick was rapid, with a half-time of unloading of ~20 s at 20°C (Fig. 6A). In contrast, unloading from RPA-coated primed ssDNA was much slower, with a half-time of ~3 min (Fig. 6C, experiment 5). The apparent stoichiometric nature of PCNA unloading by Ctf18-RFC, as suggested by the data in Fig. 5C, indicates that Ctf18-RFC forms an unusually stable complex with PCNA. Possibly, the




FIG. 6. Unloading of PCNA by RFC and Ctf18-RFC. (A) ³²P-PCNA was loaded onto nicked DNA by RFC and isolated by BioGel-A5m filtration as described for Fig. 5A and in Materials and Methods. Isolated complex containing 3 nM ³²P-PCNA was incubated at 20°C with 1 mM ATP and either no complex, 10 nM of Ctf18-RFC, or 10 nM of wild-type RFC. Aliquots were taken after 15 s, 60 s, 3 min, and 10 min, cross-linked with glutaraldehyde, and further processed for analysis. (B) unloading of PCNA from nicked DNA. Assays were similar to those in (A), and the same time points were used. Experiment 1, 0.3 nM DNA-PCNA and 0.5 nM Ctf18-RFC; experiment 2, 0.3 nM DNA-PCNA and 0.5 nM wild-type RFC; experiment 3, 3 nM DNA-PCNA and 10 nM RFC-1ΔN; experiment 4, 3 nM DNA-PCNA and 10 nM Ctf18-RFC previously incubated with a twofold molar excess of PCNA for 30 min at 0°C. (C) Unloading of PCNA from RPA-coated deca-primed SS DNA (see Fig. 3A). Assays were carried out as in (B): experiment 5, 3 nM DNA-PCNA and 10 nM Ctf18-RFC; experiment 6, 3 nM DNA-PCNA and 10 nM Ctf18-RFC previously incubated with a twofold molar excess of PCNA for 30 min at 0°C.

nature of the unloading reaction itself proceeded to yield PCNA–Ctf18-RFC as a very stable complex, thereby effectively inactivating Ctf18-RFC. This complex was detected as an uniquely migrating band by agarose gel electrophoresis (Fig. 5B and 6A). Alternatively, this type of stable complex could also be formed simply by mixing Ctf18-RFC with PCNA without nucleotide. Indeed, when Ctf18-RFC was preincubated with a twofold molar excess of PCNA prior to its addition to an unloading assay, the observed rates of unloading were substantially decreased. Half-times of unloading increased from 20 s to ~10 min for nicked DNA and from 3 min to >30 min for RPA-coated primed DNA (Fig. 6B, experiment 4, and Fig. 6C, experiment 6, respectively). Inhibition was not alleviated by increasing the temperature of the assay, from 20°C to 30°C, or by increasing salt to 100 mM NaCl (data not shown).

DISCUSSION

Comparison between yeast and human clamp loaders. During our investigation of the yeast Ctf18-RFC complex, two biochemical studies of the human complex were published that require a more in-depth comparative discussion (2, 41). In addition, earlier work had indicated that human RFC itself may function as an unloader of PCNA (6, 40, 50). To put those data in the context of the present study, we will first address the clamp unloading activities of RFC and then the activities of Ctf18-RFC.

The kinetic mechanism of PCNA loading by yeast RFC has been studied in detail and shown to follow a multistep ordered pathway with each step fueled by binding of ATP (17, 18). Initially, RFC forms a complex with PCNA; this PCNA-RFC complex then binds template-primer DNA. Hydrolysis of the bound ATP mediates release of RFC from the DNA-PCNA complex. All of these steps are proposed to take place on a msec time scale. However, the observation that RFC exhibits a high level of steady-state ATPase activity in the presence of PCNA and template-primer DNA is an indication that RFC can interact with the DNA-PCNA complex in one or more reactions that turn over ATP, with unloading of PCNA likely being one of them. Assuming that three ATPs are consumed during loading and three during unloading, yeast RFC would cycle through a loading/unloading cycle once every 3 to 4 s (Table 2) (39). If this is the case, why then do we consistently fail to observe unloading of PCNA by yeast RFC while unloading by human RFC can be readily demonstrated?

Both loading and unloading of PCNA catalyzed by RFC and ATP can be represented by the following two irreversible reactions:



If ATP is present at saturation, the rate of loading is a function of the concentrations of three variables, PCNA, DNA, and RFC, while unloading rates depend on the concentrations of only two variables: the DNA-PCNA complex and RFC. An equilibrium is reached when loading and unloading rates are equal. However, this is not an equilibrium in the sense of a classical enzymatic reaction, because each independent irreversible direction is driven by ATP hydrolysis. Therefore, the

equilibrium may be different for each species of RFC, depending on how competent this RFC is in loading and unloading, respectively. At equilibrium, for any given RFC species, the fraction of DNA-PCNA complex over total PCNA should largely be a function of the PCNA, the DNA, and the DNA-PCNA concentrations (equations 1 and 2).

This equilibrium should shift towards unloading at very low DNA-PCNA concentrations, because the reverse reaction, i.e., reloading, should slow down substantially when the concentrations of the substrates, PCNA and DNA, are below their K_d values for binding. These K_d values are 1.3 nM and 5 nM, respectively, for RFC binding to ATP and PCNA-RFC binding to DNA (17). Yet even at a starting concentration of 0.3 nM DNA-PCNA in the unloading assay, reloading by RFC of free PCNA that was either generated by spontaneous dissociation or by unloading was the predominant reaction (Fig. 6B, experiment 2). We also did not obtain evidence for PCNA unloading by RFC when the DNA-PCNA substrate was gapped DNA or primed ssDNA (Fig. 4E and data not shown). We have carried out these studies with wild-type RFC and with a truncated form of RFC, RFC-1ΔN, which removes an N-terminal 272-amino-acid domain of Rfc1 that is somewhat inhibitory to PCNA loading (Fig. 1) (16, 38, 49). Both forms of RFC exhibited the same behavior. Therefore, while the large stimulation of the steady-state ATPase activity of RFC by PCNA and effector DNA suggests that RFC catalyzes cycles of PCNA loading and unloading, our failure to measure significant unloading per se indicates that the efficiency of loading far outweighs the efficiency of unloading.

PCNA loading and unloading by Ctf18-RFC. While our results strongly indicate that under physiological conditions Ctf18-RFC is an unloader rather than a loader of PCNA, the studies of the human Ctf18-RFC complex are in part at odds with our conclusions. Both Hurwitz and colleagues and Tsurimoto and colleagues have studied the PCNA loading properties of the human five-subunit enzyme, and the Hurwitz group also studied the seven-subunit enzyme (2, 41). Similar to our studies, no fundamental biochemical differences were detected between the five-subunit and the seven-subunit human enzymes. However, both groups detected significant loading of PCNA onto primed DNA in a reaction which was not appreciably inhibited by addition of human RPA. In our system, subsaturating RPA actually stimulated PCNA loading, and it was only upon complete coating of the ssDNA with RPA that the strong and specific inhibition by RPA could be demonstrated (Fig. 3B).

However, in actuality our data on the yeast cohesion clamp loader may not be as much in disagreement with those on the human enzyme as they initially appear to be. It is evident from our studies that yeast Ctf18-RFC cycles in an equilibrium between PCNA loading and PCNA unloading and that this equilibrium is shifted strongly towards unloading in the presence of RPA. A similar equilibrium may also exist for the human complex; however, this equilibrium may be less severely disturbed by the presence of human RPA. It will be interesting to see whether the human Ctf18-RFC complex also unloads PCNA and how RPA controls these two opposing processes.

Role of RPA in regulating Ctf18-RFC function. Yeast Ctf18-RFC is both a clamp loader and a clamp unloader, and each direction of this PCNA cycle is driven by the energy of ATP




FIG. 7. Possible roles for PCNA loading and unloading in the establishment of sister chromatid cohesion. See the text for details.

hydrolysis. Does Ctf18-RFC go through cycles of PCNA loading and unloading as proposed for RFC? Our studies show that such a cycle cannot operate efficiently because the DNA substrates for PCNA loading and for unloading are quite different (compare equations 1 and 2 for RFC with equations 3 and 4 for Ctf18-RFC).



where DNA_1 is naked template-primer DNA and DNA_2 is nicked DNA or RPA-coated template-primer DNA.

Only template-primer DNA with an adjacent naked ssDNA region or coated with a mutant RPA or heterologous ssDNA binding protein is a proficient substrate for PCNA loading by Ctf18-RFC in vitro. Generally, it is assumed that any ssDNA in the cell is bound up with RPA, which should inhibit the loading activity of Ctf18-RFC. Since Ctf18-RFC is also unable to load PCNA at a nick (Table 1), it appears that so far there is no demonstrated substrate for PCNA loading. On the other hand, Ctf18-RFC efficiently unloads PCNA from nicked and gapped substrates and somewhat less efficiently from RPA-coated primed ssDNA (Fig. 5 and 6).

Surprisingly, however, we found that the inhibition by RPA is not a mere consequence of coating the ssDNA. Whereas coating by *E. coli* SSB and phage T4 gp32 protein caused only a minor inhibition of PCNA loading, coating by yeast RPA-1 Δ N actually stimulated PCNA loading by Ctf18-RFC (108 to 150% in three determinations) (Table 1). The N-terminal domain of Rpa1, lacking in RPA-1 Δ N, while not required for ssDNA binding, has been implicated in protein-protein interactions with the Rfc4 subunit of Ctf18-RFC (23). Mutations in the *RF1* gene that map to its N terminus show synthetic lethality with an *rfc4-2* mutation. Singly, these mutants show both replication defects and DNA damage checkpoint defects; the possible defects in chromatid cohesion were not investigated (23). Currently, it is not clear how the Rpa1-Rfc4 interaction could function to inhibit Ctf18-RFC while stimulating RFC, and additional specific interactions may exist. Another relevant question is whether yeast has a pathway that relieves this inhibition by a modification of Rpa1 and/or Ctf18-RFC? So far no in vivo modification of the Ctf18 subunit has been

reported. Although Rpa1 does not appear to be phosphorylated during normal DNA replication in the absence of stress or DNA damage, phosphorylation of the Rpa2 subunit persists throughout the S phase of the cell cycle (8).

A proposed role for PCNA unloading in the establishment of sister chromatid cohesion. Previous proposals for the function of Ctf18-RFC have focused on its PCNA-loading function and the potential of this loader to target establishment-specific protein(s) to the newly loaded PCNA onto replicating DNA (2, 9, 30). Specific localization to replicating DNA at cohesin binding sites may be imparted by interactions between Ctf18-RFC and cohesin (2). Since PCNA is loaded by Ctf18-RFC in the same orientation and at the same 3'-DNA structure as it is by RFC, localization of these putative proteins cannot just depend on PCNA binding, or else loading by RFC would suffice; further specificity would have to be imparted by interactions with Ctf18-RFC. Currently, it is not known which factor is supposed to be targeted to this newly loaded PCNA. One function of this factor could be to mediate bypass of the cohesin ring past the replication fork (Fig. 7). While it is still controversial whether the association of cohesin with chromosomes is peripheral in nature or whether the cohesin ring actually encircles the chromosome, as shown in Fig. 7, in both models restructuring of the fork and/or the action of specific factors may be required to allow the replication fork to bypass the obstacle formed by the cohesin ring (reviewed in reference 33). However, if cohesin does encircle the replicating chromosome, an alternative model requiring partial fork disassembly is suggested by the necessity for the DNA replication apparatus to thread through the cohesin ring.

Cohesin is loaded around the chromatin as a large 30- to 40-nm-diameter ring during late G_1 . Subsequently, sister chromatid cohesion would be established simply by passage of the replication fork through the cohesin ring (reviewed in references 34 and 48) (Fig. 7). Support for this model is derived from the ability of the cohesin ring to slide along chromatin both before and after DNA replication (26). Moreover, the act of transcription relocates cohesin rings to intergenic regions of convergent transcription, where the transcription machineries collide. A persuasive reasoning for this transcription-dependent cohesin sliding is that the transcription machinery is

simply too large to pass through the cohesin ring without disturbing its position on the chromatin. However, the replication machinery may still be larger in size than that. Polypeptides known to be associated with the fork, the MCM helicase, three DNA polymerases and their associated proteins, the Okazaki fragment maturation machinery, and perhaps also the mismatch repair machinery, already make up a complex 3 to 5 MDa in size. Although the physical size of such a complex if globular would only be 12 to 15 nm in diameter and should therefore be able to fit through the cohesin ring, there are other considerations that greatly increase the physical size of the replisome.

In order to achieve coordinate replication of the leading and lagging strands, the lagging strand is proposed to fold back onto itself, forming an extruded loop, the trombone model (42) (Fig. 7). So far, strong experimental support for this model derives only from electron-microscopic studies of the bacteriophage T7 and T4 replisomes (7, 24). The core of the small T4 replisome already has an estimated size of ~20 nm, without even taking the looped-out lagging-strand DNA into consideration. Assuming that cellular replisomes also adopt a trombone model, it is evident that the much larger eukaryotic replisome with its associated ssDNA loop coated with RPA would be very unlikely to fit through the 30- to 40-nm-diameter hole of cohesin. However, if the trombone structure could be relaxed and a linear arrangement of the replication components achieved, the replication machinery could easily pass through the cohesin ring. Since PCNA rings situated on both the leading and the lagging strand are central structural organizing elements of the replisome, unloading of PCNA by Ctf18-RFC from only one of the strands would be expected to cause a collapse of the trombone superstructure, without necessarily displacing the DNA polymerases. Continued replication would mediate passage of the fork through the cohesin ring. After replicational passage through the cohesin ring is completed, reloading of PCNA would reestablish the superstructure of the replisome. This reloading could either be accomplished by the Ctf18-RFC complex that still has the PCNA bound to it, at novel DNA structures, or at normal template-primer locations free of RPA or, alternatively, by normal RFC.

CTF18 deletion mutants and similarly *CTF8* and *DCC1* mutants show synthetic lethality not only with other mutants in cohesion and chromosome segregation but also with mutations in the lagging-strand DNA replication apparatus (45). Cohesion establishment gene *CTF7* shows synthetic lethality with *POL30* (PCNA) mutations, and overexpression of PCNA suppresses the temperature sensitivity of a *ctf7* mutant (44). Ctf7 also interacts with the Ctf18-RFC complex (21). Since interaction was also detected with RFC and with the DNA damage checkpoint Rad24-RFC complex, it is likely that this interaction with Ctf7 is mediated through the small subunits of RFC common to all these complexes. Ctf7 is a lysine acetylase, although the importance of the acetylation of Ctf7 for establishing sister chromatid cohesion has been called into question (3, 20). Therefore, it remains unclear how the demonstrated genetic link between Ctf7 and PCNA can be put into a mechanistic framework to further clarify the role of Ctf18-RFC in the establishment of chromatid cohesion through loading/unloading of PCNA.

It should be stressed that besides the Ctf18-RFC complex,

establishment of sister chromatid cohesion depends on several other proteins whose functions can be envisaged at the replication fork. Among those, Chl1 is a putative DNA helicase and Ctf4 is a Pol α -interacting protein (12, 19, 31, 36, 43). Deletions in each of these genes show phenotypes very analogous to those in the *CTF18*, *DCC1*, and *CTF8* subunit genes of the Ctf18-RFC clamp loader: failure to establish efficient sister chromatid cohesion in haploids and severe defects in chromosome disjunction during meiosis, with resulting poor spore viability in homozygous diploid mutants. Of these various establishment-associated proteins, only Chl1 has a putative PCNA-binding motif (NNHLQOFF, 643 to 650 [conserved amino acids are in boldface]), although interaction of an establishment factor with PCNA through a nonconsensus interaction domain also remains a possibility (reviewed in reference 13). How each of these proteins, perhaps in a complex, could mediate the passage of the replication fork past or through the cohesin ring remains to be determined.

ACKNOWLEDGMENTS

We thank John Majors and members of the Burgers lab for critical discussions and Jurek Majka for gifts of Rad24-RFC and RPA-1N. The truncation RPA plasmids were a generous gift from Steven Brill.

This research was supported in part by National Institutes of Health Grant GM32431. G.O.B. received partial support from the Washington University-Umeå University exchange program.

REFERENCES

- Bae, K. H., H. S. Kim, S. H. Bae, H. Y. Kang, S. Brill, and Y. S. Seo. 2003. Bimodal interaction between replication-protein A and Dna2 is critical for Dna2 function both *in vivo* and *in vitro*. *Nucleic Acids Res.* **31**:3006–3015.
- Bermudez, V. P., Y. Maniwa, I. Tappin, K. Ozato, K. Yokomori, and J. Hurwitz. 2003. The alternative Ctf18-Dcc1-Ctf8-replication factor C complex required for sister chromatid cohesion loads proliferating cell nuclear antigen onto DNA. *Proc. Natl. Acad. Sci. USA* **100**:10237–10242.
- Brands, A., and R. V. Skibbens. 2005. Ctf7p/Eco1p exhibits acetyltransferase activity—but does it matter? *Curr. Biol.* **15**:R50–R51.
- Burgers, P. M. 1999. Overexpression of multisubunit replication factors in yeast. *Methods* **18**:349–355.
- Burgers, P. M., and K. J. Gerik. 1998. Structure and processivity of two forms of *Saccharomyces cerevisiae* DNA polymerase delta. *J. Biol. Chem.* **273**:19756–19762.
- Cai, J., F. Uhlmann, E. Gibbs, R. H. Flores, C. G. Lee, B. Phillips, J. Finkelstein, N. Yao, M. O'Donnell, and J. Hurwitz. 1996. Reconstitution of human replication factor C from its five subunits in baculovirus-infected insect cells. *Proc. Natl. Acad. Sci. USA* **93**:12896–12901.
- Chastain, P. D., II, A. M. Makhov, N. G. Nossal, and J. Griffith. 2003. Architecture of the replication complex and DNA loops at the fork generated by the bacteriophage T₄ proteins. *J. Biol. Chem.* **278**:21276–21285.
- Din, S., S. J. Brill, M. P. Fairman, and B. Stillman. 1990. Cell-cycle-regulated phosphorylation of DNA replication factor A from human and yeast cells. *Genes Dev.* **4**:968–977.
- Edwards, S., C. M. Li, D. L. Levy, J. Brown, P. M. Snow, and J. L. Campbell. 2003. *Saccharomyces cerevisiae* DNA polymerase epsilon and polymerase sigma interact physically and functionally, suggesting a role for polymerase epsilon in sister chromatid cohesion. *Mol. Cell. Biol.* **23**:2733–2748.
- Eisenberg, J. C., R. Ayyagari, X. V. Gomes, and P. Burgers. 1997. Mutations in yeast proliferating cell nuclear antigen define distinct sites for interaction with DNA polymerase delta and DNA polymerase epsilon. *Mol. Cell. Biol.* **17**:6367–6378.
- Fien, K., and B. Stillman. 1992. Identification of replication factor C from *Saccharomyces cerevisiae*: a component of the leading-strand DNA replication complex. *Mol. Cell. Biol.* **12**:155–163.
- Formosa, T., and T. Nittis. 1999. Dna2 mutants reveal interactions with Dna polymerase alpha and Ctf4, a Pol alpha accessory factor, and show that full Dna2 helicase activity is not essential for growth. *Genetics* **151**:1459–1470.
- Garg, P., and P. M. Burgers. 2005. DNA polymerases that propagate the eukaryotic DNA replication fork. *Crit. Rev. Biochem. Mol. Biol.* **40**:115–128.
- Gerik, K. J., S. L. Gary, and P. M. Burgers. 1997. Overproduction and affinity purification of *Saccharomyces cerevisiae* replication factor C. *J. Biol. Chem.* **272**:1256–1262.
- Gomes, X. V., and P. M. J. Burgers. 2000. Two modes of FEN1 binding to PCNA regulated by DNA. *EMBO J.* **19**:3811–3821.

16. **Gomes, X. V., S. L. Gary, and P. M. Burgers.** 2000. Overproduction in *Escherichia coli* and characterization of yeast replication factor C lacking the ligase homology domain. *J. Biol. Chem.* **275**:14541–14549.
17. **Gomes, X. V., and P. M. Burgers.** 2001. ATP utilization by yeast replication factor C. I. ATP-mediated interaction with DNA and with proliferating cell nuclear antigen. *J. Biol. Chem.* **276**:34768–34775.
18. **Gomes, X. V., S. L. Schmidt, and P. M. Burgers.** 2001. ATP utilization by yeast replication factor C. II. Multiple stepwise ATP binding events are required to load proliferating cell nuclear antigen onto primed DNA. *J. Biol. Chem.* **276**:34776–34783.
19. **Hanna, J. S., E. S. Kroll, V. Lundblad, and F. A. Spencer.** 2001. *Saccharomyces cerevisiae* CTF18 and CTF4 are required for sister chromatid cohesion. *Mol. Cell Biol.* **21**:3144–3158.
20. **Ivanov, D., A. Schleiffer, F. Eisenhaber, K. Mechtler, C. H. Haering, and K. Nasmyth.** 2002. Eco1 is a novel acetyltransferase that can acetylate proteins involved in cohesion. *Curr. Biol.* **12**:323–328.
21. **Kenna, M. A., and R. V. Skibbens.** 2003. Mechanical link between cohesion establishment and DNA replication: Ctf7p/Eco1p, a cohesion establishment factor, associates with three different replication factor C complexes. *Mol. Cell Biol.* **23**:2999–3007.
22. **Kim, C., R. O. Snyder, and M. S. Wold.** 1992. Binding properties of replication protein A from human and yeast cells. *Mol. Cell Biol.* **12**:3050–3059.
23. **Kim, H. S., and S. J. Brill.** 2001. Rfc4 interacts with Rpa1 and is required for both DNA replication and DNA damage checkpoints in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* **21**:3725–3737.
24. **Lee, J., P. N. Chastain, T. Kusakabe, J. D. Griffith, and C. C. Richardson.** 1998. Coordinated leading and lagging strand DNA synthesis on a minicircular template. *Mol. Cell* **1**:1001–1010.
25. **Lee, S. H., and J. Hurwitz.** 1990. Mechanism of elongation of primed DNA by DNA polymerase delta, proliferating cell nuclear antigen, and activator 1. *Proc. Natl. Acad. Sci. USA* **87**:5672–5676.
26. **Lengronne, A., Y. Katou, S. Mori, S. Yokobayashi, G. P. Kelly, T. Itoh, Y. Watanabe, K. Shirahige, and F. Uhlmann.** 2004. Cohesin relocation from sites of chromosomal loading to places of convergent transcription. *Nature* **430**:573–578.
27. **Majka, J., and P. M. Burgers.** 2004. The PCNA-RFC families of DNA clamps and clamp loaders. *Prog. Nucleic Acids Res. Mol. Biol.* **78**:227–260.
28. **Majka, J., and P. M. Burgers.** 2003. Yeast Rad17/Mec3/Ddc1: a sliding clamp for the DNA damage checkpoint. *Proc. Natl. Acad. Sci. USA* **100**:2249–2254.
29. **Majka, J., B. Y. Chung, and P. M. Burgers.** 2004. Requirement for ATP by the DNA damage checkpoint clamp loader. *J. Biol. Chem.* **279**:20921–20926.
30. **Mayer, M. L., S. P. Gygi, R. Aebersold, and P. Hieter.** 2001. Identification of RFC(Ctf18p, Ctf8p, Dcc1p): an alternative RFC complex required for sister chromatid cohesion in *S. cerevisiae*. *Mol. Cell* **7**:959–970.
31. **Mayer, M. L., I. Pot, M. Chang, H. Xu, V. Aneliunas, T. Kwok, R. Newitt, R. Aebersold, C. Boone, G. W. Brown, and P. Hieter.** 2004. Identification of protein complexes required for efficient sister chromatid cohesion. *Mol. Biol. Cell* **15**:1736–1745.
32. **Merkle, C. J., L. M. Karnitz, J. T. Henry-Sánchez, and J. Chen.** 2003. Cloning and characterization of hCTF18, hCTF8, and hDCC1. *J. Biol. Chem.* **278**:30051–30056.
33. **Milutinovich, M., and D. E. Koshland.** 2003. Molecular biology. SMC complexes—wrapped up in controversy. *Science* **300**:1101–1102.
34. **Nasmyth, K., and A. Schleiffer.** 2004. From a single double helix to paired double helices and back. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **359**:99–108.
35. **Parsons, A. B., R. L. Brost, H. Ding, Z. Li, C. Zhang, B. Sheikh, G. W. Brown, P. M. Kane, T. R. Hughes, and C. Boone.** 2004. Integration of chemical-genetic and genetic interaction data links bioactive compounds to cellular target pathways. *Nat. Biotechnol.* **22**:62–69.
36. **Petronczki, M., B. Chwalla, M. F. Siomos, S. Yokobayashi, W. Helmhart, A. M. Deutschbauer, R. W. Davis, Y. Watanabe, and K. Nasmyth.** 2004. Sister-chromatid cohesion mediated by the alternative RF-CCtf18/Dcc1/Ctf8, the helicase Chl1 and the polymerase-alpha-associated protein Ctf4 is essential for chromatid disjunction during meiosis II. *J. Cell Sci.* **117**:3547–3559.
37. **Podust, L. M., V. N. Podust, J. M. Sogo, and U. Hubscher.** 1995. Mammalian DNA polymerase auxiliary proteins: analysis of replication factor C-catalyzed proliferating cell nuclear antigen loading onto circular double-stranded DNA. *Mol. Cell Biol.* **15**:3072–3081.
38. **Podust, V. N., N. Tiwari, S. Stephan, and E. Fanning.** 1998. Replication factor C disengages from proliferating cell nuclear antigen (PCNA) upon sliding clamp formation, and PCNA itself tethers DNA polymerase delta to DNA. *J. Biol. Chem.* **273**:31992–31999.
39. **Schmidt, S. L., X. V. Gomes, and P. M. Burgers.** 2001. ATP utilization by yeast replication factor C. III. The ATP-binding domains of Rfc2, Rfc3, and Rfc4 are essential for DNA recognition and clamp loading. *J. Biol. Chem.* **276**:34784–34791.
40. **Shibahara, K., and B. Stillman.** 1999. Replication-dependent marking of DNA by PCNA facilitates CAF-1-coupled inheritance of chromatin. *Cell* **96**:575–585.
41. **Shiomi, Y., A. Shinozaki, K. Sugimoto, J. Usukura, C. Obuse, and T. Tsurimoto.** 2004. The reconstituted human Chl12-RFC complex functions as a second PCNA loader. *Genes Cells* **9**:279–290.
42. **Sinha, N. K., C. F. Morris, and B. M. Alberts.** 1980. Efficient *in vitro* replication of double-stranded DNA templates by a purified T4 bacteriophage replication system. *J. Biol. Chem.* **255**:4290–4293.
43. **Skibbens, R. V.** 2004. Chl1p, a DNA helicase-like protein in budding yeast, functions in sister-chromatid cohesion. *Genetics* **166**:33–42.
44. **Skibbens, R. V., L. B. Corson, D. Koshland, and P. Hieter.** 1999. Ctf7p is essential for sister chromatid cohesion and links mitotic chromosome structure to the DNA replication machinery. *Genes Dev.* **13**:307–319.
45. **Tong, A. H., G. Lesage, G. D. Bader, H. Ding, H. Xu, X. Xin, J. Young, G. F. Berriz, R. L. Brost, M. Chang, Y. Chen, X. Cheng, G. Chua, H. Friesen, D. S. Goldberg, J. Haynes, C. Humphries, G. He, S. Hussein, L. Ke, N. Krogan, Z. Li, J. N. Levinson, H. Lu, P. Menard, C. Munyana, A. B. Parsons, O. Ryan, R. Tonikian, T. Roberts, A. M. Sdicu, J. Shapiro, B. Sheikh, B. Suter, S. L. Wong, L. V. Zhang, H. Zhu, C. G. Burd, S. Munro, C. Sander, J. Rine, J. Greenblatt, M. Peter, A. Bretscher, G. Bell, F. P. Roth, G. W. Brown, B. Andrews, H. Bussey, and C. Boone.** 2004. Global mapping of the yeast genetic interaction network. *Science* **303**:808–813.
46. **Toth, A., R. Ciosk, F. Uhlmann, M. Galova, A. Schleiffer, and K. Nasmyth.** 1999. Yeast cohesin complex requires a conserved protein, Eco1p(Ctf7), to establish cohesion between sister chromatids during DNA replication. *Genes Dev.* **13**:320–333.
47. **Tsurimoto, T., and B. Stillman.** 1991. Replication factors required for SV40 DNA replication *in vitro*. I. DNA structure-specific recognition of a primer-template junction by eukaryotic DNA polymerases and their accessory proteins. *J. Biol. Chem.* **266**:1950–1960.
48. **Uhlmann, F.** 2004. The mechanism of sister chromatid cohesion. *Exp. Cell Res.* **296**:80–85.
49. **Uhlmann, F., J. Cai, E. Gibbs, M. O'Donnell, and J. Hurwitz.** 1997. Deletion analysis of the large subunit p140 in human replication factor C reveals regions required for complex formation and replication activities. *J. Biol. Chem.* **272**:10058–10064.
50. **Yao, N., J. Turner, Z. Kelman, P. T. Stukenberg, F. Dean, D. Shechter, Z. Q. Pan, J. Hurwitz, and M. O'Donnell.** 1996. Clamp loading, unloading and intrinsic stability of the PCNA, beta and gp45 sliding clamps of human, *E. coli* and T4 replicases. *Genes Cells* **1**:101–113.
51. **Yoder, B. L., and P. M. J. Burgers.** 1991. *Saccharomyces cerevisiae* replication factor C. I. Purification and characterization of its ATPase activity. *J. Biol. Chem.* **266**:22689–22697.