

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2010

Protein misfolding induces hypoxic preconditioning via a subset of the unfolded protein response machinery

Xianrong R. Mao

Washington University School of Medicine in St. Louis

C. Michael Crowder

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Mao, Xianrong R. and Crowder, C. Michael, "Protein misfolding induces hypoxic preconditioning via a subset of the unfolded protein response machinery." *Molecular and Cellular Biology*. 30, 21. 5033-5042. (2010).

https://digitalcommons.wustl.edu/open_access_pubs/2275

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Protein Misfolding Induces Hypoxic Preconditioning via a Subset of the Unfolded Protein Response Machinery

Xianrong R. Mao and C. Michael Crowder
Mol. Cell. Biol. 2010, 30(21):5033. DOI:
10.1128/MCB.00922-10.
Published Ahead of Print 23 August 2010.

Updated information and services can be found at:
<http://mcb.asm.org/content/30/21/5033>

These include:

REFERENCES

This article cites 60 articles, 24 of which can be accessed free at: <http://mcb.asm.org/content/30/21/5033#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Protein Misfolding Induces Hypoxic Preconditioning via a Subset of the Unfolded Protein Response Machinery[▽]

Xianrong R. Mao¹ and C. Michael Crowder^{1,2*}

Departments of Anesthesiology¹ and Developmental Biology,² Washington University School of Medicine,
660 South Euclid Avenue, Saint Louis, Missouri 63110

Received 9 August 2010/Accepted 10 August 2010

Prolonged cellular hypoxia results in energy failure and ultimately cell death. However, less-severe hypoxia can induce a cytoprotective response termed hypoxic preconditioning (HP). The unfolded protein response pathway (UPR) has been known for some time to respond to hypoxia and regulate hypoxic sensitivity; however, the role of the UPR, if any, in HP essentially has been unexplored. We have shown previously that a sublethal hypoxic exposure of the nematode *Caenorhabditis elegans* induces a protein chaperone component of the UPR (L. L. Anderson, X. Mao, B. A. Scott, and C. M. Crowder, *Science* 323:630–633, 2009). Here, we show that HP induces the UPR and that the pharmacological induction of misfolded proteins is itself sufficient to stimulate a delayed protective response to hypoxic injury that requires the UPR pathway proteins IRE-1, XBP-1, and ATF-6. HP also required IRE-1 but not XBP-1 or ATF-6; instead, GCN-2, which is known to suppress translation and induce an adaptive transcriptional response under conditions of UPR activation or amino acid deprivation, was required for HP. The phosphorylation of the translation factor eIF2 α , an established mechanism of GCN-2-mediated translational suppression, was not necessary for HP. These data suggest a model where hypoxia-induced misfolded proteins trigger the activation of IRE-1, which along with GCN-2 controls an adaptive response that is essential to HP.

A lack of oxygen supply (hypoxia) poses serious challenges for cells that must then adapt to low oxygen until conditions improve or die. However, the precise cascade of events that control whether a cell adapts or dies in the face of hypoxia is unclear. Adaptive hypoxic protective mechanisms can be induced by brief sublethal exposures to hypoxia and/or ischemia. This phenomenon is called hypoxic or ischemic preconditioning and has been the subject of intense study to define intrinsic hypoxia protective mechanisms (6, 12, 30, 34). Two forms of hypoxic preconditioning (HP) have been described (12, 51). Immediate preconditioning appears within minutes after the sublethal hypoxic/ischemic episode and wanes within about 4 h; delayed preconditioning appears 12 to 24 h later and can last for days. Delayed preconditioning is thought to require changes in gene expression through new transcription (12).

The nematode *Caenorhabditis elegans* has been found to have delayed HP (10). A sublethal exposure of *C. elegans* to hypoxia induces the hypoxic protection of the animal as a whole and its myocytes and neurons with an onset of approximately 16 h and a duration of at least 36 h. As for delayed preconditioning in mammals, the mechanism in *C. elegans* for sensing hypoxia, transducing the signal, and inducing cytoprotection is unclear. We recently reported that a sublethal hypoxic exposure similar to that which produces HP induces a reporter of the unfolded protein response (UPR) in *C. elegans* (1). The *C. elegans* UPR consists of three defined branches: IRE-1–XBP-1, ATF-6, and PERK-1 (49). Misfolded proteins are sensed by IRE-1, resulting in homo-oligomerization, autophos-

phorylation, and activation. Activated IRE-1 cleaves XBP-1 mRNA with subsequent splicing to produce a new open reading frame that can be translated into the XBP-1 transcription factor. Similarly activated ATF-6 translocates to the Golgi apparatus in response to elevated misfolded proteins, where it is cleaved by proteases, producing a transcriptionally active form of ATF-6. Both XBP-1 and ATF-6 control the transcription of a large number of genes whose functions are crucial for maintaining endoplasmic reticulum (ER) homeostasis. PERK-1 (PERK-1 in mammals) acts more directly to phosphorylate translation initiation factor eIF2 α and thereby suppress general protein translation, thus reducing the nascent unfolded protein load presented to the ER (44).

As mentioned above, we have shown previously that a UPR reporter is induced by hypoxia in *C. elegans* (1). We also found that a reduction-of-function mutation in *ire-1* can decrease or increase the hypoxic survival of *C. elegans* depending on the presence or absence, respectively, of a second mutation that reduces global translation rates. Others have reported the activation of the UPR after ischemia (2, 11, 17, 18, 39, 43, 54, 57). Hypoxia also has been shown to activate the PERK-1 pathway in cell culture (4, 25). These findings together suggest that the misfolded proteins generated by hypoxia can trigger hypoxic preconditioning by the activation of the UPR that then can protect cells, perhaps by the activation of PERK-1 and translational suppression. Here, we test the various aspects of this hypothesis using *C. elegans* genetic tools.

MATERIALS AND METHODS

Strains. *C. elegans* strains were obtained from the *Caenorhabditis* Genetics Center (CGC), except where noted, and outcrossed three times prior to testing. Mutations were confirmed after outcrossing by sequencing. All strains were maintained at 20°C on NGM agar seeded with OP50 bacteria as described previously (7, 52). The strains carrying *ire-1(zc14)* and *xbp-1(zc12)* were ob-

* Corresponding author. Mailing address: Department of Anesthesiology, Washington University School of Medicine, 660 South Euclid Ave., Box 8054, St. Louis, MO 63110. Phone (314) 362-8560. Fax: (314) 362-8571. E-mail: crowderm@morpheus.wustl.edu.

[▽] Published ahead of print on 23 August 2010.

tained through outcrossing SJ30 *ire-1(zc14);hsp-4::GFP* and SJ17 *xbp-1(zc12);hsp-4::GFP* with the N2 wild type to get rid of the green fluorescent protein (GFP) transgene. The balancer *mIn1[dpy-10(e128) mIs14(p-my0-2::GFP)]* (14) was used to generate *ire-1* heterozygotes and transheterozygotes as follows. *mIn1/+* males were crossed with *ire-1(zc14)* or *ire-1(v33)* hermaphrodites. GFP-positive progeny were kept and presumed to have the genotype *zc14/mIn1* or *v33/mIn1*. *zc14/mIn1* males were crossed with *ire-1(v33)* hermaphrodites, and non-GFP-expressing F1 hermaphrodites segregating from a cross that produced 50% male progeny were assumed to have the genotype *zc14/v33* and were immediately tested. *ire-1(tm400)* and *atf-6(tm1153)* were from Shohei Mitani (Tokyo Women's Medical College, Tokyo, Japan) and the Japan National Biore-source Project (<http://www.shigen.nig.ac.jp/c.elegans/index.jsp>). *tm400* is a relatively new *ire-1* allele that was supplied as a heterozygote, since homozygotes initially were classified as lethal/sterile. However, after outcrossing a few generations, we identified *tm400* homozygotes that were viable and fertile. By using primers encompassing the deletion region of *tm400*, three cDNA products from each primer pair were identified by RT-PCR from *ire-1(tm400)* animals (data not shown). After we sequenced these products, they were identified as novel *ire-1* mRNA species, with a frameshift and early stop codon downstream of the deletion breakpoint. Thus, these mutant mRNAs likely would produce nonfunctional proteins. Given a weaker tunicamycin sensitivity phenotype than that of *ire-1(v33)*, the putative null mutant, we deduce that *tm400* mostly likely produces protein products generated by utilizing downstream in-frame ATGs (most likely Met228, which could produce a 740-amino-acid residue polypeptide with transmembrane, kinase, and ribonuclease domains).

Imaging Phsp-4::GFP. The detailed protocol for imaging Phsp-4::GFP was described previously (1).

Western blotting. After treatment, worms were harvested with M9 buffer. For hypoxic treatments without recovery, worms were harvested inside the hypoxic chamber with deoxygenated M9 buffer (52). The worm pellets were collected, and 1% SDS buffer was added before freezing. After an overnight freeze at -80°C , worm pellets were briefly sonicated and insoluble debris pelleted by centrifugation at $16,000 \times g$ for 15 min. Sample protein concentrations were determined by a NanoDrop spectrophotometer (Thermo Scientific, Wilmington, DE). Forty micrograms of total protein was loaded for each sample for SDS-PAGE. The detailed Western blotting protocol is available elsewhere (www.cellsignaling.com). All antibodies were from Cell Signaling (Beverly, MA): phospho-eIF2 α (p-eIF2 α) (ser51) (no. 9721), β -actin (13E5) (no. 4970), and anti-rat IgG horseradish peroxidase (HRP)-linked secondary antibody and detection system (no. 7077).

RT-PCR. A synchronous population (1 day post-larval stage 4) of animals was treated under conditions described in Results and the figure legends. RNA was isolated by a TRIzol freeze-cracking method. cDNA was synthesized with a RETROscript random decamer kit (Applied Biosystems, Foster City, CA) with 1 μg of total RNA as the template. Quantitative real-time PCR was performed as described previously (1) with SYBR green PCR master mix (Applied Biosystems, Foster City, CA) in an Applied Biosystems 7500-fast reverse transcription-PCR (RT-PCR) instrument with a Rox passive-reference dye. Primers were constructed to amplify an approximately 100-bp fragment. Standard PCR amplification with the primer sets produced single bands migrating at the correct size. Primers for *hsp-4* were AACGGAATATGACGTAAGCGCC (forward) and TGAGACGATTGTGGTTCGTTGGTGA (reverse), and those for *ire-1* were GCTGAACGTGGAGCCATTGCACCG (forward) and CATCCAAGTGAGA AGATATCAACTGG (reverse). Semiquantitative RT-PCR for *xbp-1* was determined with a DNAEngine Peltier thermal cycler (Bio-Rad, Hercules, CA). The exponential phases of amplification for *act-1* and *xbp-1* were empirically determined by two-cycle increments from 14 to 28 cycles. Sixteen and 20 cycles were used for *act-1* and *xbp-1* amplifications, respectively. *xbp-1* primers were GGA GAGGATCGCCGTGCCT (forward) and GATGGAGGTGGATCGGGCC TGTT (reverse). The primers created 81- and 58-bp products for unspliced and spliced forms of *xbp-1*, respectively. The products were resolved with an 8% PAGE gel.

Hypoxia and Tm incubations. Hypoxia treatment and HP protocols were similar to those published previously, except no washes were performed inside the hypoxia chamber, as we have found these to be unnecessary (10, 47). Briefly, synchronized populations of young adult animals were transferred from NGM plates to 1.5-ml Eppendorf tubes with 1 ml M9 buffer; the buffer was removed down to 100 μl , and the tubes then were placed into the hypoxia chamber as described previously (47). For HP incubations, the tubes were removed after 4 h and the worms recovered for 20 h on fresh NGM plates before a hypoxic killing incubation. Normoxic preconditioned controls were treated identically, except the preconditioning incubation instead was performed at 26°C in a normoxic chamber. For hypoxic killing incubations, worms were placed in the hypoxic

chamber as described above and incubated for 20 h. The temperature of the hypoxic incubations was 26°C , and the oxygen concentration was $<0.2\%$. Animals were scored 20 to 24 h after the completion of the hypoxic killing incubations. Animals without pharyngeal pumping and without spontaneous or touch-evoked movement were scored as dead. For tunicamycin (Tm) pretreatments, 1 day post L4 worms were washed off plates and incubated with designated concentrations of Tm in M9 buffer for 4 h at 20°C . After Tm incubation, worms were returned to regular NGM plates with food for 20 h at 20°C before they were challenged with a lethal 20-h hypoxic exposure. Worms then were scored for survival as previously described (10, 47). The Tm growth arrest assay has been published previously (1). Briefly, eggs from animals of various genotypes were laid on NGM plates containing 1 $\mu\text{g}/\text{ml}$ Tm. The fraction of worms reaching the adult stage was scored 3 days later.

RESULTS

Hypoxia induces the UPR. We first asked whether hypoxic preconditioning induces the expression of components of the UPR. We measured two markers of UPR induction, HSP-4 induction and eIF2 α phosphorylation. HSP-4 expression is transcriptionally induced in response to an increase in misfolded proteins by an IRE-1-dependent mechanism (8, 48). After 4 h of hypoxia ($<0.2\% \text{O}_2$) but no reoxygenation, expression from a GFP transgene driven by an *hsp-4* promoter was unchanged from control levels (Fig. 1A and B). However, 4 h of recovery and reoxygenation produced a marked increase in pHSP-4::GFP expression, which then returned to control levels after 20 h of recovery (Fig. 1A and B). To confirm that the transgene expression accurately reflects the native gene activity, we measured HSP-4 native transcript levels by quantitative RT-PCR (qRT-PCR) (Fig. 1C). Indeed, native HSP-4 transcript levels remained unchanged even after 8 h of hypoxia but were significantly increased after reoxygenation, with a return to control levels by 20 h. The translation factor eIF2 α is phosphorylated by the UPR components PERK-1 and GCN-2. Phosphorylated eIF2 α levels increased rapidly during hypoxic incubation but fell back to baseline within 1 h of reoxygenation (Fig. 1D). Thus, while eIF2 α is phosphorylated in response to hypoxia in *C. elegans*, the time course is inconsistent with translational suppression by p-eIF2 α contributing to the effector mechanism of hypoxic cytoprotection by HP.

Induction of the UPR produces a delayed hypoxic protective response. We next asked whether the induction of misfolded proteins is sufficient to induce a hypoxic protective response in a manner similar to that of HP. Tunicamycin (Tm) inhibits the N-glycosylation of proteins and thereby increases the load of misfolded proteins transiting the endoplasmic reticulum (29). In *C. elegans*, Tm reliably induces the UPR, presumably through its known activity to increase protein misfolding (8, 48). We pretreated wild-type *C. elegans* for 4 h with various concentrations of Tm and then allowed the animals to recover for 20 h before beginning a normally lethal hypoxic incubation. Tm induced a significant protection from hypoxia in a concentration-dependent manner (Fig. 2A and B). The onset and duration of the tunicamycin preconditioning (TmP) after recovery from Tm incubation was delayed (Fig. 2C). Significant protection compared to control incubations in buffer was not seen until 14 h after the Tm incubation. Maximal TmP was seen after a recovery time of 16 h, and the protection persisted for at least 24 h, although the magnitude of the protection was waning at that point. Notably, Tm present during the hypoxic incubation was not protective (data not shown). Thus, the




FIG. 1. Hypoxia and HP activate the UPR. HP activates an *hsp-4* promoter-GFP fusion reporter (A and B) and the endogenous *hsp-4* gene (C). (B) After HP, the level of GFP increased significantly after a 4-h recovery (*, $P < 0.001$, two-tailed t test), and it returned to the control level after 16 more hours. (C) The level of *hsp-4* mRNA was measured by qRT-PCR and was significantly elevated after a 2- and 4-h recovery from 4 h of hypoxia ($P < 0.05$, two-tailed t test), while hypoxia alone (up to 8 h) or a shorter HP incubation had no effect. (D) Hypoxia induced eIF2 α phosphorylation. The level of phosphorylated eIF2 α increased after 1 h of hypoxia and remained high under hypoxic conditions but rapidly returned to baseline during normoxic recovery. Relative band intensities normalized to no hypoxia are given. β -Actin levels decreased relative to total protein during the hypoxic incubation, thus the p-eIF2 α / β -actin ratio increased greatly. The 0-h hypoxia/0 recovery and the 4-h hypoxia/0 recovery conditions were repeated for a total of four trials, and the relative p-eIF2 α induction (1.96 ± 0.29) was statistically significant ($P < 0.01$, paired t test).

protective effect was a delayed reaction to the Tm and not due to the Tm itself. Buffer incubations also induced a hypoxic protection but with a time course distinct from that of TmP. The buffer-induced protection peaked at 4 h and fell thereafter. We speculate that the buffer induces a starvation stress or that the act of swimming in buffer for 4 h induces a brief protective stress response.

To determine whether TmP was mediated by the induction of the UPR, we measured TmP in reduction-of-function mutations in UPR genes (Fig. 3) with the hypothesis that one or more pathways within the UPR are necessary for the induction of TmP. A large deletion mutation in *pek-1* or *gcn-2* did not block Tm preconditioning (Fig. 2D and 3C, E). On the other hand, three loss-of-function mutant alleles of *ire-1* (Fig. 3B), two alleles of *atf-6* (Fig. 3D), and an allele of *xbp-1* (Fig. 3F) all were defective for Tm preconditioning (Fig. 2D). Indeed, in the *xbp-1(lf)* allele, Tm preincubation significantly reduced survival from hypoxia. These data indicate that the *ire-1*, *xbp-1*, and *atf-6* branches but not the *pek-1* or *gcn-2* branches of the UPR are essential for TmP. GRP78/Bip is a family of ER chaperones that regulate the UPR by binding to misfolded proteins, resulting in the disinhibition of UPR components, including homologs of IRE-1 and ATF-6 (42). *C. elegans* has two Bip homologs, *hsp-3* and *hsp-4*, both of which are activated by ER stress (48). A reasonable hypothesis is that an early step

in tunicamycin preconditioning is the activation of HSP-3 or HSP-4, which then disinhibit IRE-1 and ATF-6 and promote hypoxic protection. However, null mutations in *hsp-3* or *hsp-4* (Fig. 3G, H) neither blocked nor enhanced Tm preconditioning (Fig. 2D).

***ire-1* and *gcn-2* are required for hypoxic preconditioning.** We next tested whether the same UPR components were required for hypoxic preconditioning (HP). As for Tm preconditioning, HP consistently provided protection from subsequent harsh hypoxic exposure for wild-type animals (Fig. 4A and B). Again, the *pek-1* deletion mutation had no effect on HP; *pek-1(ok275)* animals were strongly protected by HP (Fig. 4B). However, unlike for TmP, *gcn-2(ok871)* completely blocked HP (Fig. 4C). *gcn-2(ok886)*, an allele with a smaller deletion that removes less of the kinase and tRNA-binding domains (Fig. 3E), also failed to exhibit a significant increase in survival after HP, although there was a trend toward protection (Fig. 4C). Two *ire-1* alleles (*v33* and *ok799*) blocked HP (Fig. 4C); however, unlike the case for Tm preconditioning, *ire-1(zc14)*, a missense mutation in the kinase domain that is thought to abolish the XBP-1 endonuclease activity of IRE-1 and behaves as a reduction-of-function allele (Fig. 3B) (8), did not block HP (Fig. 4C). Also, unlike the case for TmP, neither *atf-6* nor *xbp-1* mutation blocked HP (Fig. 4C). As for TmP, the mutation of *hsp-3* or *hsp-4* had no effect on HP (Fig. 4C). These data show




FIG. 2. Tunicamycin-induced hypoxic protection. (A) Tunicamycin (Tm) pretreatment induced protection from subsequent hypoxic injury. Worms were treated with the indicated concentrations of Tm for 4 h before being recovered for 16 h. After recovery, Tm-pretreated worms were challenged with hypoxia for 22 h, and survival was scored after another 24-h recovery. Values are means \pm standard deviations (SD) from three trials (* $P < 0.001$, paired t test, Tm versus buffer control). (C) Time course of Tm-induced hypoxia protection. The experiment was performed as described above with 10 μ g/ml Tm or buffer only with various recovery times prior to the 22-h hypoxic exposure. The control value is for animals receiving no pretreatment as opposed to buffer pretreatment. Values are means \pm SD from three trials (*, $P < 0.01$, paired t test, Tm versus buffer). (D) Wild-type (N2) or mutant animals were tested for Tm (10 μ g/ml)-induced hypoxia protection. Animals were exposed for 4 h to Tm or buffer control and then recovered for 20 h prior to a 22-h hypoxic exposure, and then they were scored 24 h later for survival. Net survival (Tm survival – buffer survival) was calculated for each genotype. Tm induced significant hypoxic protection compared to that by buffer ($P < 0.01$, paired t test) in all strains except for the *ire-1*, *atf-6*, and *xbp-1* mutants. Each bar represents the means \pm SD from a minimum of three independent trials with at least 30 animals/trial. *, $P < 0.01$, paired t test, Tm versus buffer.

that Tm preconditioning and hypoxic preconditioning both require an intact UPR but that the mechanisms are not identical. In common to both TmP and HP is a requirement for IRE-1.

Role of *ire-1* in hypoxic injury. Given the unique role among UPR components of IRE-1 in both HP and Tm preconditioning, we wanted to compare the native hypoxic sensitivity of *ire-1* alleles to that of the other UPR mutants. As previously reported (1), the missense allele *ire-1(zc14)* was significantly hypoxia resistant (Fig. 5A). In addition, *ire-1(tm400)*, a deletion allele that has the potential for an alternative translation start site downstream of the deletion (see Materials and Methods), also was hypoxia resistant. However, the two other *ire-1* deletion alleles were not hypoxia resistant. Likewise, none of the other UPR mutants were hypoxia resistant. The apparent less-severe phenotype of *ire-1(v33)* compared to those of *zc14* and *tm400* was particularly surprising given that the *v33* deletion mutation results in a frameshift and an early stop codon and is presumably a null mutation (Fig. 3B) (48). Three mechanisms might explain this result. First, other unknown mutations in the *zc14* and *tm400* mutant strains might be responsible for the resistance. Second, IRE-1 might have both hypoxic sensitivity promoting and blocking activities and *zc14* and *tm400* only disrupt the promoting activity. Third, hypoxic sensitivity might have a biphasic response to the level of activity of

IRE-1 so that the complete absence of IRE-1 function is deleterious but a partially reduced activity can protect from hypoxia death. To distinguish between these mechanisms, we measured the hypoxic sensitivity of heterozygous and transheterozygous *ire-1(zc14)*, *tm400*, and *v33* mutants (Fig. 5B). *zc14*/balancer heterozygous animals had a hypoxic sensitivity similar to that of wild-type and balancer/+ animals. However, *v33* and *tm400* heterozygotes were strongly hypoxia resistant. *zc14/v33* transheterozygotes had a hypoxic sensitivity similar to that of *zc14* homozygous animals. These data are most consistent with the third hypothesis that hypoxic sensitivity is reduced with partial but not complete loss of *ire-1* function.

We then tested if the tunicamycin sensitivities of the *ire-1* allelic combinations mirrored their hypoxic sensitivity. Indeed, in an assay of Tm-induced developmental arrest, both *zc14* and *tm400* homozygotes and *v33* heterozygotes were Tm resistant, whereas *v33* homozygotes were Tm hypersensitive, as had been reported previously (Fig. 5C) (48). Consistent with an early stop mutation and putative null phenotype of *ire-1(v33)*, the transcript levels of *v33* homozygotes were about 10-fold lower than that in wild-type or *zc14* mutant animals. *v33* heterozygotes had significantly reduced *ire-1* mRNA levels as well, which is consistent with a haploinsufficient phenotype seen in *v33* heterozygotes (Fig. 5D). This correspondence of Tm and hypoxic sensitivity phenotypes is consistent with the hypothesis




FIG. 3. UPR pathway and mutants. (A) Schematic of UPR pathways. PEK-1, IRE-1, and ATF-6 are activated in the presence of unfolded proteins in the ER lumen. These pathways can promote adaptation to unfolded proteins via translational suppression or through a transcriptional response. GCN-2 functions along with activated PEK-1 to suppress translation. (B) *ire-1* mutations. *ire-1(v33)* has an N-terminal 878-bp deletion resulting in a frameshift and stop and is a presumptive null mutation (48). *ire-1(ok799)* has a 2,093-bp deletion and 409-bp insertion and also should represent a null mutation (50). *ire-1(zc14)* has a missense mutation in a conserved residue in the kinase domain (8). *ire-1(tm400)* has a 600-bp deletion and 1-bp insertion that ends in an intron (see Wormbase.org and Materials and Methods). The mutant product is unclear. (C) *pek-1(ok275)* has a 2,073-bp deletion that produces a frameshift and stop codon (48). (D) Proteolysis of ATF-6 produces ATF-6s with only the maroon domain that is truncated by both mutations. *ok551* has a 1,900-bp deletion (49); *tm1153* has a 643-bp frameshift deletion (Wormbase.org). (E) *gcn-2(ok871)* has a 1,481-bp in-frame deletion starting and ending in exons (33); *gcn-2(ok886)* has a 1,179-bp in-frame deletion that starts and ends in exons (33). (F) *xbp-1(zc12)* has an early stop (8). (G) *hsp-3(ok1083)* has a 1,422-bp deletion that starts and ends in exons, causing frameshift (22). (H) *hsp-4(gk514)* has a 752-bp deletion that starts and ends in exons, causing frameshift (46). TM, transmembrane domain. The RWD domain was named after three major RWD-containing proteins: RING finger-containing proteins, WD-repeat-containing proteins, and yeast DEAD (DEXD)-like helicases. ΨPK, degenerate kinase domain; PK, kinase domain; HisRS, histidyl-tRNA synthetase; RB/DD, ribosome-binding and dimerization domain.

that the biphasic effect of reducing IRE-1 activity on hypoxic sensitivity is due to the response to unfolded protein stress.

Mechanism of *gcn-2*- and *ire-1*-mediated HP and hypoxia resistance. The best-established target of the GCN-2 kinase is the translation factor eIF2α. To determine whether the increase in p-eIF2α during the hypoxic preconditioning incubation (Fig. 1D) is required for the subsequent induction of HP, we measured p-eIF2α levels in the wild type and the HP-defective mutant, *gcn-2(ok871)*. In both strains, p-eIF2α levels were similarly and significantly increased relative to that of β-actin during the 4-h hypoxic preconditioning incubation (Fig. 6A and B). On the other hand, the significant hypoxic induction of p-eIF2α was blocked in *pek-1(ok275)*, a mutant with normal HP (Fig. 6C). Thus, the phosphorylation of eIF2α is neither necessary nor sufficient for HP, and the relevant GCN-2 target is unknown.

While IRE-1 has other known downstream targets (15, 20, 21, 27), XBP-1 is the best characterized. XBP-1 clearly is not

required for HP, as an *xbp-1(lf)* mutant exhibits a normal HP response (Fig. 4C). However, this result does not rule out the possibility that XBP-1 acts redundantly to induce HP or to regulate hypoxic sensitivity in general. Thus, we asked whether *ire-1* allelic differences for HP and hypoxic sensitivity phenotypes correlated with XBP-1 splicing. All three *ire-1* alleles failed to produce detectable levels of spliced XBP-1 under normal conditions or after an HP incubation (Fig. 6D and E). These data, along with the wild-type hypoxia/HP phenotypes of *xbp-1(lf)*, indicate that IRE-1 controls HP and baseline hypoxic sensitivity through an XBP-1-independent mechanism.

DISCUSSION

We showed that hypoxic preconditioning in *C. elegans* induces unfolded protein response pathways. We also found that preincubation with tunicamycin, a drug that promotes protein misfolding, is capable of producing a delayed hypoxia protec-




FIG. 4. UPR components required for hypoxic preconditioning (HP). (A and B) Wild-type (N2) animals were exposed to hypoxia (HP) or normoxia (control) incubations for 4 h and then allowed to recover for 20 h prior to a 20-h hypoxic incubation. Survival was scored after another 24-h recovery. (C) Survival from HP in the wild type and UPR mutants. Net survival (survival of HP-treated animals – survival of control animals) is plotted for each genotype. Each bar represents the means \pm standard deviations from a minimum of three independent trials with at least 30 animals/trial. *, $P < 0.01$, paired t test, HP versus control.

tion similar to that of delayed HP. Finally, we showed that distinct but overlapping components of the unfolded protein response are required for hypoxic preconditioning and tunicamycin preconditioning. These results suggest a model for hypoxic preconditioning where misfolded proteins serve as early hypoxic sensors that then signal through IRE-1 to induce an adaptive hypoxia protective response along with essential signaling from GCN-2 (Fig. 7). We now would like to place our results in the context of previous studies of protein misfolding, the UPR, and hypoxic injury/preconditioning.

A number of studies have suggested a role for the UPR in the hypoxic/ischemic injury of mammalian cells. Many studies have demonstrated an increase in UPR transcripts and protein in models of ischemic injury and preconditioning (2, 17, 39, 43, 54, 57). The induction of the UPR by hypoxia is consistent with studies of hypoxic tumor cells in which the UPR is activated (24). However, only a few studies have attempted to modulate the UPR to determine its functional role, if any, in hypoxic/ischemic injury. Tajiri et al. showed that hippocampal neurons from CHOP knockout mice were resistant to hypoxia-induced apoptosis (53). CHOP (C/EBP homologous protein) is a transcription factor induced by ER stress and is a target of ATF-6 and IRE-1; CHOP regulates the transcription of a number of genes that in general promote apoptosis (38). However, *C. elegans* does not have a known CHOP homolog. In the mouse kidney, pretreatment with tunicamycin was found to produce a long-lasting protection from ischemic injury (41). The mechanism of the protection was undetermined, although tunicamycin was shown to increase XBP-1 and GRP78 mRNA levels. Most recently, the modulation of ATF-6 has proven to be an important determinant of hypoxic/ischemic injury. The cardiac-

specific expression of an inducible form of ATF6 in transgenic mice was found to markedly reduce cardiac myocyte death after ischemia in isolated hearts (28). On the other hand, the short hairpin RNA knockdown of ATF6 expression in primary rat myocyte cultures increased cell death after a severe hypoxic incubation (13). In *C. elegans*, we find that a deletion mutant of *atf-6* blocks the induction of protection from hypoxia by tunicamycin, which is consistent with a role of ATF-6 in promoting hypoxic protection.

The requirements for the ATF-6 and IRE-1-XBP-1 pathways but not PEK-1 in Tm-induced hypoxic protection are intriguing. First, the requirements of both XBP-1 and ATF-6 are consistent with previous reports showing that these two transcription factors can target the same promoter elements and therefore coregulate the expression of certain ER stress response genes (59). XBP-1 and ATF-6 can heterodimerize as a prerequisite for binding to some promoter elements (58). Additionally, XBP-1 and ATF-6 are coordinately regulated by IRE-1, which is required for the activation of both proteins (26, 60). As for the role of *pek-1*, despite the rich literature showing that PEK-1 activation is prosurvival in ischemic/hypoxic cell death (4, 11, 25), we found no evidence for a role of PEK-1 in hypoxic cell death or preconditioning. In general in *C. elegans*, PEK-1 has been found to be dispensable for the UPR. For example, the inductions of *apy-1* and Rho subfamily member *crp-1* are controlled by *ire-1*, *xbp-1*, and *atf-6* but not *pek-1* (9, 55). Pore-forming toxins also activate *ire-1*, *xbp-1*, and *atf-6* but not *pek-1* in worms (5). However, we did find that the hypoxia-induced phosphorylation of eIF2 α required functional PEK-1.

The finding that GCN-2 is required for HP was surprising.




FIG. 5. Effect of UPR mutants on hypoxic and tunicamycin sensitivity. (A) Animals with the indicated alleles were exposed to hypoxia for 20 h without any pretreatment, and survival was scored after a 24-h recovery (*, $P < 0.0001$ versus N2 by unpaired t test). (B) Animals with different *ire-1* genetic backgrounds were tested for hypoxic survival without any pretreatment. (*, $P < 0.001$ versus N2; #, $P < 0.05$ versus *zc14/zc14* or *zc14/v33*, unpaired t test). (C) Animals with different *ire-1* genetic backgrounds were tested for sensitivity to Tm toxicity. Eggs were laid on the plates with 1 μ g/ml Tm. After 3 days, the percentage of adult worms was scored (*, $P < 0.01$ for results greater than those for N2; #, $P < 0.01$ for results less than those for N2, unpaired t test). (D) The levels of *ire-1* mRNA from the wild type and mutants were determined by quantitative RT-PCR by using a primer pair annealed 5' of *ire-1* cDNA. (*, $P < 0.01$, versus N2, *zc14/zc14*, or *zc14/mIn1* [*dpy-10(e128) mIs14(p-myo-2::GFP)*], unpaired t test).

GCN-2 is homologous to the only eIF2 α kinase found in yeast. In yeast, the *gcn-2* homolog has been found to be essential for UPR function, so it also has been known as a super-UPR component in yeast (40). However, GCN-2 was not required for hypoxia-induced eIF2 α phosphorylation (Fig. 6B) nor for TmP (Fig. 2D). Also, GCN-2 doesn't appear to be directly activated by ER stress in mammalian cells (16). The target of GCN-2 signaling in the context of HP is unknown; however, the yeast bZIP transcription factor Gcn4, a homolog of mammalian ATF4, functions downstream of Gcn2 and independently of eIF2 α to regulate yeast UPR target genes (40). In yeast, Gcn2 appears to be required for the basal expression of Gcn4, which is further activated by Ire1 during ER stress to promote UPR gene transcription. Our data are consistent with cooperativity between GCN-2 and IRE-1, but the downstream pathways in *C. elegans* are undefined.

IRE-1 was unique among the UPR genes in having essential roles in both tunicamycin and hypoxic preconditioning. However, the transduction pathway downstream of IRE-1 was distinct for the two preconditioning conditions; tunicamycin preconditioning required XBP-1, whereas hypoxic preconditioning did not. IRE-1 is classically thought to function in a

linear pathway with its downstream target being the transcription factor XBP-1. However, potential XBP-1-independent functions of IRE-1 have been reported and fall into two broad classes, mRNA degradation and protein-protein interactions (19). Regulated Ire1-dependent decay of mRNAs (RIDD) was defined originally in *Drosophila melanogaster* cells (21) and subsequently demonstrated in mammalian cells (15, 20, 37). Whether a RIDD mechanism functions in *C. elegans* is unknown. Various aspects of the RIDD pathway are similar to the role of the UPR in HP in *C. elegans*. Like HP, RIDD appears not to require XBP-1 (20). Second, in the context of ER stress, RIDD mechanisms can be induced pharmacologically and bypass the requirement for the kinase activity of IRE1 in mouse embryonic fibroblasts (20). The normal HP phenotype of the *ire-1* kinase domain mutant *zc14* suggests that IRE-1 kinase activity also is not essential for HP, although the kinase activity of *zc14* has not been directly assayed. However, Han et al. found in an HEK293-derived cell line that IRE1 kinase activity was required for RIDD (15), so the resemblance of HP and RIDD with regard to IRE-1 kinase activity is unclear. Another issue with RIDD and its role in HP is the timing. Protection after hypoxic preconditioning in *C. elegans*




FIG. 6. eIF2 α phosphorylation and XBP-1 splicing after HP. (A to C) N2, *pek-1(ok275)*, and *gcn-2(ok871)* animals were treated with hypoxia for 4 h and recovered under normoxia. Protein samples from various recovery time points were subjected to Western blotting and were probed by an antibody against p-eIF2 α (ser51). The same blots were stripped and reprobed with a β -actin antibody. Intensity values for the p-eIF2 α bands normalized to the control are indicated along with the ratio of the normalized intensities of the p-eIF2 α bands to β -actin. Four independent trials of the control and 0-h recovery time point gave normalized p-eIF2 α intensities of 1.96 ± 0.29 for N2, 1.80 ± 0.07 for *gcn-2(ok871)*, and 1.14 ± 0.15 for *pek-1(ok275)*. *P* values for the change in intensities (paired *t* test) were the following: for N2, 0.00531; for *gcn-2(ok871)*, 0.00028; and for *pek-1(ok275)*, 0.399. (D and E) The unspliced and spliced forms of *xbp-1* mRNA were amplified by RT-PCR in N2 and *ire-1* mutant animals under control conditions (D) and after a 2-h recovery from a 4-h HP incubation (E). Spliced *xbp-1* was undetectable in all three *ire-1* mutant alleles under both conditions.

occurs about 16 h after preconditioning (10), whereas RIDD is thought to act fully within 4 to 8 h to degrade certain RNAs (15, 21). These disparate time courses suggest that RIDD itself is not the effector of protection after preconditioning; rather, if involved in HP, RIDD would act earlier and upstream of the effector mechanism, perhaps in transducing the preconditioning stimulus.

Alternatively, IRE-1 could control HP in *C. elegans* via proteins that have been found to interact with IRE1 (27). Mammalian IRE1 has been shown to form a complex with TRAF2, ASK-1, JNK, and ASK1-interacting protein (AIP1), which together promote apoptotic cell death in models of ER stress (23, 27, 32, 56). This IRE1 pathway is thought to be independent of XBP1 because an endonuclease-deficient IRE1 was




FIG. 7. Working model for the role of the UPR in HP and TmP in *C. elegans*. Both hypoxia and tunicamycin inhibit protein folding and thereby activate signaling through IRE-1 and ATF-6 pathways. IRE-1 is required for both HP and TmP. GCN-2 is required for HP only, and ATF-6 is required for TmP only. The mechanisms downstream of IRE-1 and GCN-2 to induce HP are unknown.

competent to interact with TRAF2 and activate JNK (56). Paradoxically, in *C. elegans*, the overexpression of the *C. elegans* homolog of JNK, JNK-1, increases life span and thermal and oxidative stress resistance (36). Thus, a plausible hypothesis is that limited hypoxia activates IRE-1, stimulating JNK-1, which promotes the transcription of proadaptive gene products, perhaps similar to those that increase life span and stress resistance. More prolonged hypoxia also could act through an IRE-1/JNK-1 pathway to promote cell death, the more typical output of the JNK1 pathway in mammalian models. One output of the JNK pathway that might reasonably regulate HP is macroautophagy. Autophagy has been shown to protect against ER stress (3, 35) and is activated by ER stress by an IRE1-, JNK-, and TRAF2-dependent mechanism (35). We have previously shown that macroautophagy is activated by hypoxia and protects against hypoxic injury in *C. elegans* (45). Thus, the activation of autophagy is a plausible candidate as the effector of IRE-1-dependent HP. Another IRE1-interacting protein is USP14. USP14, a ubiquitin-specific protease, has been shown to interact directly with IRE1 α in HEK293 cells (31). Kinase-dead IRE1 α is capable of recruiting USP14 to a complex that includes members of the ER-associated protein degradation (ERAD) machinery. The association of UPR14 with kinase-dead IRE1 α correlated with the inhibition of ERAD, whereas autophosphorylated IRE1 α did not bind UPR14 and was incompetent for UPR14-mediated ERAD inhibition. Although not directly tested, this IRE1 function presumably would be independent of XBP-1 and therefore have characteristics consistent with the mediation of HP. Future studies will be aimed at elucidating the mechanism(s) whereby IRE-1 and GCN-2 regulate HP in *C. elegans* and determining whether this mechanism is operant in mammalian cells.

ACKNOWLEDGMENTS

This work was supported by R01-NS045905 from the National Institute of Neurological Disorders and Stroke, the McKnight Endowment Fund for Neuroscience, the American Heart Association, and the International Anesthesia Research Society.

REFERENCES

- Anderson, L. L., X. Mao, B. A. Scott, and C. M. Crowder. 2009. Survival from hypoxia in *C. elegans* by inactivation of aminoacyl-tRNA synthetases. *Science* **323**:630–633.
- Azfer, A., J. Niu, L. M. Rogers, F. M. Adamski, and P. E. Kolattukudy. 2006. Activation of endoplasmic reticulum stress response during the development of ischemic heart disease. *Am. J. Physiol. Heart Circ. Physiol.* **291**:H1411–H1420.
- Bernales, S., K. L. McDonald, and P. Walter. 2006. Autophagy counterbalances endoplasmic reticulum expansion during the unfolded protein response. *PLoS Biol.* **4**:e423.
- Bi, M., C. Naczi, M. Koritzinsky, D. Fels, J. Blais, N. Hu, H. Harding, I. Novoa, M. Varia, J. Raleigh, D. Scheuner, R. J. Kaufman, J. Bell, D. Ron, B. G. Wouters, and C. Koumenis. 2005. ER stress-regulated translation increases tolerance to extreme hypoxia and promotes tumor growth. *EMBO J.* **24**:3470–3481.
- Bischof, L. J., C. Y. Kao, F. C. Los, M. R. Gonzalez, Z. Shen, S. P. Briggs, F. G. van der Goot, and R. V. Aronin. 2008. Activation of the unfolded protein response is required for defenses against bacterial pore-forming toxin in vivo. *PLoS Pathog.* **4**:e1000176.
- Bolli, R. 2007. Preconditioning: a paradigm shift in the biology of myocardial ischemia. *Am. J. Physiol. Heart Circ. Physiol.* **292**:H19–H27.
- Brenner, S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* **77**:71–94.
- Calfon, M., H. Zeng, F. Urano, J. H. Till, S. R. Hubbard, H. P. Harding, S. G. Clark, and D. Ron. 2002. IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* **415**:92–96.
- Caruso, M. E., S. Jenna, M. Bouchecareilh, D. L. Baillie, D. Boismenu, D. Halawani, M. Latterich, and E. Chevet. 2008. GTPase-mediated regulation of the unfolded protein response in *Caenorhabditis elegans* is dependent on the AAA+ ATPase CDC-48. *Mol. Cell.* **28**:4261–4274.
- Dasgupta, N., A. M. Patel, B. A. Scott, and C. M. Crowder. 2007. Hypoxic preconditioning requires the apoptosis protein CED-4 in *C. elegans*. *Curr. Biol.* **17**:1954–1959.
- DeGracia, D. J. 2004. Acute and persistent protein synthesis inhibition following cerebral reperfusion. *J. Neurosci. Res.* **77**:771–776.
- Dirnagl, U., K. Becker, and A. Meisel. 2009. Preconditioning and tolerance against cerebral ischaemia: from experimental strategies to clinical use. *Lancet Neurol.* **8**:398–412.
- Doroudgar, S., D. J. Thuerlauf, M. C. Marcinko, P. J. Belmont, and C. C. Glembotski. 2009. Ischemia activates the ATF6 branch of the endoplasmic reticulum (ER) stress response. *J. Biol. Chem.* **284**:29735–29745.
- Edgley, M. L., and D. L. Riddle. 2001. LG II balancer chromosomes in *Caenorhabditis elegans*: mT1(II;III) and the mIn1 set of dominantly and recessively marked inversions. *Mol. Genet. Genomics* **266**:385–395.
- Han, D., A. G. Lerner, L. Vande Walle, J.-P. Upton, W. Xu, A. Hagen, B. J. Backes, S. A. Oakes, and F. R. Papa. 2009. IRE1 α kinase activation modes control alternate endonuclease outputs to determine divergent cell fates. *Cell* **138**:562–575.
- Harding, H. P., I. Novoa, Y. Zhang, H. Zeng, R. Wek, M. Schapira, and D. Ron. 2000. Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol. Cell* **6**:1099–1108.
- Hayashi, T., A. Saito, S. Okuno, M. Ferrand-Drake, and P. H. Chan. 2003. Induction of GRP78 by ischemic preconditioning reduces endoplasmic reticulum stress and prevents delayed neuronal cell death. *J. Cereb. Blood Flow Metab.* **23**:949–961.
- Hayashi, T., A. Saito, S. Okuno, M. Ferrand-Drake, R. L. Dodd, T. Nishi, C. M. Maier, H. Kinouchi, and P. H. Chan. 2003. Oxidative damage to the endoplasmic reticulum is implicated in ischemic neuronal cell death. *J. Cereb. Blood Flow Metab.* **23**:1117–1128.
- Hetz, C., and L. H. Glimcher. 2009. Fine-tuning of the unfolded protein response: assembling the IRE1 α interactome. *Mol. Cell* **35**:551–561.
- Hollien, J., J. H. Lin, H. Li, N. Stevens, P. Walter, and J. S. Weissman. 2009. Regulated Ire1-dependent decay of messenger RNAs in mammalian cells. *J. Cell Biol.* **186**:323–331.
- Hollien, J., and J. S. Weissman. 2006. Decay of endoplasmic reticulum-localized mRNAs during the unfolded protein response. *Science* **313**:104–107.
- Kapulkin, W. J., B. G. Hiester, and C. D. Link. 2005. Compensatory regulation among ER chaperones in *C. elegans*. *FEBS Lett.* **579**:3063–3068.
- Kim, I., C. W. Shu, W. Xu, C. W. Shiau, D. Grant, S. Vasile, N. D. Cosford, and J. C. Reed. 2009. Chemical biology investigation of cell death pathways activated by endoplasmic reticulum stress reveals cytoprotective modulators of ASK1. *J. Biol. Chem.* **284**:1593–1603.
- Koumenis, C., M. Bi, J. Ye, D. Feldman, and A. C. Koong. 2007. Hypoxia and the unfolded protein response. *Methods Enzymol.* **435**:275–293.
- Koumenis, C., C. Naczi, M. Koritzinsky, S. Rastani, A. Diehl, N. Sonenberg, A. Koromilas, and B. G. Wouters. 2002. Regulation of protein synthesis by hypoxia via activation of the endoplasmic reticulum kinase PERK and phosphorylation of the translation initiation factor eIF2 α . *Mol. Cell Biol.* **22**:7405–7416.
- Lee, K., W. Tirasophon, X. Shen, M. Michalak, R. Prywes, T. Okada, H. Yoshida, K. Mori, and R. J. Kaufman. 2002. IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev.* **16**:452–466.
- Luo, D., Y. He, H. Zhang, L. Yu, H. Chen, Z. Xu, S. Tang, F. Urano, and W. Min. 2008. AIP1 is critical in transducing IRE1-mediated endoplasmic reticulum stress response. *J. Biol. Chem.* **283**:11905–11912.
- Martindale, J. J., R. Fernandez, D. Thuerlauf, R. Whittaker, N. Gude, M. A. Sussman, and C. C. Glembotski. 2006. Endoplasmic reticulum stress gene induction and protection from ischemia/reperfusion injury in the hearts of transgenic mice with a tamoxifen-regulated form of ATF6. *Circ. Res.* **98**:1186–1193.
- Merksamer, P. I., A. Trusina, and F. R. Papa. 2008. Real-time redox measurements during endoplasmic reticulum stress reveal interlinked protein folding functions. *Cell* **135**:933–947.
- Murry, C. E., R. B. Jennings, and K. A. Reimer. 1986. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation* **74**:1124–1136.
- Nagai, A., H. Kadowaki, T. Maruyama, K. Takeda, H. Nishitoh, and H. Ichijo. 2009. USP14 inhibits ER-associated degradation via interaction with IRE1 α . *Biochem. Biophys. Res. Commun.* **379**:995–1000.
- Nishitoh, H., A. Matsuzawa, K. Tobiume, K. Saegusa, K. Takeda, K. Inoue, S. Hori, A. Kakizuka, and H. Ichijo. 2002. ASK1 is essential for endoplasmic reticulum stress-induced neuronal cell death triggered by expanded polyglutamine repeats. *Genes Dev.* **16**:1345–1355.
- Nukazuka, A., H. Fujisawa, T. Inada, Y. Oda, and S. Takagi. 2008. Semaphorin controls epidermal morphogenesis by stimulating mRNA translation via eIF2 α in *Caenorhabditis elegans*. *Genes Dev.* **22**:1025–1036.
- Obrenovitch, T. P. 2008. Molecular physiology of preconditioning-induced brain tolerance to ischemia. *Physiol. Rev.* **88**:211–247.
- Ogata, M., S. Hino, A. Saito, K. Morikawa, S. Kondo, S. Kanemoto, T. Murakami, M. Taniguchi, I. Tani, K. Yoshinaga, S. Shiosaka, J. A. Hammarback, F. Urano, and K. Imaizumi. 2006. Autophagy is activated for cell survival after endoplasmic reticulum stress. *Mol. Cell Biol.* **26**:9220–9231.
- Oh, S. W., A. Mukhopadhyay, N. Srivizaka, F. Jiang, R. J. Davis, and H. A. Tissenbaum. 2005. JNK regulates lifespan in *Caenorhabditis elegans* by modulating nuclear translocation of forkhead transcription factor/DAF-16. *Proc. Natl. Acad. Sci. U. S. A.* **102**:4494–4499.
- Oikawa, D., M. Tokuda, and T. Iwakaki. 2007. Site-specific cleavage of CD59 mRNA by endoplasmic reticulum-localized ribonuclease, IRE1. *Biochem. Biophys. Res. Commun.* **360**:122–127.
- Oyadomari, S., and M. Mori. 2004. Roles of CHOP/GADD153 in endoplasmic reticulum stress. *Cell Death Differ.* **11**:381–389.
- Paschen, W., C. Auenberg, S. Hotop, and T. Mengesdorf. 2003. Transient cerebral ischemia activates processing of xbp1 messenger RNA indicative of endoplasmic reticulum stress. *J. Cereb. Blood Flow Metab.* **23**:449–461.
- Patil, C. K., H. Li, and P. Walter. 2004. Gcn4p and novel upstream activating sequences regulate targets of the unfolded protein response. *PLoS Biol.* **2**:E246.
- Prachasilchai, W., H. Sonoda, N. Yokota-Ikeda, S. Oshikawa, C. Aikawa, K. Uchida, K. Ito, T. Kudo, K. Imaizumi, and M. Ikeda. 2008. A protective role of unfolded protein response in mouse ischemic acute kidney injury. *Eur. J. Pharmacol.* **592**:138–145.
- Quinones, Q. J., G. G. de Ridder, and S. V. Pizzo. 2008. GRP78: a chaperone with diverse roles beyond the endoplasmic reticulum. *Histol. Histopathol.* **23**:1409–1416.
- Roberts, G. G., M. J. Di Loreto, M. Marshall, J. Wang, and D. J. DeGracia. 2007. Hippocampal cellular stress responses after global brain ischemia and reperfusion. *Antioxid. Redox. Signal.* **9**:2265–2275.
- Rzymiski, T., and A. L. Harris. 2007. The unfolded protein response and integrated stress response to anoxia. *Clin. Cancer Res.* **13**:2537–2540.
- Samokhvalov, V., B. A. Scott, and C. M. Crowder. 2008. Autophagy protects against hypoxic injury in *C. elegans*. *Autophagy* **4**:1034–1041.
- Sasagawa, Y., K. Yamanaka, and T. Ogura. 2007. ER E3 ubiquitin ligase HRD-1 and its specific partner chaperone BiP play important roles in ERAD and developmental growth in *Caenorhabditis elegans*. *Genes Cells* **12**:1063–1073.
- Scott, B. A., M. S. Avidan, and C. M. Crowder. 2002. Regulation of hypoxic death in *C. elegans* by the insulin/IGF receptor homolog DAF-2. *Science* **296**:2388–2391.
- Shen, X., R. E. Ellis, K. Lee, C. Y. Liu, K. Yang, A. Solomon, H. Yoshida, R. Morimoto, D. M. Kurnit, K. Mori, and R. J. Kaufman. 2001. Complementary signaling pathways regulate the unfolded protein response and are required for *C. elegans* development. *Cell* **107**:893–903.
- Shen, X., R. E. Ellis, K. Sakaki, and R. J. Kaufman. 2005. Genetic interac-

- tions due to constitutive and inducible gene regulation mediated by the unfolded protein response in *C. elegans*. *PLoS Genet.* **1**:e37.
50. Shim, J., T. Umemura, E. Nothstein, and C. Rongo. 2004. The unfolded protein response regulates glutamate receptor export from the endoplasmic reticulum. *Mol. Biol. Cell* **15**:4818–4828.
 51. Sommerschild, H. T., and K. A. Kirkeboen. 2002. Preconditioning-endogenous defence mechanisms of the heart. *Acta Anaesthesiol. Scand.* **46**:123–137.
 52. Stiernagle, T. 11 February 2006, posting date. Maintenance of *C. elegans*. In D. Fay (ed.), *Wormbook*. The *C. elegans* Research Community. doi:10.1895/wormbook.1.101.1.
 53. Tajiri, S., S. Oyadomari, S. Yano, M. Morioka, T. Gotoh, J. I. Hamada, Y. Ushio, and M. Mori. 2004. Ischemia-induced neuronal cell death is mediated by the endoplasmic reticulum stress pathway involving CHOP. *Cell Death Differ.* **11**:403–415.
 54. Truettner, J. S., K. Hu, C. L. Liu, W. D. Dietrich, and B. Hu. 2009. Subcellular stress response and induction of molecular chaperones and folding proteins after transient global ischemia in rats. *Brain Res.* **1249**:9–18.
 55. Uccelletti, D., A. Pascoli, F. Farina, A. Alberti, P. Mancini, C. B. Hirschberg, and A. C. Palleschi. 2008. APY-1, a novel *Caenorhabditis elegans* apyrase involved in unfolded protein response signalling and stress responses. *Mol. Biol. Cell* **19**:1337–1345.
 56. Urano, F., X. Wang, A. Bertolotti, Y. Zhang, P. Chung, H. P. Harding, and D. Ron. 2000. Coupling of stress in the ER to activation of JNK protein kinases by transmembrane protein kinase IRE1. *Science* **287**:664–666.
 57. Wang, S., F. M. Longo, J. Chen, M. Butman, S. H. Graham, K. G. Haglid, and F. R. Sharp. 1993. Induction of glucose regulated protein (grp78) and inducible heat shock protein (hsp70) mRNAs in rat brain after kainic acid seizures and focal ischemia. *Neurochem. Int.* **23**:575–582.
 58. Yamamoto, K., T. Sato, T. Matsui, M. Sato, T. Okada, H. Yoshida, A. Harada, and K. Mori. 2007. Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6alpha and XBP1. *Dev. Cell* **13**:365–376.
 59. Yamamoto, K., H. Yoshida, K. Kokame, R. J. Kaufman, and K. Mori. 2004. Differential contributions of ATF6 and XBP1 to the activation of endoplasmic reticulum stress-responsive cis-acting elements ERSE, UPRE and ERSE-II. *J. Biochem.* **136**:343–350.
 60. Yoshida, H., T. Matsui, A. Yamamoto, T. Okada, and K. Mori. 2001. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* **107**:881–891.