

2007

Identification and characterization of an RTX toxin in the emerging pathogen *Kingella kingae*

Thomas E. Kehl-Fie
Washington University School of Medicine in St. Louis
Joseph W. St. Geme III
Duke University Medical Center

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Kehl-Fie, Thomas E. and St. Geme, Joseph W. III, "Identification and characterization of an RTX toxin in the emerging pathogen *Kingella kingae*." *Journal of Bacteriology*. 189, 2. 430-436. (2007).
https://digitalcommons.wustl.edu/open_access_pubs/2487

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Identification and Characterization of an RTX Toxin in the Emerging Pathogen *Kingella kingae*

Thomas E. Kehl-Fie and Joseph W. St. Geme III
J. Bacteriol. 2007, 189(2):430. DOI: 10.1128/JB.01319-06.
Published Ahead of Print 10 November 2006.

Updated information and services can be found at:
<http://jb.asm.org/content/189/2/430>

	<i>These include:</i>
REFERENCES	This article cites 24 articles, 9 of which can be accessed free at: http://jb.asm.org/content/189/2/430#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Identification and Characterization of an RTX Toxin in the Emerging Pathogen *Kingella kingae*[∇]

Thomas E. Kehl-Fie^{1,2} and Joseph W. St. Geme III^{2*}

Edward Mallinckrodt Department of Pediatrics and Department of Molecular Microbiology, Washington University School of Medicine, 660 S. Euclid Ave., St. Louis, Missouri 63110,¹ and Departments of Pediatrics and Molecular Genetics and Microbiology, Duke University Medical Center, Children's Health Center, Durham, North Carolina 27710²

Received 18 August 2006/Accepted 30 October 2006

Kingella kingae is an emerging bacterial pathogen that is increasingly recognized as the causative agent of a variety of pediatric diseases, including septic arthritis and osteomyelitis. The pathogenesis of *K. kingae* disease is believed to begin with colonization of the upper respiratory tract. In the present study, we examined interactions between *K. kingae* and cultured respiratory epithelial cells and observed potent cytotoxicity, detected by both microscopy and lactic acid dehydrogenase (LDH) release assays. Experiments with synovial and macrophage cell lines revealed cytotoxicity for these cell types as well. Using mariner mutagenesis and a screen for loss of cytotoxicity, a genetic locus encoding an RTX toxin system was identified. Disruption of the *K. kingae* RTX locus resulted in a loss of cytotoxicity for respiratory epithelial, synovial, and macrophage cell lines. DNA sequence analysis demonstrated that the RTX locus is flanked by insertion elements and has a reduced G+C content compared to that of the whole genome. Two relatively less invasive *Kingella* species, *K. oralis* and *K. denitrificans*, were found to be noncytotoxic and to lack the RTX region, as determined by LDH release assays and Southern blotting. We concluded that *K. kingae* expresses an RTX toxin that has wide cellular specificity and was likely acquired horizontally. The possible roles for this toxin in the pathogenesis of *K. kingae* disease include breaching of the epithelial barrier and destruction of target tissues, such as synovium (joint lining).

Kingella kingae is a fastidious gram-negative bacterium and is a member of the *Neisseriaceae* family. Until recently, *K. kingae* was believed to be a rare pathogen and to exist primarily as a commensal organism in the upper respiratory tract. However, improvements in culture techniques and molecularly based detection methods have led to the recognition that *K. kingae* is an important cause of a variety of pediatric diseases, including osteomyelitis, septic arthritis, bacteremia, and endocarditis (8, 16, 19, 22, 25). In several recent studies, *K. kingae* has been identified as a leading cause of osteomyelitis and septic arthritis in children and the most common etiology of culture-negative cases of these infections in pediatric patients (8, 16, 19, 23).

Invasive disease due to *K. kingae* begins with asymptomatic colonization of the upper respiratory tract. One study demonstrated that more than 70% of young children are colonized with *K. kingae* at least once per year and that the same strain of *K. kingae* is able to persist in the respiratory tract of children for at least 2 months (22, 24, 25). In order to cause invasive disease, *K. kingae* must first breach the respiratory epithelium, which allows access to the underlying intravascular space and permits dissemination of the bacterium to distant sites, such as bones, joints, and endocardium. Infection at these sites is characterized by tissue destruction and inflammation.

To better understand the essential processes of respiratory tract colonization and breaching of the respiratory epithelium,

we examined *K. kingae* in assays with respiratory epithelial cell lines. We observed potent cytotoxicity for cultured respiratory epithelial cells. Microscopy and lactic acid dehydrogenase (LDH) release experiments revealed that *K. kingae* was also cytotoxic to macrophage-like cells and synovial cells, two cell types that the organism encounters in the joint (4). Using the mariner element and transposon mutagenesis, we identified a locus encoding an RTX toxin system flanked by insertion elements. Disruption of this locus resulted in loss of toxicity for all cell types tested. Further analyses suggested that the RTX locus was acquired via horizontal gene transfer.

MATERIALS AND METHODS

Bacterial strains, growth conditions, and plasmid construction. The strains used in this study are listed in Table 1. All strains were stored in either brain heart infusion (BHI) broth with 30% glycerol or Luria-Bertani (LB) broth with 15% glycerol at -80°C . *Escherichia coli* was routinely grown in LB broth or on LB agar plates supplemented with 30 $\mu\text{g}/\text{ml}$ chloramphenicol or 100 $\mu\text{g}/\text{ml}$ ampicillin as appropriate. *K. kingae* and *K. oralis* were grown on chocolate agar or TSA II blood agar plates (Becton-Dickson, Franklin Lakes NJ) at 37°C with 5% CO_2 . *K. denitrificans* was grown on TSA II blood agar plates. Chocolate agar was supplemented with 50 $\mu\text{g}/\text{ml}$ kanamycin as appropriate. Plasmid pRTXAC was constructed by first amplifying the *rtxAC* region from *K. kingae* strain 269-492 using primers GATCGGATCCTTACTGCGCTAGGTGCTAATACATTTT GCGG and GATCGAATTCATGGATAAATTTTCAGAACTAGGCAGCAT CGCG containing BamHI and EcoRI restriction sites, respectively, and then ligating the fragment into pTrc99A.

Cell lines. The cell lines used in this study were obtained from the American Tissue Culture Collection or the Duke Comprehensive Cancer Center cell culture facility and are listed in Table 2. Most cell lines were maintained at 37°C with 7.5% CO_2 ; the only exception was SW982 cells, which were maintained at 37°C without CO_2 .

Cytotoxicity assays. Bacteria were cultured on appropriate agar media overnight and were then resuspended in BHI broth to an optical density at 600 nm of 0.32. For microscopy studies, $\sim 3 \times 10^6$ bacteria were added to confluent cell

* Corresponding author. Mailing address: Department of Pediatrics, Duke University Medical Center, Children's Health Center, Room T901, DUMC 3352, Durham, NC 27710. Phone: (919) 681-4080. Fax: (919) 681-2714. E-mail: j.stgeme@duke.edu.

[∇] Published ahead of print on 10 November 2006.

TABLE 1. Strains and plasmids used in this study

Strain or plasmid	Description	Source or reference
Strains		
269-492	<i>K. kingae</i> clinical isolate	St. Louis Children's Hospital
ATCC 23330	<i>K. kingae</i> nasal isolate	ATCC
ATCC 23332	<i>K. kingae</i> blood isolate	ATCC
97-982	<i>K. kingae</i> clinical isolate	St. Louis Children's Hospital
05-001-1818	<i>K. kingae</i> clinical isolate	St. Louis Children's Hospital
KK01	Nonspreading/noncorroding derivative of 269-492	This study
KK03	Spreading/corroding derivative of 269-492	This study
60H11T1	Noncytotoxic transposon mutant of 269-492	This study
67A6T1	Noncytotoxic transposon mutant of 269-492	This study
97G1T1	Noncytotoxic transposon mutant of 269-492	This study
ATCC 33394	<i>K. denitrificans</i> type strain	ATCC
ATCC 51147	<i>K. oralis</i> type strain	ATCC
DH5α	<i>E. coli</i> F ⁻ φ80dlacZΔM15 Δ(lacZYA-argF)U169 deoR recA1 endA1 hsdR17(r _K ⁻ m _K ⁺) <i>phoA supE441 thi-1 gyrA96 relA1</i>	17
RTXAC	DH5α containing <i>rtxA</i> C in pRTXAC	This study
WAM716	DH5α containing <i>rtxB</i> D in pWAM716	5
RTXAC/WAM716	DH5α containing pRTXAC and pWAM716	This study
Plasmids		
pWAM716	<i>rtxB</i> and <i>rtxD</i> from <i>E. coli</i> in pACYC184	5
pRTXAC	<i>rtxA</i> and <i>rtxC</i> from <i>K. kingae</i> cloned into pTrc99A	This study

monolayers in 24-well tissue culture plates. To quantitate cytotoxicity, we used LDH release as a surrogate marker for cell death. For LDH release assays, ~7.5 × 10⁵ bacteria were added to confluent cell monolayers in 96-well tissue culture plates. Each sample was assayed in triplicate. For microscopy and LDH release assays, plates were centrifuged for 5 min at 1,000 rpm and then incubated for 10 min at 37°C. Samples were then either fixed and stained with Giemsa stain for examination by microscopy or incubated for 20 min at room temperature and assayed for LDH release. LDH release was assayed using a Cytotox One kit (Promega, Madison, WI) according to the manufacturer's instructions, using excitation and emission wavelengths of 540 nm and 590 nm, respectively. The maximal LDH release was defined as 100% and was determined by adding the lysis solution (Cytotox One kit) to uninfected monolayers, determining the absorbance, and then subtracting the background value. In complementation assays LDH release was assayed as described above, except that bacteria were grown overnight in LB medium at 37°C with the appropriate antibiotics, back diluted 1:10, and then grown for two additional hours at 37°C. In these assays ~2 × 10⁶ bacteria were added to each well.

Transposon library construction and screening. Chromosomal DNA was prepared from *K. kingae* strain 269-492 using a Wizard genomic DNA kit (Promega, Madison, WI) according to the manufacturer's instructions. To create a transposon library, chromosomal DNA from *K. kingae* 269-492 was mutagenized using the Himar1 transposase and pFalcon2, a plasmid that contains a mariner transposon derivative called Solo, which carries the *aphA3* kanamycin resistance gene (11). Mutagenesis was performed as described by Hendrixson et al. (11), except that the buffer exchanges described in the protocol were accomplished with a Qiagen II gel extraction kit (QIAGEN, Valencia CA) used according to the manufacturer's instructions instead of using DNA precipitation. To transform *K. kingae*, bacteria were grown for 12 h on chocolate agar and were resuspended to an optical density at 600 nm of 0.8 in BHI medium supplemented with 2% bovine

serum albumin and 0.5 mM CaCl₂. Aliquots of bacteria and mutagenized DNA were mixed in 24-well plates. The plates were incubated for 30 min at room temperature and were then supplemented with an equal volume of BHI medium plus 2% yeast extract and 4% horse plasma and incubated for 1 h at 37°C. Subsequently, the transformation reaction mixtures were plated onto chocolate agar containing kanamycin. In order to assess the randomness of transposon insertion, chromosomal DNA was extracted from individual transformants, digested with ClaI, and examined by Southern hybridization, using the *aphA3* cassette from pFalcon2 as a probe.

To identify mutants lacking cytotoxic activity, individual transformants were screened using LDH release assays. To confirm that the loss of cytotoxicity was due to the transposon insertion, chromosomal DNA was isolated from noncytotoxic mutants and retransformed into the parent *K. kingae* strain 269-492. The resulting transformants were then examined using LDH release assays. Following confirmation of the noncytotoxic phenotype, the sequence surrounding the transposon insertion site was determined using arbitrary PCR. The initial PCR was performed using arbitrary primer ARB1 (5' GGCCACGCGTCTGACTAGT ACNNNNNNNNNGATAT 3') or ARB6 (5' GGCCACGCGTCTGACTAGT ACNNNNNNNNNACGCC 3') and specific primer Solo5'Arb#1 (5' GCCC GGAATCATTTGAAGGTTG 3') or Solo3'Arb#1 (5' CGCGTCGCGACGC GTCAATTCGAGG 3'). Solo5'Arb#1 anneals at the 5' end of the Solo transposon, and Solo3'Arb#1 anneals at the 3' end of the Solo transposon. In the second round of amplification we utilized ARB2 (5' GGCCACGCGTCTGCA CTAGTAC 3'), which anneals to the 5' end of ARB 1 and ARB 6, and Solo5'outN (5' AATATGCATTTAATACTAGCGACGCC 3') or Solo3'outN (5' CGCTCTTGAAGGGAAGTATGTTG 3'), which are external to Solo5'Arb#1 and Solo3'Arb#1, respectively. The PCR products from the second round of amplification were gel purified and sequenced (Seqwright, Houston, TX) using either Solo5'outN or Solo3'outN, as appropriate.

Electron microscopy. Bacteria were allowed to adsorb onto Formvar/carbon-coated grids for 1 min and were then washed in distilled H₂O and stained with 1% aqueous uranyl acetate (Ted Pella Inc., Redding, CA) for 1 min. Excess liquid was gently wicked off, and the grids were allowed to air dry. Samples were viewed with a JEOL 1200EX transmission electron microscope (JEOL USA, Peabody, MA) at an accelerating voltage of 80 kV.

Southern analysis for *rtx* locus. Approximately 1 μg of chromosomal DNA from each strain was digested with BspHI, separated by agarose gel electrophoresis, and then transferred to a nitrocellulose membrane. The DNA probe was generated by PCR amplifying a 0.9-kb region containing *rtxC* and 0.3 kb of the 5' end of *rtxA* from *K. kingae* strain 269-469 using primers GCAGAAACG GCTACACCAGTTTGTAG and CAGAAGTACAGCAGCATCGCGTGG. The probe was labeled using the ECL direct nucleic acid labeling system (GE Healthcare, Piscataway, NJ) and was then incubated with the membrane at 42°C in blocking solution (ECL direct nucleic acid labeling system). Subsequently, the

TABLE 2. Cell lines used in this study

Cell line	Medium ^a	Description
Chang	MEM + 1× NEAA + 10% FCS	Human conjunctival cells
A549	MEM + 10% FCS	Human type II pneumocytes
HEp-2	DMEM + 10 % FCS	Human laryngeal cells
Hig-82	Ham's F12 + 10% FCS	Rabbit synoviocytes
RAW 264.7	DMEM + 10% FCS	Murine macrophage-like cells

^a Abbreviations: MEM, Eagle minimal essential medium; NEAA, nonessential amino acids; FCS, fetal calf serum; DMEM, Dulbecco's modified Eagle's medium.




FIG. 1. *K. kingae* cytotoxicity. (A) Light microscopy evidence of *K. kingae* cytotoxicity for Chang cells. The top panel shows an intact monolayer after incubation for 10 min with *E. coli* DH5 α , and the bottom panel shows a destroyed monolayer after incubation for 10 min with *K. kingae* strain 269-492. Bars = 100 μ M. (B and C) LDH release assays with *K. kingae* strains 269-492, 23330, 23332, 97-982, and 05-001-1818 and *E. coli* DH5 α with respiratory epithelial cells (B) and synovial and macrophage-like cells (C).

membrane was washed two times with 0.5 \times SSC and 0.4% (wt/vol) sodium dodecyl sulfate at 55°C for 10 min and then rinsed two times with 2 \times SSC at room temperature for 5 min (1 \times SSC is 0.15 M NaCl plus 0.015 M sodium citrate). Hybridization was detected using Supersignal West Pico (Pierce, Rockford, IL).

Nucleotide sequence accession number. The sequence corresponding to the *K. kingae* RTX locus has been deposited in the GenBank under accession number EF067866.

RESULTS

***K. kingae* is cytotoxic to a range of cell types.** To examine the interaction between *K. kingae* and respiratory epithelial cells, we inoculated wild-type strain 269-492 onto monolayers of Chang (human conjunctiva), HEp-2 (human larynx), and A549 (human type II pneumocyte) cells, incubated the preparations for various times, and then viewed them by light microscopy. *E. coli* DH5 α was included as a negative control. As shown in Fig. 1A, we observed that *K. kingae* induced rounding and detachment of cells from the monolayer. In an effort to confirm and quantitate this apparent cytotoxicity, we used an LDH release assay. As shown in Fig. 1B, incubation of *K. kingae* strain 269-492 and four additional *K. kingae* strains with Chang, A549, and HEp-2 cells for 10 min at 37°C resulted in 70% of the maximal LDH release.

To extend these observations and to study the cellular specificity of *K. kingae* cytotoxicity, we examined whether *K. kingae* was cytotoxic to Hig-82 rabbit synoviocytes and RAW 264.7 murine macrophage-like cells. All five *K. kingae* strains tested were highly cytotoxic to these cells, and the LDH release was more than 80% of the maximal LDH release (Fig. 1C). These results demonstrate that *K. kingae* expresses a potent cytotoxin with broad cell specificity, including human and nonhuman cells.

Cytotoxicity is independent of colony morphology. In previous work on *K. kingae* workers identified two interchanging colony morphologies, referred to as spreading/corroding and nonspreading/noncorroding, which are believed to reflect phase-variable expression of pili (7, 12, 13). The spreading/corroding morphology (SC type) correlates with expression of pili, and the nonspreading/noncorroding (N type) morphology correlates with reduced expression of pili (7, 12, 20). To determine if colony morphology and the density of pili influence *K. kingae* cytotoxicity, we recovered two morphological variants of strain 269-492, designated KK01 and KK03. As predicted, examination by negative staining transmission electron microscopy confirmed that KK01 has reduced numbers of pili and that KK03 expresses abundant pili (Fig. 2). Based on colony morphology, the phenotypes of the two variants appear to be stable, with more than 99% of colonies maintaining the parental morphology. As shown in Fig. 3, strains 269-492, KK01, and KK03 all displayed similar levels of cytotoxicity in assays with respiratory, synovial, and macrophage-like cells, suggesting that cytotoxicity is independent of colony morphology and the level of piliation.

***K. kingae* mutants 60H11T1, 69A6T1, and 97G1T1 lack cytotoxicity.** To identify the factors responsible for *K. kingae* cytotoxicity, we used the Himar1 transposase and pFalcon2 to generate a transposon library in strain 269-492. Assessment of 10 transformants by Southern hybridization confirmed that there was random insertion of the Solo transposon (data not shown). Using the LDH release assay, the library was screened for mutants that lacked cytotoxicity for Chang cells. Utilizing this approach, we recovered three independent noncytotoxic mutants, designated 60H11T1, 69A6T1, and 97G1T1 (Fig. 4A).




FIG. 2. Transmission electron micrographs of *K. kingae* strain 269-492 (top left), an N-type variant (bottom left) designated KK01, and an SC-type variant (top right) designated KK03 after negative staining with uranyl acetate. Bars = 100 nM.




FIG. 3. Cytotoxicity of spreading/corroding and nonspreading/non-corroding variants of *K. kingae*. *K. kingae* strains KK01, KK03, and 269-492 were assayed for cytotoxicity for respiratory epithelial cells (A) and synovial and macrophage-like cells (B) by the LDH release assay. *E. coli* DH5α was used as a negative control.

To identify the sites of transposon insertion in 60H11T1, 69A6T1, and 97G1T1, the transposon junctions were amplified using arbitrary PCR and were then sequenced. The resulting sequences were examined for homology to the results from the *K. kingae* sequencing project (T. E. Kehl-Fie et al., unpublished data). All three transposon insertions were located on an approximately 10-kb contig. In mutant 60H11T1, the transposon inserted into a gene encoding a protein with a high level of homology to MbxA, an RTX toxin from *Moraxella bovis*. In mutant 69A6T1, the transposon inserted immediately upstream of the predicted start site of the *mbxA* homolog. In mutant 97G1T1, the transposon inserted into a gene encoding a protein with a high level of homology to the *M. bovis* MbxD protein, a component of the type I secretion apparatus for export of MbxA. Further analysis of the contig revealed the presence of genes with homology to the genes encoding MbxC, MbxB, and TolC (Table 3 and Fig. 5), indicating that *K. kingae* possesses all of the genes necessary for production and secretion of an active RTX toxin (designated *rtxA*, *rtxB*, *rtxC*, *rtxD*, and *tolC*). To confirm that the RTX locus is responsible for the observed cytotoxicity, the *K. kingae* *rtxA* and *rtxC* genes (the structural and lipidating components of the locus) were cloned




FIG. 4. Loss of cytotoxicity of *K. kingae* mutants 60H11T1, 69A6T1, and 97G1T1. (A) LDH release assays with *K. kingae* strain 269-492 and mutants 60H11T1, 69A6T1, and 97G1T1 using Chang epithelial cells. *E. coli* DH5α was used as a negative control. (B) LDH release assays with *E. coli* strains DH5α, RTXAC, WAM716, and RTXAC/WAM716 using Chang epithelial cells.

into pTrc99A, creating pRTXAC. This construct was introduced into WAM716 (DH5α carrying a plasmid that contains *rtxB* and *rtxD*, encoding the *E. coli* type I secretion system), generating strain RTXAC/WAM716. As shown in Fig. 4B, RTXAC/WAM716 was found to be cytotoxic to Chang cells, providing strong evidence that the *K. kingae* locus is responsible for the observed *K. kingae* cytotoxicity.

To extend our characterization of mutants 60H11T1, 69A6T1, and 97G1T1, we examined these strains using cytotoxicity assays with respiratory epithelial, synovial, and macrophage-like cells and hemolysis assays on sheep blood agar plates. In assays with HEp-2 and A549 cells, mutants 60H11T1 and 97G1T1 were virtually devoid of cytotoxic activity and mutant 69A6T1 had markedly reduced cytotoxic activity (Fig. 6A). In assays with Hig-82 and RAW 264.7 cells, mutants 60H11T1 and 97G1T1 were found to be noncytotoxic, while

TABLE 3. *K. kingae* RTX locus homology

<i>K. kingae</i> gene product	Best homolog	% Identity ^a	% Similarity ^a
TolC	<i>N. meningitidis</i> TolC	64	82
RtxA	<i>M. bovis</i> MbxA	72	85
RtxC	<i>M. bovis</i> MbxC	73	87
RtxD	<i>N. meningitidis</i> RtxD	81	91
RtxB	<i>M. bovis</i> MbxB	83	91

^a Levels of identity and similarity were determined by comparing the entire *K. kingae* gene product and the entire sequence of the closest homolog.




FIG. 5. Diagram of the *K. kingae* RTX locus of strain 269-492, showing the locations of transposon insertions in mutants 60H11T1, 69A6T1, and 97G1T1.

mutant 69A6T1 remained highly cytotoxic, comparable to strain 269-492 (Fig. 6B). The observation that mutant 69A6T1 remained cytotoxic to Hig-82 and RAW 264.7 cells suggests that this strain produces reduced levels of functional toxin, a conclusion consistent with the finding that the insertion in this strain is immediately upstream of *rtxA*. When grown on sheep blood agar plates, 60H11T1 and 97G1T1 were found to be nonhemolytic and 69A6T1 was minimally hemolytic (data not shown).

To further explore the apparent increased *K. kingae* cytotoxicity for synovial and macrophage-like cells compared to the cytotoxicity for respiratory epithelial cells, we prepared serial dilutions of wild-type strain 269-492 and then measured LDH release. As shown in Fig. 7, Hig-82 and RAW 264.7 cells were found to be more sensitive than Chang and A549 cells, exhibiting similar levels of cytotoxicity with two- to fourfold fewer bacteria.

***K. kingae* RTX region appears to have been horizontally acquired.** The presence of all five genes necessary for production and secretion of an RTX toxin in a single gene cluster on the chromosome raised the possibility that the *K. kingae* RTX locus was acquired via horizontal gene transfer, similar to




FIG. 6. Cytotoxicity of *K. kingae* mutants 60H11T1, 69A6T1, and 97G1T1 for different cell types. LDH release assays were performed by using *K. kingae* strain 269-492 and mutants 60H11T1, 69A6T1, and 97G1T1 with respiratory epithelial cells (A) and synovial and macrophage-like cells (B). *E. coli* DH5α was used as a negative control.




FIG. 7. LDH release assays with *K. kingae* strain 269-492 comparing respiratory epithelial, synovial, and macrophage-like cells when different bacterial inocula were used.

speculation concerning other RTX toxins (6). To examine this hypothesis, we examined the flanking sequence and found that downstream of the RTX locus is a region with homology to insertion elements from *M. bovis*, *Neisseria meningitidis*, and *Haemophilus influenzae*. Further analysis revealed that the *K. kingae* RTX locus has a G+C content of 40%, which is significantly lower than the G+C content of the *K. kingae* genome (47%). To determine if the RTX locus is present in other *Kingella* species, *K. oralis* and *K. denitrificans* were examined to determine their cytotoxicity for respiratory, synovial, and macrophage-like cell lines using LDH release assays. Both *K. oralis* and *K. denitrificans* were found to be noncytotoxic to all cell types tested (Fig. 8). To assess whether this lack of cytotoxicity was due to inactivation or absence of the cytotoxin region, chromosomal DNA from *K. oralis* and *K. denitrificans* was examined by Southern hybridization using a 5' fragment of *rtxA* and *rtxC* as a probe. As expected, all five *K. kingae* strains examined possessed the cytotoxin region, while both *K. oralis* and *K. denitrificans* did not react with the probe (Fig. 9).

DISCUSSION

In this study we examined interactions between the emerging pediatric pathogen *K. kingae* and cultured respiratory epithelial cells. We observed that *K. kingae* expresses a potent cytotoxin capable of rapidly killing a variety of human respiratory epithelial cell lines. Additional investigation established that *K. kingae* is also cytotoxic to synovial cells and macrophage-like cells, two cell types that the bacterium is likely to




FIG. 8. LDH release assays examining *K. kingae* strain 269-492, *K. denitrificans* type strain, and *K. oralis* type strain cytotoxicity for respiratory epithelial cells, synovial cells, and macrophage-like cells. *E. coli* DH5α was used as a negative control.




FIG. 9. Southern analysis of chromosomal DNA from *K. kingae* (Kk) strains 269-492, 23330, 23332, 97-982, and 05-001-1817, the *K. denitrificans* type strain, the *K. oralis* type strain, and *E. coli* DH5α, using a 0.9-kb fragment containing *rtxC* and the 5' end of *rtxA* from *K. kingae* strain 269-492 as a probe.

encounter in the course of septic arthritis (4). By screening a *K. kingae* mariner transposon library for loss of cytotoxicity we identified a locus encoding an RTX toxin system. This locus is essential for cytotoxicity for respiratory epithelial, synovial, and macrophage-like cells.

RTX toxins have been divided into the following three categories based on cellular specificity: the hemolysins, which exhibit toxicity for a wide range of cell types (including erythrocytes); the cytotoxins, which exhibit toxicity for a wide but defined range of cell types; and the leukotoxins, which exhibit very narrow cell type and species specificity (21). The wide cellular specificity of the *K. kingae* RTX toxin suggests that this toxin belongs with the hemolysins. While the RTX hemolysins, cytotoxins, and leukotoxins are distinguished by different levels of specificity, all three classes possess several conserved motifs, including an amino-terminal hydrophobic domain implicated in pore formation, a species-specific lipid modification, a conserved calcium binding repeat domain, and a type I secretion signal (21). The *K. kingae* RTX toxin contains an amino-terminal hydrophobic domain, potential lipidation sites, and a calcium binding motif (data not shown). It is noteworthy that exposure of cells to sublytic doses of HlyA from *E. coli* or LktA from a variety of species results in a range of cellular reactions, including calcium fluxes, secretion of proinflammatory cytokines, and induction of apoptosis (2, 3, 10, 15, 18). Given the inflammatory nature of the disease caused by *K. kingae*, the possibility that the *K. kingae* cytotoxin may contribute to the inflammatory process is intriguing.

Examination of the *K. kingae* RTX locus revealed five genes, designated *rtxA*, *rtxC*, *rtxD*, *rtxB*, and *tolC*. Further analysis demonstrated that downstream of the locus is a region with homology to insertion elements from *M. bovis*, *Neisseria* species, and *Haemophilus* species. The identification of *tolC* within the locus is notable, since in most instances this gene is not physically associated with the toxin locus (the *M. bovis* RTX locus and the *Bordetella pertussis* adenylate cyclase locus are

exceptions) (1, 9). Although arranged in a noncanonical order, the *K. kingae* RTX locus exhibits marked homology with the *M. bovis* RTX locus. The *K. kingae* *rtxA*, *rtxC*, and *rtxB* genes encode proteins that have more than 70% identity with their *M. bovis* homologs (Table 3). The two remaining genes in the locus, *tolC* and *rtxD*, encode proteins with substantial homology to their *M. bovis* homologs but even greater homology to their *N. meningitidis* counterparts. These observations could be explained by acquisition of the RTX locus and then reacquisition of *tolC* and *rtxD* from either *K. kingae* itself or *N. meningitidis*. The results of the homology analysis combined with the observation that *K. denitrificans* and *K. oralis* lack an RTX locus suggest that the *K. kingae* RTX locus was acquired via horizontal gene transfer from either *M. bovis* or a common donor organism. The horizontal acquisition hypothesis is further supported by the identification of insertion elements flanking the *M. bovis* RTX locus (14).

In assays with synovial and macrophage-like cells, we observed two- to fourfold increases in sensitivity to *K. kingae* cytotoxic activity compared to the sensitivity observed with respiratory cells. This increased sensitivity is consistent with clinical observations that suggest that *K. kingae* colonizes the respiratory tract asymptotically but causes significant tissue destruction in the joint. It is possible that reduced sensitivity and other factors, such as mucus and other barriers, could combine to produce asymptomatic carriage of *K. kingae*.

Given the broad cellular specificity of the *K. kingae* RTX toxin, there are a number of possible roles for the cytotoxin during colonization of the respiratory tract and development of invasive disease. First, expression of the cytotoxin in the respiratory tract may result in damage to epithelial cells and tight junctions, disrupting innate physical defenses, allowing binding to respiratory mucosa and invasion of the bloodstream. Second, expression of the cytotoxin in the joint may promote inflammation directly by damaging the synovium or indirectly by promoting the release of proinflammatory cytokines, resulting in the recruitment of immune cells and inflammation. Third, the cytotoxin may aid *K. kingae* in immune evasion by killing macrophages and neutrophils.

In this paper we report the identification of a *K. kingae* RTX toxin that may play multiple roles in the pathogenesis of *K. kingae* disease, including colonization of the respiratory tract, invasion of the bloodstream, and damage to the joints. Given that septic arthritis is probably a terminal path for *K. kingae*, it is intriguing to speculate that the toxin may have been acquired and maintained to increase the area of the respiratory tract that *K. kingae* is capable of colonizing, coincidentally increasing the pathogenic potential. Further study of the *K. kingae* RTX toxin may increase our understanding of the balance between commensal and pathogen that is common among microbial pathogens, including other members of the *Neisseriaceae* family.

ACKNOWLEDGMENTS

We thank Shane Cotter for thoughtful discussions. Additionally, we thank Wandy Beatty for assistance with electron microscopy, Rodney Welch for providing strain WAM716, and Pablo Yagupsky for offering general advice about handling *K. kingae*.

This work was supported by NIH training grant T32-GM07067 to T.K.F.

REFERENCES

- Angelos, J. A., J. F. Hess, and L. W. George. 2003. An RTX operon in hemolytic *Moraxella bovis* is absent from nonhemolytic strains. *Vet. Microbiol.* **92**:363–377.
- Atapattu, D. N., and C. J. Czuprynski. 2005. *Mannheimia haemolytica* leukotoxin induces apoptosis of bovine lymphoblastoid cells (BL-3) via a caspase-9-dependent mitochondrial pathway. *Infect. Immun.* **73**:5504–5513.
- Bhakdi, S., and E. Martin. 1991. Superoxide generation by human neutrophils induced by low doses of *Escherichia coli* hemolysin. *Infect. Immun.* **59**:2955–2962.
- Bremell, T., A. Abdelnour, and A. Tarkowski. 1992. Histopathological and serological progression of experimental *Staphylococcus aureus* arthritis. *Infect. Immun.* **60**:2976–2985.
- Felmlee, T., and R. A. Welch. 1988. Alterations of amino acid repeats in the *Escherichia coli* hemolysin affect cytolytic activity and secretion. *Proc. Natl. Acad. Sci. USA* **85**:5269–5273.
- Frey, J., and P. Kuhnert. 2002. RTX toxins in *Pasteurellaceae*. *Int. J. Med. Microbiol.* **292**:149–158.
- Froholm, L. O., and K. Bovre. 1972. Fimbriation associated with the spreading-corroding colony type in *Moraxella kingii*. *Acta Pathol. Microbiol. Scand. B* **80**:641–648.
- Gene, A., J. J. Garcia-Garcia, P. Sala, M. Sierra, and R. Huguet. 2004. Enhanced culture detection of *Kingella kingae*, a pathogen of increasing clinical importance in pediatrics. *Pediatr. Infect. Dis. J.* **23**:886–888.
- Glaser, P., H. Sakamoto, J. Bellalou, A. Ullmann, and A. Danchin. 1988. Secretion of cyclolysin, the calmodulin-sensitive adenylate cyclase-haemolysin bifunctional protein of *Bordetella pertussis*. *EMBO J.* **7**:3997–4004.
- Gleason, T. G., C. W. Houlgrave, A. K. May, T. D. Crabtree, R. G. Sawyer, W. Denham, J. G. Norman, and T. L. Pruett. 1998. Hemolytically active (acylated) alpha-hemolysin elicits interleukin-1beta (IL-1beta) but augments the lethality of *Escherichia coli* by an IL-1- and tumor necrosis factor-independent mechanism. *Infect. Immun.* **66**:4215–4221.
- Hendrixson, D. R., B. J. Akerley, and V. J. DiRita. 2001. Transposon mutagenesis of *Campylobacter jejuni* identifies a bipartite energy taxis system required for motility. *Mol. Microbiol.* **40**:214–224.
- Henrichsen, J., L. O. Froholm, and K. Bovre. 1972. Studies on bacterial surface translocation. 2. Correlation of twitching motility and fimbriation in colony variants of *Moraxella nonliquefaciens*, *M. bovis*, and *M. kingii*. *Acta Pathol. Microbiol. Scand. B* **80**:445–452.
- Henrikson, S. D. 1969. Corroding bacteria from the respiratory tract. 1. *Moraxella kingii*. *Acta Pathol. Microbiol. Scand.* **75**:85–90.
- Hess, J. F., and J. A. Angelos. 2006. The *Moraxella bovis* RTX toxin locus *mbx* defines a pathogenicity island. *J. Med. Microbiol.* **55**:443–449.
- Ludwig, A., and W. Goebel. 2000. Dangerous signals from *E. coli* toxin. *Nat. Med.* **6**:741–742.
- Moumle, K., J. Merckx, C. Glorion, P. Berche, and A. Ferroni. 2003. Osteoarticular infections caused by *Kingella kingae* in children: contribution of polymerase chain reaction to the microbiologic diagnosis. *Pediatr. Infect. Dis. J.* **22**:837–839.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Uhlen, P., A. Laestadius, T. Jahnukainen, T. Soderblom, F. Backhed, G. Celsi, H. Brismar, S. Normark, A. Aperia, and A. Richter-Dahlfors. 2000. Alpha-haemolysin of uropathogenic *E. coli* induces Ca²⁺ oscillations in renal epithelial cells. *Nature* **405**:694–697.
- Verdier, I., A. Gayet-Ageron, C. Ploton, P. Taylor, Y. Benito, A. M. Freydiere, F. Chotel, J. Berard, P. Vanhems, and F. Vandenesch. 2005. Contribution of a broad range polymerase chain reaction to the diagnosis of osteoarticular infections caused by *Kingella kingae*: description of twenty-four recent pediatric diagnoses. *Pediatr. Infect. Dis. J.* **24**:692–696.
- Weir, S., and C. F. Marrs. 1992. Identification of type 4 pili in *Kingella denitrificans*. *Infect. Immun.* **60**:3437–3441.
- Welch, R. A. 2001. RTX toxin structure and function: a story of numerous anomalies and few analogies in toxin biology. *Curr. Top. Microbiol. Immunol.* **257**:85–111.
- Yagupsky, P. 2004. *Kingella kingae*: from medical rarity to an emerging paediatric pathogen. *Lancet Infect. Dis.* **4**:358–367.
- Yagupsky, P., R. Dagan, C. W. Howard, M. Einhorn, I. Kassis, and A. Simu. 1992. High prevalence of *Kingella kingae* in joint fluid from children with septic arthritis revealed by the BACTEC blood culture system. *J. Clin. Microbiol.* **30**:1278–1281.
- Yagupsky, P., R. Dagan, F. Prajrod, and M. Merires. 1995. Respiratory carriage of *Kingella kingae* among healthy children. *Pediatr. Infect. Dis. J.* **14**:673–678.
- Yagupsky, P., N. Peled, and O. Katz. 2002. Epidemiological features of invasive *Kingella kingae* infections and respiratory carriage of the organism. *J. Clin. Microbiol.* **40**:4180–4184.