

2004

Sequence organization and insertion specificity of the novel chimeric ISHp609 transposable element of *Helicobacter pylori*

Dangeruta Kersulyte

Washington University School of Medicine in St. Louis

Awdesh Kalia

Washington University School of Medicine in St. Louis

MaoJun Zhang

Washington University School of Medicine in St. Louis

Hae-Kyung Lee

Washington University School of Medicine in St. Louis

Dharmalingam Subramaniam

Washington University School of Medicine in St. Louis

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Kersulyte, Dangeruta; Kalia, Awdesh; Zhang, MaoJun; Lee, Hae-Kyung; Subramaniam, Dharmalingam; Kiuduliene, Levute; and Chalkauskas, Henrika, "Sequence organization and insertion specificity of the novel chimeric ISHp609 transposable element of *Helicobacter pylori*." *Journal of Bacteriology*.186,22. 7521-7528. (2004).
https://digitalcommons.wustl.edu/open_access_pubs/2715

Authors

Dangeruta Kersulyte, Awdesh Kalia, MaoJun Zhang, Hae-Kyung Lee, Dharmalingam Subramaniam, Levute Kiuduliene, and Henrika Chalkauskas

Sequence Organization and Insertion Specificity of the Novel Chimeric IS *Hp609* Transposable Element of *Helicobacter pylori*

Dangeruta Kersulyte, Awdhesh Kalia, MaoJun Zhang,
Hae-Kyung Lee, Dharmalingam Subramaniam, Levute
Kiuduliene, Henrikas Chalkauskas and Douglas E. Berg
J. Bacteriol. 2004, 186(22):7521. DOI:
10.1128/JB.186.22.7521-7528.2004.

Updated information and services can be found at:
<http://jb.asm.org/content/186/22/7521>

These include:

SUPPLEMENTAL MATERIAL

[Supplemental material](#)

REFERENCES

This article cites 25 articles, 17 of which can be accessed free
at: <http://jb.asm.org/content/186/22/7521#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Sequence Organization and Insertion Specificity of the Novel Chimeric ISHp609 Transposable Element of *Helicobacter pylori*†

Dangeruta Kersulyte,¹ Awdhesh Kalia,¹ MaoJun Zhang,¹ Hae-Kyung Lee,^{1,2}
Dharmalingam Subramaniam,³ Levute Kiuduliene,⁴
Henrikas Chalkauskas,⁵ and Douglas E. Berg^{1,3*}

Department of Molecular Microbiology¹ and Department of Internal Medicine,³ Washington University School of Medicine, St. Louis, Missouri; Department of Clinical Pathology, St. Mary's Hospital, Catholic University Medical College, Uijungbu, Korea²; and Institute of Biotechnology⁴ and Clinic of Gastroenterology, Vilnius University Hospital,⁵ Vilnius, Lithuania

Received 2 July 2004/Accepted 29 July 2004

Here we describe ISHp609 of *Helicobacter pylori*, a new member of the IS605 mobile element family that is novel and contains two genes whose functions are unknown, *jhp960* and *jhp961*, in addition to homologs of two other *H. pylori* insertion sequence (IS) element genes, *orfA*, which encodes a putative serine recombinase-transposase, and *orfB*, whose homologs in other species are also often annotated as genes that encode transposases. The complete four-gene element was found in 10 to 40% of strains obtained from Africa, India, Europe, and the Americas but in only 1% of East Asian strains. Sequence comparison of 10 representative ISHp609 elements revealed higher levels of DNA sequence matches (99%) than those seen in normal chromosomal genes (88 to 98%) or in other IS elements (95 to 97% for IS605, IS606, and IS607) from the same *H. pylori* populations. Sequence analysis suggested that ISHp609 can insert at many genomic sites with its left end preferentially next to TAT, with no target specificity for its right end, and without duplicating or deleting target sequences. A deleted form of ISHp609, containing just *jhp960* and *jhp961* and 37 bp of *orfA*, found in reference strain J99, was at the same chromosomal site in 15 to 40% of the strains from many geographic regions but again in only 1% of the East Asian strains. The abundance and sequence homogeneity of ISHp609 and of this nonmobile remnant suggested a recent bottleneck and then rapid spread in *H. pylori* populations, possibly selected by the contributions of the elements to bacterial fitness.

Insertion sequence (IS) elements are a diverse group of specialized DNA segments that move to new sites in prokaryotic and eukaryotic genomes by mechanisms that do not require extensive DNA sequence homology (for reviews see references 5, 7, and 19). They cause insertion mutations and genome rearrangements, affect nearby gene expression, and help mediate the spread of resistance and virulence determinants within and among species. Many IS elements specify just a single transposase protein that acts in concert with host proteins at each end of the element to mediate insertion. Other, more complex elements, such as Tn7, specify two or more proteins that act together as the transposase, plus additional proteins that help select insertion sites or affect the efficiency of transposition. Host proteins can also affect the frequency or specificity of transposition of certain elements. Many, but not all, species of elements terminate in short inverted repeats (size range, 9 to 40 bp) and generate short direct repeats of target sequences (typically 2 to 9 bp) when they transpose.

Each of the four known species of IS elements in *H. pylori* (IS605, IS606, IS607, and ISHp608) belongs to the distinctive IS605 mobile element family, and each seems to be chimeric, containing two transposition-related genes, *orfA* and *orfB*, that

may have different phylogenetic origins (12, 14, 16). The IS605 element family is divisible into two subfamilies based on *orfA* homologies; in one subfamily, represented by IS607, *orfA* encodes a putative serine recombinase that helps IS607 transpose to multiple sites (in contrast, most serine recombinases mediate site-specific recombination), and in the other subfamily, represented by IS605, IS606, and ISHp608 (28 to 36% amino acid identity for sequences encoded by *orfA* genes), *orfA* encodes a transposase related to that encoded by IS200, a single-gene element that is widespread in enteric species (5, 19, 25) whose product is distinct from serine recombinase proteins. The proteins encoded by *orfB* genes of IS605 family members exhibit 25 to 35% amino acid identity to one another; their homologs in other species are often annotated in sequence databases as putative transposases, and they are also homologs of the protein encoded by *gipA*, a *Salmonella* prophage gene that enhances bacterial growth in Peyer's patches (22).

IS605, IS607, and ISHp608 each have been found to transpose in *Escherichia coli* (12, 14, 16). With each of the two elements tested (IS607 and ISHp608), transposition depended on *orfA* and not on *orfB*. Hence, the constant presence of *orfB* in IS605 family members suggested either involvement in transposition in certain species or a contribution to bacterial fitness (11a). Inspection of sequences at sites of insertion in *H. pylori* and *E. coli* indicated that (i) IS605, IS607, and ISHp608 insert with their left (*orfA*) ends immediately downstream of specific AT-rich sequences (5'-TTTAA or 5'-TTTAAC for IS605, 5'-TTTAT for IS606, and 5'-TTAC for ISHp608), and their right (*orfB*) ends seem to join to target

* Corresponding author. Mailing address: Box 8230, Department of Molecular Microbiology, 4940 Parkview Place, Washington University School of Medicine, St. Louis, MO 63110. Phone: (314) 362-2772. Fax: (314) 362-1232. E-mail: berg@borcim.wustl.edu.

† Supplemental material for this article may be found at <http://jlb.asm.org>.

TABLE 1. Primers^a

Primer	Sequence	Location
ISHp609 primers		
609LE	CATTAAACTTTCAATTTAATTTTG	At the left end, forward
609R-t1	CTAAAAACTCTTTAATCTTATTTAAAGTA	At the right end, reverse
609.F1	CACAACAGGTATTAATGCTT	1,083 bp from the right end, in <i>orfB</i> , forward
609.R1	CTTTAGCTCTTGTTTCCAGC	1,985 bp from the left end, in <i>orfB</i> , reverse
609t2-1 ^b	TTACCTGCAAGCCATTAAGG	1,037 bp from the right end, in <i>orfB</i> , forward
609t2-2 ^b	TTATGCATAAAGTTATTCAGC	1,535 bp from the left end, in <i>orfB</i> , reverse
FlankL	CTTTATTTGGGAGATTTAGAAGC	2,266 bp from the right end, in <i>orf1</i> , forward
609.R6	CCATTCTGCAATAATAGCTTCTC	346 bp from the left end, in <i>orf2</i> , reverse
609R3t1	TAAATAGCTTTTCTACTAACTCATA	1,001 bp from the left end, in <i>orfA</i> , reverse
609.R5	ATTAGCGTAGTTGTAAAGGTT	1,199 bp from the left end, in <i>orfB</i> , reverse
Other primers		
jhp959	GTAGCACAACTCTTATGCGATG	177 bp from the 3' end of <i>jhp959</i> , forward
jhp962	TGTATGTCATGCTGAGCGAAAAC	371 bp from the 3' end of <i>jhp962</i> , reverse
CR2-LF	CATACTAGACTTAAAGGACAGCA	Left of ISHp609 in CR2 clone ^c
CR2-RF	ATTTTGATTTTGTCTGTTACTTCGCAC	Right of ISHp609 in CR2 clone ^c
B43-LF	GCTATGGCAGAAAACATCTAT	Left of ISHp609 in HUP-B43
B43-RF	GAGTTTCTAAAATAATTCCTCAT	Right of ISHp609 in HUP-B43
B79-LF	CAGATTTACCCAACTCACT	Left of ISHp609 in HUP-B79
B79-RF	AAGGTTTTATCAAAGCCTATGC	Right of ISHp609 in HUP-B79
LitA7-LF	GAGGTAATTTTTGAAATTTTAAACAC	Left of ISHp609 in LitA7
LitA7-RF	ATCAGGCTTTAGGGTATTCTTT	Right of ISHp609 in LitA7
LitA38-LF	TTTATCCTTTCTTTATTATTTAACTTTT	Left of ISHp609 in LitA38
LitA38-RF	ATTTTATGCGATTGCATTGAAAAG	Right of ISHp609 in LitA38

^a Sequences of additional primers used for sequencing are available on request.

^b ISHp609^{var} specific (Chennai4 strain).

^c See reference 6.

DNAs nonspecifically; (ii) in contrast, IS607 whose *orfA* transposase gene is unrelated to the other genes, inserted preferentially between adjacent GG nucleotides in target DNA; (iii) none of these elements duplicated or deleted sequences at sites of insertion; and (iv) none contained terminal inverted repeats (12, 14, 16). The *H. pylori* ISHp609 element described here is nonrandomly distributed geographically and is unique among elements of the IS605 family in containing four open reading frames instead of the usual two.

MATERIALS AND METHODS

General methods. Standard procedures were used for *H. pylori* growth on brain heart infusion agar (Difco) containing 10% horse blood in a microaerobic atmosphere (27). High-molecular-weight genomic DNA was isolated by a hexadecyltrimethylammonium method (3).

Specific PCR were carried out in 20- μ l mixtures containing 5 to 10 ng of DNA, 0.1 U of *Taq* polymerase (Biolase; Midwest Scientific, St. Louis, Mo.) or the Expand High Fidelity *Taq-Pwo* polymerase mixture (Boehringer-Mannheim, Indianapolis, Ind.), 2.5 pmol of each primer, and each deoxynucleoside triphosphate at a concentration of 0.2 mM in a standard buffer for 30 cycles with the following cycling parameters: denaturation at 94°C for 30 s, annealing at a temperature appropriate for the primer sequence (generally 50°C) for 30 s, and DNA synthesis at 72°C for an appropriate time (1 min per kb). PCR primers are listed in Table 1.

DNA sequencing was carried out by using a Big Dye Terminator DNA sequencing kit (Perkin-Elmer) and an ABI automated sequencer. Direct sequencing on chromosomal DNA was done with 5 μ l of chromosomal DNA (1 to 2 μ g), 1 μ l of primer (10 pmol per μ l), and 6 μ l of Big Dye under the following conditions: 96°C for 5 min and then 90 cycles of denaturation at 96°C for 10 s, annealing at a temperature appropriate for the primer for 5 s, and extension at 60°C for 4 min under oil-free conditions (Perkin-Elmer 2400).

DNA sequence editing and analysis were performed with programs in the GCG package (Genetics Computer Group, Madison, Wis.), programs and data in the *H. pylori* genome sequence databases (2, 24), and BLAST and Pfam (version 14.0) homology search programs (<http://www.ncbi.nlm.nih.gov/BLAST/BLAST.cgi>; <http://pfam.wustl.edu/hmmsearch.shtml>).

Phylogenetic analysis. *H. pylori* OrfA and OrfB IS element protein sequences were aligned with CLUSTALX by using a PAM250 amino acid substitution

matrix. OrfA and OrfB neighbor-joining phylogenies were constructed by using the Jones-Taylor-Thornton distance matrix and variable rates among sites (modeled with a gamma distribution shape parameter, $\alpha = 0.5$). Gaps in the protein alignments were deleted in pairwise comparisons. Phylogenetic analysis and DNA diversity calculation were done with Mega2.1 (www.megasoftware.net).

Bacterial strains and plasmids. Most *H. pylori* strains used were obtained from the Berg laboratory collection and have been described in detail elsewhere (4, 13, 15, 16, 20). The *H. pylori* strains used in this study included 71 Peruvian strains, 24 Spanish strains, 47 Lithuanian strains from Vilnius, 28 African strains (16 strains from Soweto in South Africa and 12 strains from Gambia), 69 Indian strains (48 strains from Calcutta, 15 strains from Chennai, and 6 strains from the Santal tribe), 71 Japanese strains (24 strains from Ube and 47 strains from Fukui), 46 strains from Alaskan natives, 47 Chinese strains (40 strains from YunNan and 7 strains from Changle), and 47 Korean strains.

Plasmids carrying an intact ISHp609 element marked with chloramphenicol resistance downstream of the *orfB* stop codon were constructed, and transposition assays in *E. coli* were carried out as described previously for ISHp608 (16).

GenBank accession numbers of DNA sequences. The sequence of ISHp609 from strain HUP-B43 corresponds to nucleotides 14337 through 16733 of the 33,671-nucleotide sequence of the plasticity zone in GenBank accession number AY487825. The other three full-length ISHp609 elements were accessioned in GenBank as follows: from strain HUP-B79, nucleotides 287 through 2684 of the 3,084 nucleotides in accession number AY639112; from strain LitA7, nucleotides 381 through 2794 of the 2,875 nucleotides in accession number AY639110; and from strain LitA38, nucleotides 20 through 2417 of the 2,603 nucleotides in accession number AY639111. The full length of the rare type of ISHp609 element (ISHp609^{var}) from strain Chen4 corresponds to nucleotides 313 through 2346 of the 2,496 nucleotides in accession number AY639119. The GenBank accession numbers for partial ISHp609 sequences (all but 24 nucleotides on the left and 34 nucleotides on the right due to the positions of PCR binding sites) were as follows: strain HUP-B80, AY639113; Alaska64, AY639115; Alaska97, AY639116; AfricaR48, AY639114; I-86, AY639117; and SJM27, AY639118.

RESULTS

ISHp609 discovery and sequence. The 2.4-kb ISHp609 element was discovered while the plasticity zone of strain HUP-B43 (accession number AY487825) was being sequenced as an insertion in gene *jhp928* (the gene designations used match




FIG. 1. Structures of ISHp609 and related elements. (A) Full-length predominant ISHp609 type with four characteristic open reading frames and 99% DNA identity, independent of geographic origin. Sequence analysis of 10 full-length elements from Spanish strains (HUP-B43, HUP-B79, and HUP-B80; accession numbers AY487825, AY639112, and AY639113), Lithuanian strains (Lit-7 and Lit38; accession numbers AY639110 and AY639111), Alaskan strains (Al64 and Al97; accession numbers AY639115 and AY639116), a Peruvian strain (SJM27; accession number AY639118), an Indian strain (I-86; accession number AY639117), and an African strain (R48; accession number AY639114) showed limited internal divergence. In strain Lit7 *orfA* contained a frameshift due to a 14-bp duplication; in strain HUP-B79 *orfA* contained an in-frame stop codon due to a G-to-T substitution; and in strain I-86 *orf1* contained an in-frame stop codon due to a C-to-T substitution. (B) ISHp609var. This rare variant element was found in one Indian strain (Chennai4; accession number AY639119) with 81% DNA identity to the predominant type. ISHp609var has full-length *orfA* and *orfB* genes, although *orfB* is probably inactive due to a frameshift. It lacks *orf2* and most of *orf1*, and it has the first 79 bp of *orf1*, but there is no start codon at its left end. (C) *orf1*-2 remnant. This DNA segment consists of *orf1* (*jhp960* in strain J99), *orf2* (*jhp961* in strain J99), and the first 37 bp of *orfA* and is located between homologs of *jhp959* and *jhp962* in most or all strains (as determined by PCR with primers *jhp959* and *jhp962*, primers *jhp959* and 609.R6, and primers FlankL and *jhp962*).

those of reference strains J99 [*jhp*] and 26695 [*hp*] when homologies are high [2, 24]). It contained four open reading frames designated *orf1*, *orf2*, *orfA*, and *orfB* (Fig. 1A).

The ISHp609 *orfA* sequence (210 codons) is homologous to the *orfA* sequence of IS607 (32% protein identity and 58% protein similarity) (Fig. 2A), and its 5' end specifies a helix-turn-helix domain (conserved domain cd01104, HTH_MlrA; E value, $3e^{-04}$) that might mediate DNA binding. Extrapolation from IS607 studies (14) suggested that *orfA* might encode a transposase. The ISHp609 *orfA* product also belongs to a widespread protein family whose members are generally designated serine recombinases or site-specific integrase-resolvases. One of the strongest homologies is with a putative serine recombinase gene of *Thermoanaerobacter tengcongensis* MB4T (*tte0714*; 47% protein identity and 68% protein similarity).

The ISHp609 *orfB* sequence (409 codons) exhibits weak homology to the IS605 *orfB* sequence (22% protein identity and 42% protein similarity) (Fig. 2B) and more generally to members of a widespread gene family that are annotated as genes that encode transposases but that also include the *gipA* virulence gene of *Salmonella*. Again, one of the strongest homologies is with an open reading frame in *T. tengcongensis* MB4T (*tte0715*; 42% protein identity and 58% protein similarity), which is adjacent to the *orfA* homolog, *tte0714*. The C-terminal tetracysteine Zn(II) binding motif CX₍₂₎CX₍₁₅₎CX₍₂₎C (C4-type zinc finger) found in *GipA* (22) and the OrfB proteins encoded in IS605, IS606, IS607, and ISHp608 was absent from

the ISHp609 OrfB protein and also from its close homolog in *T. tengcongensis* (Fig. 2C).

Two short coding sequences in ISHp609, *orf1* (85 codons) and *orf2* (61 codons), closely matched two genes of reference strain J99 whose functions are unknown, *jhp960* (99% DNA identity and 100% protein identity) and *jhp961* (98% DNA identity, 96% protein identity, and 98% protein similarity). These open reading frames are located between *jhp959* and *jhp962* in strain J99, not within *jhp928*, but they are absent from the other fully sequenced genome (strain 26695) (24). Although annotated as a gene whose function is unknown, *orf1* belongs to a gene family whose protein products generally contain a characteristic ligand binding domain next to a helix-turn-helix DNA binding domain (Pfam domain 03681, UPF0150; 14 to 68 bp; E value, $1e^{-10}$), and the protein encoded by *orf2* exhibits a conserved predicted periplasmic or secreted lipoprotein motif throughout its length (61 amino acids; cluster of orthologous groups, COG1724; E value, $6e^{-13}$). Small genes related to *orf1* and *orf2* are found in many bacterial species, in some cases together (for example, *ssr1765* and *ssr1766* in *Synechocystis* sp. strain PCC 6803 [GenBank accession numbers BAA16930 and BAA16931], with 46% amino acid identity and 70% amino acid similarity to *orf1* and 35% amino acid identity and 38% amino acid similarity to *orf2*, respectively). However, most homologs of either *orf1* or *orf2* occur singly and are not linked to a homolog of the other gene in other bacterial species. No *orfA* or *orfB* homologs were found next to *ssr1765* and *ssr1766* in *Synechocystis*, nor were *orf1* or *orf2* homologs found next to the *orfA* and *orfB* homologs (*tte0714* and *tte0715*) in *T. tengcongensis*.

ISHp609 elements and adjacent DNAs from three additional strains (strains HUP-B79, LitA7, and LitA38) were sequenced by primer walking on genomic DNAs in order to better understand the structure of the element and its insertion specificity. All three elements were 2.4 kb long, contained four open reading frames (*orf1*, *orf2*, *orfA*, and *orfB*) matching the open reading frames found in strain HUP-B43, and were 99% identical in DNA sequence to one another. A comparison of sequences flanking ISHp609 with corresponding empty sites in the 26695 and/or J99 genome showed that the element was located at a different genetic locus in each strain (Fig. 3A). These results, coupled with PCR data (see below), indicate that ISHp609 is transposable and that it can insert into many genomic sites. Sequence comparisons also identified the probable termini of the element and its insertion specificity. Based on data summarized in Fig. 3A, the ends of ISHp609 seemed to be 5'-TAT or CAT on the left and 5'-CAT on the right. The left end was inserted preferentially next to 5'-TAT, whereas there seemed to be no target specificity for the right end, and there was no evidence of target sequence duplication or deletion during insertion.

Further IS element sequencing was done by using nearly full-length ISHp609 PCR products from six additional strains from five geographic regions (Spain, Africa, Alaska, India, and Peru) and with primers located just within the ISHp609 ends (primers 609LE and 609R-t1) (GenBank accession numbers AY636113 to AY636118). Each element had the same number and organization of open reading frames, and each was $\geq 99\%$ identical in DNA sequence to other elements, regardless of the geographic origin. The ratio of synonymous to




FIG. 2. Phylogenetic relationships among *H. pylori* IS elements. (A) OrfA and homologs. Two subfamilies were identified, one represented by OrfAs of IS605 (accession number NP_208326), IS606 (accession number AAD11513), and ISHp608 (accession number AAL06576), which are not considered to encode serine recombinases based on amino acid homologies, and the other represented by IS607 (accession number AAF05600), ISHp609 (accession number AAR83266.1), ISHp609var (accession number AY639119), and the closest homolog in *T. tengcongensis* (*T. teng.*) (*tte0714*; accession number AAM23976), which are thought to encode serine recombinases. Branches with significant bootstrap support (≥ 50) are indicated. Bar = 1 amino acid substitution per site. (B) OrfB and homologs. OrfBs of *H. pylori* IS605 (accession number NP_208324), IS606 (accession number AAD11514), IS607 (accession number AAF05601), ISHp608 (accession number AAL06577), and *Salmonella*'s GipA protein (accession number NP_752781) form an OrfB subfamily different from that of ISHp609 (accession number AAR83267.1), ISHp609var (accession number AY639119), and the closest homolog in *T. tengcongensis* (*tte0715*; accession number AAM23977). (C) Partial C-terminal sequence alignment of *H. pylori* IS element OrfBs, GipA (22), and the corresponding sequence in *T. tengcongensis*. A single C-terminal Zn(II) binding tetracysteine motif, CX₍₂₎CX₍₁₅₎CX₍₂₎C (C4-type zinc finger), is well conserved among IS605, IS606, IS607, and ISHp608 OrfBs and GipA and might potentially facilitate DNA or RNA binding or protein-protein interaction. Notably, this motif is not present in ISHp609 OrfB (both predominant and variant).

nonsynonymous changes in the four open reading frames ranged from 3.2 to 7.0 (average, 5.0) (Table 2), values that are typical of *H. pylori* housekeeping genes (1, 9).

In addition to this predominant ISHp609 type, we also found one variant element (strain Chennai4) based on weaker-than-normal PCR amplification with *orfB*-specific primers 609.F1 and 609.R1. This element, ISHp609var, exhibited only 81% DNA homology to other ISHp609 elements and consisted of the 5' end of *orf1* (79 bp, 79% DNA homology) without an obvious start codon, linked directly to full-length *orfA* and *orfB* (although *orfB* contains a frameshift due to a 5-bp internal deletion) (Fig. 1B). ISHp609var was inserted into an unknown chromosomal site with its inferred left end next to a TAT sequence, which suggested insertion specificity matching that of the predominant ISHp609 type (Fig. 3B).

In further experiments, a plasmid clone of ISHp609 from strain HUP-B43, marked with a chloramphenicol resistance determinant, was used to select for transposition to the F factor pOX38 in *E. coli* in a standard mating-out assay, essentially as done previously with IS607 and ISHp608 (14, 16).

However, no transposition ($<10^{-9}$) of this marked ISHp609 element was detected in any of four repetitions of this assay. In contrast, IS607 and ISHp608 transposed at frequencies of about 10^{-7} in equivalent assays (14, 16).

ISHp609 geographic distribution and structural analysis. The frequency of ISHp609 carriage in various *H. pylori* populations was estimated by PCR with 479 strains by using two *orfB*-specific primers (primers 609.F1 and 609.R1). ISHp609 was found in 35 to 40% of the strains from Europe (Spain and Lithuania), in 20 to 40% of the strains from the Americas (Peru, Guatemala, and Alaska), and in 10 to 15% of the strains from India and Africa but in only 1% of the strains from East Asia (Table 3). All 479 strains, including 68 additional strains from India, were tested for the variant type (ISHp609var) that had been found in one Indian strain by using specific primers (primers 609t2-1 and 609t2-2). No additional strains harboring this variant were found. This seemed to be reminiscent of the single case of a rare type 3 ISHp608 element found in one Indian strain but not in 116 other Indian strains or in 606 strains from other regions (16).




FIG. 3. Terminal sequences of ISHp609, ISHp609var, and the *orf1-2* remnant and their sites of insertion in *H. pylori*. ISHp609, ISHp609var, and *orf1-2* remnant termini are in uppercase type. Flanking DNA and empty sites in reference strains 26695 and J99 (gene designations beginning with *hp* and *jhp*, respectively) are in lowercase type. (A) ISHp609 predominant type. (B) ISHp609var type. Terminal sequences were extrapolated from a comparison with sequences of the predominant ISHp609 type, because its flanking sequences did not have homology with known *H. pylori* sequences and therefore the site of insertion (empty site) was not known. (C) Predicted left end of the *orf1-2* remnant and its flanking sequences. Sites of insertion could not be determined precisely due to local sequence heterogeneity in strains lacking this element (intergenic region between *jhp959* and *jhp962*) (see Fig. S2 in the supplemental material). (D) Predicted right end of the *orf1-2* remnant compared to the corresponding region of *orfA* in the ISHp609 sequence.

Strain J99 contains close homologs of *orf1* and *orf2* (*jhp960* and *jhp961*) and the first 37 bp of *orfA*, as noted above. This suggested that this strain might contain a remnant (deleted) version of ISHp609 (Fig. 1C). Related segments of the same size were found by PCR (with primers FlankL and 609.R6) in 15 to 40% of the strains from various geographic regions, some of which also contained full-length ISHp609 elements in other genome locations. As observed with full-length ISHp609, however, this remnant was also rare in East Asian strains (1%) (Table 3). Further PCR tests with *jhp959*- and *jhp962*-specific primers located the remnant between *jhp959* and *jhp962*, as in strain J99, in at least 64 of the 72 cases tested. This inference was confirmed by sequencing PCR products generated with *jhp959* and *jhp962* primers from five strains (LitA10, LitA11, HUP-B50, HUP-B84, and India27); in each case *orf1* (*jhp960*),

orf2 (*jhp961*), and the 5' end of *orfA* were inserted between *jhp959* and *jhp962*, exactly as they are in J99 (predicted left and right ends of the remnant element and insertion site are shown in Fig. 3C and D, respectively). The conservation of the chro-

TABLE 3. Distribution of ISHp609 in different geographic regions

Strain origin	No. of strains	No. (%) ISHp609 positive ^a	No. (%) with <i>orf1-2</i> remnant only ^b
Europe			
Lithuania	47	17 (36)	11 (23)
Spain	24	10 (42)	3 (13)
The Americas			
Alaska	46	20 (43)	12 (26)
Peru	71	12 (17)	13 (18)
Guatemala	29	8 (28)	6 (21)
Africa	28	3 (11)	11 (39)
South Asia: India	69	11 (16)	14 (20)
East Asia			
Japan	71	0	1 (1)
China	47	0	1 (2)
Korea	47	2 (4)	0

^a ISHp609-positive strains were identified by PCR with four sets of primers (primers 609.F1 and 609.R1, primers FlankL and 609R3t1, primers FlankL and 609.R5, and primers FlankL and 609.R1 [Table 1]). Thirty-three of 83 ISHp609 elements were truncated, mostly from the right end. PCR tests indicated that 16 of the 83 ISHp609-positive strains also contained the *orf1-2* remnant (see below), in each case at the location occupied in strain J99, between *jhp959* and *jhp962*.

^b The *orf1-2* remnant contains *orf1*, *orf2*, and the 5' end of *orfA*, as in strain J99 (see text). PCR (with primers *jhp959* and *jhp962*, primers *jhp959* and 609.R6, and primers FlankL and *jhp962*) located it between *jhp959* and *jhp962* in 64 of 72 cases, as in strain J99 (exceptions might have been due to sequence divergence in primer binding sites).

TABLE 2. DNA divergence in ISHp609

Gene	Length (bp)	π_{JC}^a	π_{SIC}^b	π_{NSIC}^c	Ks/Ka ^d
<i>orf1</i>	255	0.009	0.028	0.004	7
<i>orf2</i>	183	0.02	0.045	0.014	3.2
<i>orfA</i>	633	0.008	0.018	0.005	3.6
<i>orfB</i>	1,227	0.01	0.03	0.005	6
Overall	2,298	0.01175	0.03025	0.007	4.95

^a Nucleotide diversity per site. Nucleotide diversity calculations were corrected for multiple hits by using the Jukes-Cantor (JC) correction.

^b Nucleotide diversity per synonymous site.

^c Nucleotide diversity per nonsynonymous site.

^d Ratio of synonymous to nonsynonymous changes.

mosomal location and of *orf1*, *orf2*, and the 5' end of *orfA* sequences suggests that this block of DNA (called the *orf1-2* remnant) resulted from a deletion within a full-length *ISHp609* element and loss of transposition ability and that it spread through *H. pylori* populations by interstrain recombination. High sequence homology (99%) was also found throughout this segment, except for the 3' end of the *jhp959* sequence, which is highly variable (70 to 90% DNA identity) (see Fig. S1 in the supplemental material).

The connection between *jhp959* and *jhp962* was highly conserved among all *H. pylori* strains, independent of the geographic origin; PCR with primers *jhp959* and *jhp962* showed that *jhp959* and *jhp962*, which flanked the *orf1-2* remnant whenever it was present, were next to one another in at least 96% of the *orf1-2* remnant-free strains. This included reference strain 26695, in which the *hp422* gene, a close homolog of *jhp962* (100% protein identity), is adjacent to *hp423*, a distant homolog of *jhp959* (49% protein identity and 64% protein similarity).

To try to define a precise insertion site of the *orf1-2* remnant, we sequenced the putative empty site region in 15 strains that lacked this remnant. Here too *jhp959* sequences and the intergenic region leading to *jhp962* were highly divergent, both in length and in base substitution differences (25 to 30% DNA divergence) (see Fig. S2 in the supplemental material). Consequently, no unique empty site sequence could be discerned. The reason for this extreme diversity is not known. One possible explanation involves slipped strand mispairing during normal DNA replication; an alternative explanation involves deletion of an element and error-prone gap repair.

Other deletions in *ISHp609* were also common; up to one-half of *ISHp609* elements were truncated, mostly from the right end, terminating at different points in *orfB* or *orfA* sequences (Table 2). A distinct 0.9-kb internal deletion due to recombination between 11-bp direct repeats (CCTT[T/G]C TAAAA) located in *orf1* and the 3' end *orfA* was found in *ISHp609* in three of nine Peruvian strains and two of nine Spanish strains (this study). This sequence matched a separately reported sequence from a Costa Rican strain (clone CR2 in reference 6; accession number AF326626). This 0.9-kb deletion was not found in any *ISHp609* element from 15 Lithuanian, 10 Alaskan, 8 Guatemalan, 4 Indian, 3 African, and 2 Korean strains. In all six cases the *ISHp609* element with the 0.9-kb internal deletion was inserted at the same chromosomal site that was defined by PCR (with primers CR2-LF and CR2-RF). Because the 0.9-kb deletion is less widely disseminated than the *orf1-2* remnant, it may have arisen more recently, but it also may have been spread by interstrain recombination. Full-length *ISHp609* sequences from two other strains (one Lithuanian strain and one Guatemalan strain) were found in the same location as the elements containing the 0.9-kb internal deletion.

PCR was used to test for *ISHp609* at each of the other four chromosomal sites identified by sequencing (Fig. 3A) (locations in strains HUP-B43, HUP-B79, LitA7, and LitA38) and also the *jhp959-jhp962* location of the *orf1-2* remnant (with primers in Table 2) in 60 *ISHp609*-positive strains from Spain, Peru, Guatemala, Lithuania, Alaska, Africa, India, and Korea. *ISHp609* was found at the HUP-B43 site (in the *jhp928* homolog) in two of four *ISHp609*-positive Indian strains but not in the other 56 *ISHp609*-positive strains from various countries. No other strain was found to carry *ISHp609* at the specific

sites containing this element in strains LitA7, LitA38, and HUP-B79. Also, no full-length *ISHp609* element was found between *jhp959* and *jhp962*, the site normally occupied by the *orf1-2* remnant.

DISCUSSION

The *ISHp609* element described here is the fifth member of the *IS605* mobile element family found in *H. pylori* and is novel since it contains four open reading frames instead of the usual two. In addition to *orfA* and *orfB*, homologs of which are present in other *H. pylori* IS elements, it also has two other open reading frames, *orf1* and *orf2*, which are members of short, conserved gene families whose functions are unknown that are widespread among eubacterial species. A divergent variant of *ISHp609* was also found, but in only 1 of the 479 strains tested.

ISHp609 was more abundant in strains from Europe, the Americas, and South Asia (10 to 40%) than in strains from East Asia (1%) (Table 3). This is reminiscent of the *ISHp608* distribution (16). The difference between East Asian and other strains in terms of the relative abundance of the *ISHp608* and *ISHp609* elements is in accord with the major geographic differences in the *H. pylori* gene pool, as determined by PCR or multilocus sequence typing of virulence-associated and ordinary housekeeping genes (1, 9, 15, 26, 28), and also with the geographic differences in the sequences of *orfA* and *orfB* from *IS605* and *IS607* elements (11a). This geographic partitioning can be ascribed to *H. pylori*'s preferential transmission within families and local communities rather than in sweeping epidemics (10, 21).

The sequences of *ISHp609* elements from various geographic regions are more closely related to one another (99%) than normal chromosomal gene sequences from the same populations are. This is reminiscent of patterns seen with IS elements of natural isolates of *E. coli*, where the sequences within each IS element type are also more highly conserved than the sequences of chromosomal genes are (11). This, plus the frequent occurrence of elements on bacterial plasmids, suggested that *E. coli* IS elements were spread easily among bacterial lineages by conjugation and transposition (18). Such easy spread would have allowed little time to accumulate neutral variation and would have resulted in the observed sequence homogeneity. Most strains of *H. pylori* are readily transformable, and extensive interstrain recombination is evident in the gene pool (1, 8, 13, 23), although it is restricted geographically because transmission is highly localized. The distribution and sequence uniformity of *ISHp609* suggest that this element entered the *H. pylori* gene pool recently on the evolutionary time scale (perhaps after separation of East Asian strains from other *H. pylori* strains) and spread rapidly, leading to its present abundance in some regions (up to 40% of the strains). This suggests that there was selection for *ISHp609* carriage, stemming from a significant contribution to *H. pylori* host fitness and/or strong molecular drive (selfish DNA). The similar sequence uniformity and distribution of the *orf1-2* remnant (*orf1*, *orf2*, and 5' end of *orfA*) also favor a selection model, but one in which *orf1* and/or *orf2* contributes to fitness. The abundance of truncated (inactive) *ISHp609* elements also suggests such dynamics, but with selection for mutations that rendered many

elements inactive. This is reminiscent of the dynamics of P elements in *Drosophila* (17).

The ISHp609 gene(s) that mediates transposition has not been identified experimentally because we could not, in several tries, detect this element's movement in *E. coli*. The many genomic locations of ISHp609 in *H. pylori* populations imply that there is easy transposition in this host and thus perhaps involvement of a host factor that *E. coli* K-12 lacks. The constancy of the genomic positions of two ISHp609 deletion variants (*orf1-2* remnant and 0.9-kb internal deletion) suggests that neither *orfB* (intact in the 0.9-kb deletion element) nor *orf1* and *orf2* (intact in the *orf1-2* remnant) are sufficient for ISHp609 transposition. Rather, *orfA* of ISHp609 may be needed for movement of this element, as is the case with *orfA* homologs of IS607 and ISHp608 (14, 16).

Other questions concerning transposition mechanisms emerged from the analysis of ISHp609 sequences. The OrfA phylogenies (Fig. 2A) illustrate two distinct subfamilies, one represented by the *orfA* genes of IS607 and ISHp609, which appear to encode serine recombinases, and the other represented by the *orfA* genes of IS605, IS606, and ISHp608, which encode another type of transposition-associated function. Despite ISHp609 *orfA*'s homology to IS607's putative transposase gene, its insertion specificity (downstream of 5'-TAT) seems to be more closely related to that of the IS605-IS606-ISHp608 subfamily (downstream of 5'-TTTAA or 5'-TTTAAAC, 5'-TTTAT, and 5'-TTAC) than to that of IS607 (between GG). Further studies are needed to determine if this stems from functional differences in IS element-encoded proteins or in host factors with which they interact. In terms of *orfB* phylogenies (Fig. 2B), the *orfB* gene of ISHp609 is more distant from the *gipA* *Salmonella* virulence gene than the genes of other IS605 family members are. Therefore, together with its homolog in *T. tengcongensis*, this gene could be considered a member of a distinct subfamily. Most striking in this regard is the C-terminal tetracysteine Zn(II) binding motif CX₍₂₎CX₍₁₅₎CX₍₂₎C (C4-type zinc finger) that could potentially facilitate DNA or RNA binding or protein-protein interaction. This motif is found in *GipA* and the *orfB* products of IS605, IS606, IS607, and ISHp608, but it is absent from the ISHp609 *orfB* product (both predominant and rare variant types) and is also absent from the close homolog in *T. tengcongensis* (Fig. 2C). Although the functions of IS element *orfB* genes are not known, the zinc finger motif difference between OrfB of ISHp609 and the other elements hints at possible differences in the interactions with nucleic acids or other cellular constituents, which in turn might affect transposition or control of host gene expression.

ACKNOWLEDGMENTS

We thank Giedrius Dailide and Shumin Tan for valuable discussions and help.

This research was supported by grants RO1 AI38166, RO3 AI49161, RO1 DK53727, RO1 DK63041, and P30 DK52574 from the National Institutes of Health.

REFERENCES

- Achtman, M., T. Azuma, D. E. Berg, Y. Ito, G. Morelli, Z. J. Pan, S. Suerbaum, S. A. Thompson, A. van der Ende, and L. J. van Doorn. 1999. Recombination and clonal groupings within *Helicobacter pylori* from different geographical regions. *Mol. Microbiol.* **32**:459-470.
- Alm, R. A., L. S. Ling, D. T. Moir, B. L. King, E. D. Brown, P. C. Doig, D. R. Smith, B. Noonan, B. C. Guild, B. L. deJonge, G. Carmel, P. J. Tummino, A. Caruso, M. Uria-Nickelsen, D. M. Mills, C. Ives, R. Gibson, D. Merberg, S. D. Mills, Q. Jiang, D. E. Taylor, G. F. Vovis, and T. J. Trust. 1999. Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* **397**:176-180.
- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. A. Struhl. 1994. Current protocols in molecular biology, supplement 27 CPMB, page 2.4.1. Greene Publishing and Wiley Interscience, New York, N.Y.
- Berg, D. E., R. H. Gilman, J. Lelwala-Guruge, K. Srivastava, Y. Valdez, J. Watanabe, J. Miyagi, N. S. Akopyants, A. Ramirez-Ramos, T. H. Yoshiwara, S. Recavarren, and R. Leon-Barua. 1997. *Helicobacter pylori* populations in individual Peruvian patients. *Clin. Infect. Dis.* **25**:996-1002.
- Berg, D. E., and M. M. Howe (ed.). 1989. Mobile DNA. American Society for Microbiology, Washington, D.C.
- Chanto, G., A. Occhialini, N. Gras, R. A. Alm, F. Megraud, and A. Marais. 2002. Identification of strain-specific genes located outside the plasticity zone in nine clinical isolates of *Helicobacter pylori*. *Microbiology* **148**:3671-3680.
- Craig, N. C., R. Craigie, M. Gellert, and A. M. Lambowitz (ed.). 2002. Mobile DNA II. ASM Press, Washington, D.C.
- Falush, D., C. Kraft, N. S. Taylor, P. Correa, J. G. Fox, M. Achtman, and S. Suerbaum. 2001. Recombination and mutation during long-term gastric colonization by *Helicobacter pylori*: estimates of clock rates, recombination size, and minimal age. *Proc. Natl. Acad. Sci. USA* **98**:15056-15061.
- Falush, D., T. Wirth, B. Linz, J. K. Pritchard, M. Stephens, M. Kidd, M. J. Blaser, D. Y. Graham, S. Vacher, G. I. Perez-Perez, Y. Yamaoka, F. Megraud, K. Otto, U. Reichard, E. Katzowitz, X. Wang, M. Achtman, and S. Suerbaum. 2003. Traces of human migrations in *Helicobacter pylori* populations. *Science* **299**:1582-1585.
- Han, S. R., H. C. Zschausch, H. G. Meyer, T. Schneider, M. Loos, S. Bhakdi, and M. J. Mæurer. 2000. *Helicobacter pylori*: clonal population structure and restricted transmission within families revealed by molecular typing. *J. Clin. Microbiol.* **38**:3646-3651.
- Hartl, D. L., and S. A. Sawyer. 1988. Why do unrelated insertion sequences occur together in the genome of *Escherichia coli*? *Genetics* **118**:537-541.
- Kalia, A., A. K. Mukhopadhyay, G. Dailide, Y. Ito, T. Azuma, B. C. Y. Wong, and D. E. Berg. 2004. Evolutionary dynamics of insertion sequences in *H. pylori*. *J. Bacteriol.* **186**:7508-7520.
- Kersulyte, D., N. S. Akopyants, S. W. Clifton, B. A. Roe, and D. E. Berg. 1998. Novel sequence organization and insertion specificity of IS605 and IS606: chimaeric transposable elements of *Helicobacter pylori*. *Gene* **223**:175-186.
- Kersulyte, D., H. Chalkauskas, and D. E. Berg. 1999. Emergence of recombinant strains of *Helicobacter pylori* during human infection. *Mol. Microbiol.* **31**:31-43.
- Kersulyte, D., A. K. Mukhopadhyay, M. Shirai, T. Nakazawa, and D. E. Berg. 2000. Functional organization and insertion specificity of IS607, a chimeric element of *Helicobacter pylori*. *J. Bacteriol.* **182**:5300-5308.
- Kersulyte, D., A. K. Mukhopadhyay, B. Velapatino, W. W. Su, Z. J. Pan, C. Garcia, V. Hernandez, Y. Valdez, R. S. Mistry, R. H. Gilman, Y. Yuan, H. Gao, T. Alarcon, M. Lopez-Brea, G. Balakrish Nair, A. Chowdhury, S. Datta, M. Shirai, T. Nakazawa, R. Ally, I. Segal, B. C. Wong, S. K. Lam, F. O. Olfat, T. Boren, L. Engstrand, O. Torres, R. Schneider, J. E. Thomas, S. Czinn, and D. E. Berg. 2000. Differences in genotypes of *Helicobacter pylori* from different human populations. *J. Bacteriol.* **182**:3210-3218.
- Kersulyte, D., B. Velapatino, G. Dailide, A. K. Mukhopadhyay, Y. Ito, L. Cahuayme, A. J. Parkinson, R. H. Gilman, and D. E. Berg. 2002. Transposable element ISHp608 of *Helicobacter pylori*: nonrandom geographic distribution, functional organization, and insertion specificity. *J. Bacteriol.* **184**:992-1002.
- Kidwell, M. G., and D. R. Lisch. 2001. Perspective: transposable elements, parasitic DNA, and genome evolution. *Evol. Int. J. Org. Evol.* **55**:1-24.
- Lawrence, J. G., H. Ochman, and D. L. Hartl. 1992. The evolution of insertion sequences within enteric bacteria. *Genetics* **131**:9-20.
- Mahillon, J., and M. Chandler. 1998. Insertion sequences. *Microbiol. Mol. Biol. Rev.* **62**:725-774.
- Mukhopadhyay, A. K., D. Kersulyte, J. Y. Jeong, S. Datta, Y. Ito, A. Chowdhury, S. Chowdhury, A. Santra, S. K. Bhattacharya, T. Azuma, G. B. Nair, and D. E. Berg. 2000. Distinctiveness of genotypes of *Helicobacter pylori* in Calcutta, India. *J. Bacteriol.* **182**:3219-3227.
- Owen, R. J., and J. Kerry. 2003. Tracing clonality of *Helicobacter pylori* infecting family members from analysis of DNA sequences of three house-keeping genes (*ureI*, *atpA* and *ahpC*), deduced amino acid sequences, and pathogenicity-associated markers (*cagA* and *vacA*). *J. Med. Microbiol.* **52**:515-524.
- Stanley, T. L., C. D. Ellermeier, and J. M. Schlauch. 2000. Tissue-specific gene expression identifies a gene in the lysogenic phage Gifsy-1 that affects *Salmonella enterica* serovar Typhimurium survival in Peyer's patches. *J. Bacteriol.* **182**:4406-4413.
- Suerbaum, S., J. M. Smith, K. Bapumia, G. Morelli, N. H. Smith, E. Kunstmann, I. Dyrek, and M. Achtman. 1998. Free recombination within *Helicobacter pylori*. *Proc. Natl. Acad. Sci. USA* **95**:12619-12624.

24. Tomb, J. F., O. White, A. R. Kerlavage, R. A. Clayton, G. G. Sutton, R. D. Fleischmann, K. A. Ketchum, H. P. Klenk, S. Gill, B. A. Dougherty, K. Nelson, J. Quackenbush, L. Zhou, E. F. Kirkness, S. Peterson, B. Loftus, D. Richardson, R. Dodson, H. G. Khalak, A. Glodek, K. McKenney, L. M. Fitzgerald, N. Lee, M. D. Adams, E. K. Hickey, D. E. Berg, J. D. Gocayne, T. R. Utterback, J. D. Peterson, J. M. Kelley, M. D. Cotton, J. M. Weidman, C. Fujii, C. Bowman, L. Watthey, E. Wallin, W. S. Hayes, M. Borodovsky, P. D. Karp, H. O. Smith, C. M. Fraser, and J. C. Venter. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **388**:539–547.
25. Turlan, C., and M. Chandler. 2000. Playing second fiddle: second-strand processing and liberation of transposable elements from donor DNA. *Trends Microbiol.* **8**:268–274.
26. van der Ende, A., Z. J. Pan, A. Bart, R. W. van der Hulst, M. Feller, S. D. Xiao, G. N. Tytgat, and J. Dankert. 1998. *cagA*-positive *Helicobacter pylori* populations in China and The Netherlands are distinct. *Infect. Immun.* **66**:1822–1826.
27. Westblom, T. U. 1991. Laboratory diagnosis and handling of *Helicobacter pylori*, p 81–91. *In* B. J. Marshall, R. W. McCallum, and R. L. Guerrant (ed.), *Helicobacter pylori* in peptic ulceration and gastritis. Blackwell Scientific Publications, Boston, Mass.
28. Yamaoka, Y., M. S. Osato, A. R. Sepulveda, O. Gutierrez, N. Figura, J. G. Kim, T. Kodama, K. Kashima, and D. Y. Graham. 2000. Molecular epidemiology of *Helicobacter pylori*: separation of *H. pylori* from East Asian and non-Asian countries. *Epidemiol. Infect.* **124**:91–96.