

2014

Draft genome sequence of *Acetobacter aceti* strain 1023, a vinegar factory isolate

John E. Hung
Washington University in St Louis

Christopher P. Mill
Washington University in St Louis

Sandra W. Clifton
Washington University School of Medicine in St. Louis

Vincent Magrini
Washington University School of Medicine in St. Louis

Ketaki Bhide
Purdue University

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Hung, John E.; Mill, Christopher P.; Clifton, Sandra W.; Magrini, Vincent; Bhide, Ketaki; Francois, Julie A.; Ransome, Aaron E.; Fulton, Lucinda; Thimmapuram, Jyothi; Wilson, Richard K.; and Kappock, T. Joseph, "Draft genome sequence of *Acetobacter aceti* strain 1023, a vinegar factory isolate." *Genome Announcements*. 2, 3. e00550-14. (2014).
https://digitalcommons.wustl.edu/open_access_pubs/2933

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

John E. Hung, Christopher P. Mill, Sandra W. Clifton, Vincent Magrini, Ketaki Bhide, Julie A. Francois, Aaron E. Ransome, Lucinda Fulton, Jyothi Thimmapuram, Richard K. Wilson, and T. Joseph Kappock

Draft Genome Sequence of *Acetobacter aceti* Strain 1023, a Vinegar Factory Isolate

John E. Hung,^{a*} Christopher P. Mill,^{a*} Sandra W. Clifton,^{b*} Vincent Magrini,^{b,c} Ketaki Bhide,^d Julie A. Francois,^{a*} Aaron E. Ransome,^{a*} Lucinda Fulton,^b Jyothi Thimmapuram,^d Richard K. Wilson,^{b,c} T. Joseph Kappock^e

Department of Chemistry, Washington University in St. Louis, St. Louis, Missouri, USA^a; The Genome Institute, Washington University School of Medicine, St. Louis, Missouri, USA^b; Department of Genetics, Washington University School of Medicine, St. Louis, Missouri, USA^c; Bioinformatics Core, Discovery Park, Purdue University, West Lafayette, Indiana, USA^d; Department of Biochemistry, Purdue University, West Lafayette, Indiana, USA^e

* Present address: John E. Hung, Sigma-Aldrich Company, St. Louis, Missouri, USA; Christopher P. Mill, Department of Biochemistry, St. Jude Children's Research Hospital, Memphis, Tennessee, USA; Sandra W. Clifton, Department of Chemistry and Biochemistry, Advanced Center for Genome Technology, University of Oklahoma, Norman, Oklahoma, USA; Julie A. Francois, Monsanto Company, Creve Coeur, Missouri, USA; Aaron E. Ransome, GEVO, Englewood, Colorado, USA.

The genome sequence of *Acetobacter aceti* 1023, an acetic acid bacterium adapted to traditional vinegar fermentation, comprises 3.0 Mb (chromosome plus plasmids). *A. aceti* 1023 is closely related to the cocoa fermenter *Acetobacter pasteurianus* 386B but possesses many additional insertion sequence elements.

Received 16 May 2014 Accepted 20 May 2014 Published 5 June 2014

Citation Hung JE, Mill CP, Clifton SW, Magrini V, Bhide K, Francois JA, Ransome AE, Fulton L, Thimmapuram J, Wilson RK, Kappock TJ. 2014. Draft genome sequence of *Acetobacter aceti* strain 1023, a vinegar factory isolate. *Genome Announc.* 2(3):e00550-14. doi:10.1128/genomeA.00550-14.

Copyright © 2014 Hung et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](#).

Address correspondence to T. Joseph Kappock, kappock@purdue.edu.

Acetic acid bacteria (AAB) are acidophilic aerobic alphaproteobacteria with many uses in food processing (1, 2). *Acetobacter aceti* strain 1023, a traditional rice vinegar mash surface isolate (3), was used in pioneering studies of AAB physiology (4). The continual selection of vinegar strains has favored acetic acid/ethanol resistance traits and disfavored wasteful overoxidation, in which acetic acid is lost as CO₂ (5). Whole-genome sequencing of *A. aceti* 1023 was used to identify adaptations in this highly domesticated vinegar strain.

A. aceti 1023 was propagated at 30°C in yeast-peptone-dextrose medium supplemented with 2% ethanol. Genomic DNA was used to prepare plasmid (4.1- and 6.1-kb inserts in plasmid pOTW13) and fosmid (40-kb inserts in pCC-FOS1) libraries, as previously described (6, 7). Using PCAP (8), paired-end Sanger reads were assembled (28,731 reads, 76% input) into 337 contigs >1 kb (total, 3.2 Mb; N₅₀, 17,669 bp), as was disclosed in a preliminary form (9).

Genomic DNA libraries were analyzed by 454 GS-FLX pyrosequencing using both fragment (564,984 reads, 140 Mb total) and mate-pair (3-kb insert; 468,069 reads, 66 Mb total) libraries. A hybrid assembly of Sanger and 454 reads using Newbler (version 2.9) furnished 33 scaffolds composed of 193 contigs (>0.5 kb) and 3.0 Mb total sequence at 72-fold coverage. The scaffolds were ordered with Mauve (version 2.3.1) (10), using the complete genome sequence of *Acetobacter pasteurianus* 386B (11) as the template. The NCBI Prokaryotic Genome Annotation Pipeline (version 2.5) and BLASTn analysis predicted 2,650 open reading frames, 66 pseudogenes, and 47 functional RNAs. At least eight scaffolds (0.07 Mb total) appeared to originate from plasmids, as judged by the presence of *repBA* and plasmid partitioning genes. As is typical for the low-copy-number AAB “cryptic” plasmids (12), the plasmid scaffolds contain few genes that clearly confer a phenotype.

A phylogenetic analysis of AAB GroEL sequences (13) grouped *A. aceti* 1023 with *A. pasteurianus* and *Acetobacter pomorum*, not *A. aceti* NBRC 14818 or ATCC 23746. Central carbon metabolism is more straightforward in *A. aceti* 1023 and *A. pasteurianus* strains, which use a specialized citric acid cycle containing *aarC* (14), than in *A. aceti* NBRC 14818, which has greater metabolic versatility (15–17). As judged by gene synteny and sequence similarity, *A. aceti* 1023 has a particularly close relationship to *A. pasteurianus* 386B, a cocoa fermenter (11). However, *A. pasteurianus* 386B lacks numerous insertion sequence (IS) elements present in the vinegar strains *A. aceti* 1023 and *A. pasteurianus* NBRC 3283 (18). As anticipated from Southern blots (19–21), *A. aceti* 1023 contains IS1380, IS1452, and IS12538, with minimal copy numbers of 64, 4, and 1, respectively. The adaptation of a common ancestor to different fermentation milieus involved divergent histories of transposable element acquisition in *A. aceti* 1023 and *A. pasteurianus* 386B.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [JEOA00000000](#). The version described in this paper is the first version, [JEOA01000000](#).

ACKNOWLEDGMENTS

This work was supported by funds from the College of Agriculture at Purdue University and grants from the Danforth Foundation at Washington University, the Herman Frasch Foundation for Research in Agricultural Chemistry (531HF-02), and the National Science Foundation (MCB 0347250 and an REU supplement for J.E.H.) to T.J.K.

We thank the staff of the Genome Institute at Washington University, including Catrina Fronick, Michael Becker, Kim Delehaunty, Kevin Haub, Shelly O’Laughlin, and Brenda Theising, for library construction and DNA sequencing and finishing. We thank the staff of the Purdue University Genomics Core Facility for library construction and DNA sequencing, and Phillip San Miguel and Rick Westerman for advice and

assistance with preliminary assemblies. We also thank Koichi Kondo (Mizkan Group Corporation) for the gift of the title strain and Akiko Okamoto-Kainuma (Tokyo University of Agriculture) for guidance.

REFERENCES

- Cleenwerck I, De Vos P. 2008. Polyphasic taxonomy of acetic acid bacteria: an overview of the currently applied methodology. *Int. J. Food Microbiol.* 125:2–14. <http://dx.doi.org/10.1016/j.ijfoodmicro.2007.04.017>.
- Raspor P, Goranovic D. 2008. Biotechnological applications of acetic acid bacteria. *Crit. Rev. Biotechnol.* 28:101–124. <http://dx.doi.org/10.1080/07388550802046749>.
- Ohmori S, Masai H, Arima K, Beppu T. 1980. Isolation and identification of acetic acid bacteria for submerged acetic acid fermentation at high temperature. *Agric. Biol. Chem.* 44:2901–2906. <http://dx.doi.org/10.1271/bbb1961.44.2901>.
- Beppu T. 1994. Genetic organization of *Acetobacter* for acetic acid fermentation. *Antonie Van Leeuwenhoek* 64:121–135. <http://dx.doi.org/10.1007/BF00873022>.
- Saeki A, Taniguchi M, Matsushita K, Toyama H, Theeragool G, Lotong N, Adachi O. 1997. Microbiological aspects of acetate oxidation by acetic acid bacteria, unfavorable phenomena in vinegar fermentation. *Biosci. Biotechnol. Biochem.* 61:317–323. <http://dx.doi.org/10.1271/bbb.61.317>.
- Magrini V, Warren WC, Wallis J, Goldman WE, Xu J, Mardis ER, McPherson JD. 2004. Fosmid-based physical mapping of the *Histoplasma capsulatum* genome. *Genome Res.* 14:1603–1609. <http://dx.doi.org/10.1101/gr.2361404>.
- Wild J, Hradecna Z, Szybalski W. 2002. Conditionally amplifiable BACs: switching from single-copy to high-copy vectors and genomic clones. *Genome Res.* 12:1434–1444. <http://dx.doi.org/10.1101/gr.130502>.
- Huang X, Wang J, Aluru S, Yang SP, Hillier L. 2003. PCAP: a whole-genome assembly program. *Genome Res.* 13:2164–2170. <http://dx.doi.org/10.1101/gr.1390403>.
- Francois JA, Kappock TJ. 2007. Alanine racemase from the acidophile *Acetobacter aceti*. *Protein Expr. Purif.* 51:39–48. <http://dx.doi.org/10.1016/j.pep.2006.05.016>.
- Rissman AI, Mau B, Biehl BS, Darling AE, Glasner JD, Perna NT. 2009. Reordering contigs of draft genomes using the Mauve aligner. *Bioinformatics* 25:2071–2073. <http://dx.doi.org/10.1093/bioinformatics/btp356>.
- Illegghems K, De Vuyst L, Weckx S. 2013. Complete genome sequence and comparative analysis of *Acetobacter pasteurianus* 386B, a strain well-adapted to the cocoa bean fermentation ecosystem. *BMC Genomics* 14: 526. <http://dx.doi.org/10.1186/1471-2164-14-526>.
- Fukaya M, Iwata T, Entani E, Masai H, Uozumi T, Beppu T. 1985. Distribution and characterization of plasmids in acetic acid bacteria. *Agric. Biol. Chem.* 49:1349–1355. <http://dx.doi.org/10.1271/bbb1961.49.1349>.
- Huang CH, Chang MT, Huang L, Chua WS. 2014. Molecular discrimination and identification of *Acetobacter* genus based on the partial heat shock protein 60 gene (*hsp60*) sequences. *J. Sci. Food Agric.* 94:213–218. <http://dx.doi.org/10.1002/jsfa.6231>.
- Mullins EA, Francois JA, Kappock TJ. 2008. A specialized citric acid cycle requiring succinyl-coenzyme A (CoA):acetate CoA-transferase (AarC) confers acetic acid resistance on the acidophile *Acetobacter aceti*. *J. Bacteriol.* 190:4933–4940. <http://dx.doi.org/10.1128/JB.00405-08>.
- Matsutani M, Hirakawa H, Yakushi T, Matsushita K. 2011. Genome-wide phylogenetic analysis of *Gluconobacter*, *Acetobacter*, and *Gluconacetobacter*. *FEMS Microbiol. Lett.* 315:122–128. <http://dx.doi.org/10.1111/j.1574-6968.2010.02180.x>.
- Sakurai K, Arai H, Ishii M, Igarashi Y. 2011. Transcriptome response to different carbon sources in *Acetobacter aceti*. *Microbiology* 157:899–910. <http://dx.doi.org/10.1099/mic.0.045906-0>.
- Sakurai K, Yamazaki S, Ishii M, Igarashi Y, Arai H. 2013. Role of the glyoxylate pathway in acetic acid production by *Acetobacter aceti*. *J. Biosci. Bioeng.* 115:32–36. <http://dx.doi.org/10.1016/j.jbiosc.2012.07.017>.
- Azuma Y, Hosoyama A, Matsutani M, Furuya N, Horikawa H, Harada T, Hirakawa H, Kuhara S, Matsushita K, Fujita N, Shirai M. 2009. Whole-genome analyses reveal genetic instability of *Acetobacter pasteurianus*. *Nucleic Acids Res.* 37:5768–5783. <http://dx.doi.org/10.1093/nar/gkp612>.
- Kondo K, Horinouchi S. 1997. A new insertion sequence IS1452 from *Acetobacter pasteurianus*. *Microbiology* 143:539–546. <http://dx.doi.org/10.1099/00221287-143-2-539>.
- Kondo K, Horinouchi S. 1997. Characterization of an insertion sequence, IS12528, from *Gluconobacter suboxydans*. *Appl. Environ. Microbiol.* 63: 1139–1142.
- Takemura H, Horinouchi S, Beppu T. 1991. Novel insertion sequence IS1380 from *Acetobacter pasteurianus* is involved in loss of ethanol-oxidizing ability. *J. Bacteriol.* 173:7070–7076.