

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2014

Symptoms of pain do not correlate with rotator cuff tear severity

Robert H. Brophy

Washington University School of Medicine in St. Louis

Matthew V. Smith

Washington University School of Medicine in St. Louis

Rick W. Wright

Washington University School of Medicine in St. Louis

et al

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Brophy, Robert H.; Smith, Matthew V.; Wright, Rick W.; and et al, "Symptoms of pain do not correlate with rotator cuff tear severity." *The Journal of Bone and Joint Surgery*. 96, 10. 793-800. (2014).

https://digitalcommons.wustl.edu/open_access_pubs/2947

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Symptoms of Pain Do Not Correlate with Rotator Cuff Tear Severity

A Cross-Sectional Study of 393 Patients with a Symptomatic Atraumatic Full-Thickness Rotator Cuff Tear

Warren R. Dunn, MD, MPH, John E. Kuhn, MD, MS, Rosemary Sanders, BA, Qi An, MS, Keith M. Baumgarten, MD, Julie Y. Bishop, MD, Robert H. Brophy, MD, James L. Carey, MD, MPH, G. Brian Holloway, MD, Grant L. Jones, MD, C. Benjamin Ma, MD, Robert G. Marx, MD, MS, Eric C. McCarty, MD, Sourav K. Poddar, MD, Matthew V. Smith, MD, Edwin E. Spencer, MD, Armando F. Vidal, MD, Brian R. Wolf, MD, MS, and Rick W. Wright, MD,
on behalf of the MOON Shoulder Group

Background: For many orthopaedic disorders, symptoms correlate with disease severity. The objective of this study was to determine if pain level is related to the severity of rotator cuff disorders.

Methods: A cohort of 393 subjects with an atraumatic symptomatic full-thickness rotator-cuff tear treated with physical therapy was studied. Baseline pretreatment data were used to examine the relationship between the severity of rotator cuff disease and pain. Disease severity was determined by evaluating tear size, retraction, superior humeral head migration, and rotator cuff muscle atrophy. Pain was measured on the 10-point visual analog scale (VAS) in the patient-reported American Shoulder and Elbow Surgeons (ASES) score. A linear multiple regression model was constructed with use of the continuous VAS score as the dependent variable and measures of rotator cuff tear severity and other non-anatomic patient factors as the independent variables. Forty-eight percent of the patients were female, and the median age was sixty-one years. The dominant shoulder was involved in 69% of the patients. The duration of symptoms was less than one month for 8% of the patients, one to three months for 22%, four to six months for 20%, seven to twelve months for 15%, and more than a year for 36%. The tear involved only the supraspinatus in 72% of the patients; the supraspinatus and infraspinatus, with or without the teres minor, in 21%; and only the subscapularis in 7%. Humeral head migration was noted in 16%. Tendon retraction was minimal in 48%, midhumeral in 34%, glenohumeral in 13%, and to the glenoid in 5%. The median baseline VAS pain score was 4.4.

Results: Multivariable modeling, controlling for other baseline factors, identified increased comorbidities ($p = 0.002$), lower education level ($p = 0.004$), and race ($p = 0.041$) as the only significant factors associated with pain on presentation. No measure of rotator cuff tear severity correlated with pain ($p > 0.25$).

Conclusions: Anatomic features defining the severity of atraumatic rotator cuff tears are not associated with the pain level. Factors associated with pain are comorbidities, lower education level, and race.

Level of Evidence: Prognostic Level III. See Instructions for Authors for a complete description of levels of evidence.

Peer Review: This article was reviewed by the Editor-in-Chief and one Deputy Editor, and it underwent blinded review by two or more outside experts. It was also reviewed by an expert in methodology and statistics. The Deputy Editor reviewed each revision of the article, and it underwent a final review by the Editor-in-Chief prior to publication. Final corrections and clarifications occurred during one or more exchanges between the author(s) and copyeditors.

Disclosure: One or more of the authors received payments or services, either directly or indirectly (i.e., via his or her institution), from a third party in support of an aspect of this work. In addition, one or more of the authors, or his or her institution, has had a financial relationship, in the thirty-six months prior to submission of this work, with an entity in the biomedical arena that could be perceived to influence or have the potential to influence what is written in this work. No author has had any other relationships, or has engaged in any other activities, that could be perceived to influence or have the potential to influence what is written in this work. The complete **Disclosures of Potential Conflicts of Interest** submitted by authors are always provided with the online version of the article.



A commentary by Frederick A. Matsen III, MD, is linked to the online version of this article at jbsj.org.

For many orthopaedic disorders, such as osteoarthritis, symptoms correlate with disease severity¹⁻³. However, symptoms related to rotator cuff tears are not as predictable. It is well known that many people in the general population have an asymptomatic rotator cuff tear⁴⁻⁶. Patients may become asymptomatic following rotator cuff repair despite evidence that the repair failed on magnetic resonance imaging (MRI) or ultrasound imaging⁷. Nonoperative treatment for full-thickness rotator cuff tears is often successful despite a lack of tendon healing⁸⁻¹¹. As a result, the link between rotator cuff tears and symptoms is not clearly established. Pain is a frequently cited symptom associated with rotator cuff tears and is often used as an indication for surgery¹²⁻¹⁵. The objective of this investigation was to determine if the pain level is related to the severity of the rotator cuff disease. We used a cross-sectional design to test the hypothesis that increased pain correlates with greater rotator cuff tear severity.

Materials and Methods

The MOON (Multicenter Orthopaedic Outcomes Network) Shoulder Group is a team of sixteen fellowship-trained orthopaedic surgeons and research personnel from nine geographically dispersed sites, representing both academic and private practice settings, within the United States. This group was formed to conduct large multicenter studies on conditions of the shoulder.

From May 2004 through October 2006, the MOON Shoulder Group met regularly to formulate research questions of interest; develop and standardize radiographic and MRI protocols; assemble validated behavioral and patient-oriented outcome assessment forms for data collection; and conduct validation studies on MRI classification of rotator cuff tears¹⁶, classification of rotator cuff tears based on arthroscopic videos¹⁷, and radiographic findings associated with rotator cuff disease¹⁸. In addition, the group performed systematic reviews of the literature to evaluate rehabilitation following rotator cuff repairs¹⁹, summarize the literature regarding indications for surgical treatment of rotator cuff tears²⁰, and determine the effectiveness of physical therapy in treating rotator cuff disease while developing an evidence-based protocol²¹.

TABLE I Demographic Features of Cohort

	No. of Patients	Female (N = 190)	Male (N = 203)
Age* (yr)	393	63.2 ± 10.2	62.1 ± 9.9
BMI* (kg/m ²)	388	28.3 ± 7.2	29.0 ± 5.1
Involved side	390		
Nondominant		27% (51/188)	35% (70/202)
Dominant		73% (137/188)	65% (132/202)
Race	389		
Other		7% (13/187)	4% (9/202)
Black		12% (23/187)	4% (8/202)
White		81% (151/187)	92% (185/202)
Education	392		
High school or less		27% (51/189)	35% (72/203)
Some college		31% (59/189)	21% (43/203)
Bachelor's degree		20% (38/189)	19% (39/203)
Graduate degree		22% (41/189)	24% (49/203)
Marital status	391		
Other		5% (10/189)	3% (7/202)
Divorced		20% (37/189)	9% (18/202)
Married		48% (90/189)	82% (165/202)
Single		10% (18/189)	3% (6/202)
Widowed		18% (34/189)	3% (6/202)
Employment	392		
Full-time		38% (72/189)	55% (112/203)
Part-time		12% (23/189)	7% (14/203)
Retired		34% (64/189)	32% (65/203)
Homemaker		9% (17/189)	0% (0/203)
Not working		7% (13/189)	6% (12/203)
Smoking	390		
No		90% (170/188)	90% (182/202)
Yes		10% (18/188)	10% (20/202)

*The values are given as the mean and standard deviation.

TABLE II Features of Rotator Cuff Disease in the Cohort

	No. of Patients	Female (N = 190)	Male (N = 203)
Duration of symptoms	390		
<1 mo		6% (12/189)	9% (18/201)
1-3 mo		22% (41/189)	22% (44/201)
4-6 mo		22% (41/189)	18% (37/201)
7-12 mo		18% (34/189)	11% (23/201)
>12 mo		32% (61/189)	39% (79/201)
Involved tendons	383		
Supraspinatus		75% (139/186)	70% (137/197)
Supraspinatus + infraspinatus ± teres minor		20% (37/186)	22% (44/197)
Subscapularis		5% (10/186)	8% (16/197)
Retraction	390		
Minimal		49% (92/188)	48% (96/202)
Midhumeral		34% (63/188)	35% (70/202)
Glenohumeral		13% (25/188)	12% (25/202)
Glenoid		4% (8/188)	5% (11/202)
Superior migration	380		
No		87% (163/187)	82% (158/193)
Yes		13% (24/187)	18% (35/193)
Torn tendons	384		
1		75% (139/186)	70% (139/198)
≥2		25% (47/186)	30% (59/198)
MRI quantity of supraspinatus	387		
Normal		40% (74/186)	53% (107/201)
<25% atrophy		31% (58/186)	22% (45/201)
25-50% atrophy		22% (40/186)	13% (27/201)
50-75% atrophy		6% (12/186)	7% (15/201)
Complete atrophy		1% (2/186)	3% (7/201)
Acromion shape	377		
Type I		13% (23/182)	10% (20/195)
Type II		70% (127/182)	73% (142/195)
Type III		18% (32/182)	17% (33/195)
Acromiohumeral interval* (mm)	339	8.0, 10.0, 11.0 (9.8 ± 6.8)	8.0, 10.0, 11.0 (10.2 ± 7.2)

*The values are given as the lower quartile, median, upper quartile (mean and standard deviation).

The indications for surgery for atraumatic rotator cuff tears are not clear^{12,20,22}, and our research group could not develop standard indications for surgery by consensus. Therefore, the group conducted a prospective cohort study to assess the nonoperative treatment of atraumatic full-thickness rotator cuff tears with use of the physical therapy protocol derived from the systematic review. We expected that nonoperative treatment would succeed for some patients, who would then decline surgical intervention, while it would fail for others, who would then undergo rotator cuff repair. By identifying features that distinguish these groups, we sought insight into the risk factors for failure of nonoperative management to better define the appropriate indications for surgery. To test the hypothesis that pain correlates with measures of rotator cuff tear severity, we performed a cross-sectional study of data derived from this population when patients initially presented to the treating physician. Sample-size requirements for that study were formulated with the events-per-variable approach, with ten events as the minimum number to avoid overfitting of the model. Baseline data on the first 393 subjects enrolled into the ongoing prospective cohort study were used in the present study.

Institutional Review Board Approval

Institutional review board approval was obtained at all participating sites before enrollment of the first patient.

Inclusion and Exclusion Criteria

All patients eighteen to 100 years old with an MRI-documented full-thickness atraumatic symptomatic rotator cuff tear were invited to participate. Exclusion criteria included an injury that precipitated the pain, pain determined to be related to the cervical spine, scapular pain, previous shoulder surgery, glenohumeral arthritis, inflammatory arthritis, adhesive capsulitis, a previous proximal humeral fracture, a symptomatic contralateral rotator cuff tear, and dementia.

Protocol

Patients were recruited over a four-year period (January 2007 to January 2011). At the initial visit, patients completed a questionnaire that detailed demographic

data and included the following validated patient-reported outcome measures: the Short Form-12 (SF-12)²³, American Shoulder and Elbow Surgeons (ASES) score²⁴, Western Ontario Rotator Cuff (WORC) Index²⁵, Single Assessment Numeric Evaluation (SANE) score²⁶, Self-Administered Comorbidity Questionnaire (SCQ)²⁷, and the Shoulder Activity Scale²⁸. Patient demographic information including age, sex, race, employment status, Workers' Compensation or automobile insurance claims, and tobacco use history were also collected at the time of enrollment.

Physicians at each site examined patients and images and recorded information regarding shoulder motion, strength, tenderness, provocative signs, and descriptors of radiographs and MRI scans. The following MRI features used to assess the severity of a rotator cuff tear had reasonably high interobserver agreement when studied by our research group¹⁶ and were used in this analysis: (1) the rotator cuff tendons that were seen to be involved by the tear on MRI (72% agreement, kappa = 0.55), (2) retraction in the sagittal plane seen on MRI as described by Patte²⁹ (63% agreement, kappa = 0.44), (3) presence of superior humeral head migration on standing anteroposterior radiographs (52% agreement, kappa = 0.26), and (4) amount of supraspinatus atrophy³⁰ seen on MRI (59% agreement, kappa = 0.25).

Statistical Methods

To evaluate the association between shoulder pain and other baseline factors, a multivariable linear multiple regression model was fit with use of a continuous visual analog scale (VAS) pain score (item 3 from the ASES patient form²⁴) as the dependent variable. Independent variables included in the model were age, sex, body mass index (BMI), duration of symptoms, activity level, handedness, education, occupation, race, smoking status, patient expectations, tendons torn, retraction, humeral head migration, and amount of atrophy in the supraspinatus.

We did not assume linearity of covariate effects but only assumed smoothed relationships, using restricted cubic regression splines. Missing

values of predictor variables were imputed with use of multiple imputation incorporating predictive mean matching and flexible additive imputation models as implemented in the *aregImpute* function available in the Hmisc package in R³¹. This procedure involves predicting missing values on the basis of nonmissing observed data—essentially finding matches for the missing values. A regression model is fit to the observed data, then the missing values for a particular covariate are matched to the closest set of observed values on the basis of the regression fit, and then missing values are imputed on the basis of the closest set of observed values. This process was repeated ten times; hence, the imputed values were averaged over ten imputations. Pooling of low-prevalence categories was performed for data reduction to preserve degrees of freedom in the model, and to avoid convergence problems with the model. Specifically, American Indian, Asian, and Hawaiian were collapsed into an “other” category for race; disabled and unemployed were combined into a “not working” category for employment status; and no high school and some high school were collapsed into a “high school or less” category for education. Statistical analysis was performed with free open-source R statistical software (www.r-project.org).

Population and Rotator Cuff Demographics

The group evaluated 2233 patients who presented with a rotator cuff tear during the enrollment period; 1280 of these patients were excluded because of a traumatic tear (38%), previous surgery (11%), bilateral disease (8%), neck disorder (6%), frozen shoulder (2%), glenohumeral dislocation (3%), rheumatoid disease (1%), or fracture (1%). Of the remaining 953 patients eligible to enroll in the study, 452 (47%) elected to do so. The first 393 patients in this cohort were included in the present cross-sectional study, in which we used data from the first visit at the time of the initial enrollment in the study.

The average age of the patients who enrolled was sixty-two years, whereas the average age of those who did not was fifty-eight years ($p < 0.001$).




Fig. 1

Plot of the effects of the significant predictors in the model. The gray bars represent the 95% CI for the mean effect. The mean effect of comorbidity on pain was determined by comparing those with an SCQ score of 2 with those with a score of 10; the VAS pain score (ases3) was increased by 1.07 (95% CI: 0.31, 1.83) in the latter group. HS = high school, Grad = graduate, and Bach = Bachelor's.

Equal numbers of men and women enrolled, whereas 63% of those who did not enroll were male; this was a significant difference ($p < 0.001$).

Demographic Data for the Study Population

The median age of the study population was sixty-one years (range, thirty-one to ninety years), and 52% were male. The dominant arm was affected in 69% of the subjects; 90% were nonsmokers. Other demographic features, including race, education level, and employment status, are listed in Table I stratified by sex. Twenty-three percent of the patients had already tried some physical therapy, 40% had received injections, and 80% had tried nonsteroidal anti-inflammatory drugs. The median baseline VAS pain score was 4.4.

MRI Features of the Rotator Cuff Tears

The tears involved only the supraspinatus in 72% of the patients. Retraction was minimal in 48% of the patients and was to the midpart of the humeral head in 34%. Superior humeral head migration was present in 16% of the patients. Tear characteristics stratified by sex are listed in Table II.

Source of Funding

Sources of funding included an unrestricted gift from the Arthrex Corporation, and a research funding grant specific to the investigation from the National Football League (NFL) Foundation, which were used to support research personnel and pay for supplies and computer hardware and software. Career development grants from the National Institutes of Health (NIH) and the American Orthopaedic Society for Sports Medicine (AOSSM) supported Dr. Dunn.

Results

Factors Associated with Pain

Greater pain was associated with an increased number of comorbidities ($p = 0.002$), a lower education level ($p = 0.004$), and race ($p = 0.041$). Measures of tear severity, including the tendons involved ($p = 0.5$), amount of retraction ($p = 0.9$), presence of humeral head migration ($p = 0.3$), and amount of fatty degeneration of the supraspinatus ($p = 0.4$), were not associated with pain. These findings were adjusted for the other covariates included in the model—specifically, duration of symptoms, occupation, smoking status, activity level, BMI, dominant arm, and patient expectations. A summary-of-effects plot is shown in Figure 1 with the 95% confidence intervals (CIs) for the mean effects.

While the SCQ score, a lower level of education, and race were significant factors, the respective point estimates shown in Figure 1 do not represent a clinically meaningful difference in the VAS pain score, which is 1.4 cm for subjects with rotator cuff disease³². However, using the nomogram in Figure 2, one can estimate the cumulative effects of the predictors on pain. For instance, the nomogram can be used to estimate the effect of race, education level, and comorbidity (allowing the other factors depicted on the nomogram to default to the left-most category, which contributes no points to the calculation) as follows. A sum




Fig. 2
Nomogram for the model predicting an individual patient's VAS pain score. First, each variable is marked on the appropriate scale, and the number of points for each is derived from the "Points" scale at the top of the nomogram. Then the points are totaled, and the total value is marked on the "Total Points" scale. Viewing down, one then derives the predicted pain score on the scale labeled "Linear Predictor." For ease of interpretation not all predictors included in the model are shown in the nomogram. HS = high school, Grad = graduate, Bach = Bachelor's, and SSp = supraspinatus.

of the points for a subject of black race, an SCQ of 10, and a high-school education (~45 points for black race, ~40 points for an SCQ of 10, and 29 points for a high-school education) equals 114 total points, with a predicted VAS pain score of 5.6. When this is compared with a subject of white race who has an SCQ of 2 and a graduate degree (~25 points for white, ~12 points for an SCQ of 2, and ~2 points for a graduate degree equals a total of 39 points, with a predicted VAS pain score of ~3.3, the difference in the predicted VAS pain scores is 2.3, which would be considered a clinically meaningful difference.

Discussion

This cross-sectional study demonstrated no correlation between the level of pain reported by patients and any anatomic measure of rotator cuff tear severity. The level of pain did correlate with nonanatomic features, including increased number of comorbidities, race, and lower education level.

There are several study limitations, the foremost of which is the cross-sectional design using only baseline data from a prospective cohort, which made causal inferences difficult. Other limitations include the potential for selection bias, as patients who were less willing to have surgery may have been more inclined to participate and patients may have self-selected on the basis of their level of symptoms, and performance bias, as some patients may have received medications, acupuncture, or other pain-relieving treatments that we did not examine. Also, generalizability may be limited as many patients presented with a history of trauma and were excluded from the study. Despite these potential limitations, our study included nearly 400 subjects from multiple practices across the United States. Thus, the results of this study may be generalizable to the symptomatic patients with a rotator cuff tear in the United States population within the limitations described above.

Patients with atraumatic orthopaedic conditions typically present with pain, and that pain often correlates with the severity of the disorder¹⁻³. This does not appear to be the case with rotator cuff tears as our data suggest that there is no relationship between pain severity and rotator cuff tear severity. There are abundant data to suggest that the relationship between pain and rotator cuff tears is not robust. Nonoperative treatment of symptomatic, atraumatic, full-thickness rotator cuff tears is successful in approximately 75% of patients⁸⁻¹¹. Patients who have a failed repair of a rotator cuff tear are often pain-free, and their outcomes may be indistinguishable from those in patients with an intact repair⁷. Finally, the prevalence of asymptomatic rotator cuff tears is high. If one uses a conservative estimate that 10% of the population over the age of sixty-five years in the United States has a full-thickness rotator cuff tear⁶, then on the basis of 2010 census data³³ more than four million citizens in the United States have a full-thickness rotator cuff tear. With this very large number as a denominator and the approximately 250,000 rotator cuff surgical procedures performed each year in the United States³⁴ as a numerator, only 6% of patients with a full-thickness rotator cuff tear have surgery each year. These different lines of evidence suggest that many rotator cuff tears are not associated with pain.

Natural history studies have demonstrated that, in some patients, pain may be associated with enlargement of an asymptomatic rotator cuff tear. Yamaguchi et al. studied progression of asymptomatic rotator cuff tears in twenty-three patients with a contralateral symptomatic rotator cuff tear. The asymptomatic shoulder became symptomatic over a period averaging 2.8 years after the initial ultrasound study³⁵. Nine of the twenty-three remained asymptomatic, and fourteen became symptomatic. Of the nine who remained asymptomatic, two had progression of the asymptomatic rotator cuff tear. Of the fourteen who became symptomatic, seven had rotator cuff tear progression. While this study demonstrated that those with tear progression are more likely to develop pain, it also showed that tear progression can develop without pain and that pain can develop without tear progression. The average 10-cm VAS pain score in the symptomatic group was 2.9 cm higher than that in the asymptomatic group.

In the study by Moosmayer et al., fifty patients with an asymptomatic rotator cuff tear and a contralateral symptomatic rotator cuff tear were followed over three years³⁶. Eighteen patients (36%) developed symptoms, with an increase of 3.3 cm in the 10-cm VAS pain score, and the increase in tear size over time was greater in the newly symptomatic group than in the asymptomatic group. Confounding this was the finding that the long head of the biceps tendon developed more pathologic changes in the newly symptomatic group than in the asymptomatic group. What is not known is whether the increase in pain with rotator cuff tear progression diminishes with exercise or with time. One might expect that patients with larger rotator cuff tears would report a longer duration of symptoms, but this is not the case³⁷.

It appears that the majority of rotator cuff tears are asymptomatic. Rotator cuff tears can progress in some patients; this progression may occur without symptoms in some but will be associated with symptoms in many. The amount of pain associated with tear progression (approximately 3 cm on a 10-cm pain VAS) is less than the median value seen in our study (4.4 cm), suggesting that the pain with cuff tear progression may not be great enough to drive patients to seek medical attention. The data also suggest that patients with a rotator cuff tear may develop pain without tear progression, suggesting that other sources of pain, such as the long head of the biceps, may be present^{38,39}.

Indications for surgical repair of chronic, atraumatic rotator cuff tears are not clearly defined and have little uniformity^{12,20,22}, which may explain the widespread geographic variation in rotator cuff surgery rates⁴⁰. Patients in whom a rotator cuff repair has failed report outcome scores that are not significantly different from those for patients whose repair has healed, unless the outcome score includes a large component for strength, in which case those with a healed repair have better scores^{7,11}. Weakness or loss of function may be a better indication for rotator cuff repair than pain in patients with an atraumatic symptomatic rotator cuff tear.

In conclusion, patients who present with shoulder pain without a history of an injury and with MRI evidence of a rotator cuff tear present a dilemma to physicians as the literature lacks high-level evidence to help us make decisions regarding appropriate treatment. In this large cross-sectional

study of patients with an atraumatic, symptomatic full-thickness rotator cuff tear, the patient's pain had no correlation with anatomic measures of the severity of the rotator cuff tear. Because the relationship between pain and the presence of a rotator cuff tear is not robust, when a patient presents with pain as the primary symptom, it may not be accurate to assume that the accompanying rotator cuff tear is responsible for the pain. ■

NOTE: The authors acknowledge the following research personnel from their respective institutions: Vanderbilt University: Brooke Rode, BA; Washington University in St. Louis: Linda Burnworth, Amanda Haas, MA, and Deb Hanson; University of Iowa: Carla Britton, PhD; Hospital for Special Surgery: Samuel Chu, Jessica Ryu, Patrick Grimm, Kaitlyn Lillemoe, and Brian Boyle; Ohio State University: Angela Pedroza, BS; University of California-San Francisco: May Shishido; Orthopaedic Institute: Kari Caspers; and Knoxville Orthopaedic Clinic: Lori Sharp, PA-C, and Jeff Jamigan, PA-C.

Warren R. Dunn, MD, MPH
University of Wisconsin,
Research Park Clinic Sports Medicine Clinic,
621 Science Drive,
Madison, WI 53711

John E. Kuhn, MD, MS
Rosemary Sanders, BA
Vanderbilt University Medical Center,
4200 MCE South Tower,
1215 21st Avenue South,
Nashville, TN 37232.
E-mail address for J.E. Kuhn: j.kuhn@vanderbilt.edu

Qi An, MS
St. Jude Children's Research Hospital,
262 Danny Thomas Place,
Mail Stop 723,
Memphis, TN 38105-3768

Keith M. Baumgarten, MD
Sports Medicine & Shoulder Surgery Orthopedic Institute,
810 East 23rd Street,
Sioux Falls, SD 57117

Julie Y. Bishop, MD
Grant L. Jones, MD
OSU Sports Medicine Center,
2050 Kenny Road, Suite 3300,
Columbus, OH 43221-3502

Robert H. Brophy, MD
Matthew V. Smith, MD
Department of Orthopaedic Surgery,

Washington University School of Medicine,
14532 South Outer Forty Drive,
Chesterfield, MO 63017

Rick W. Wright, MD
Department of Orthopaedic Surgery,
Washington University School of Medicine,
One Barnes-Jewish Hospital Plaza,
11300 West Pavilion,
St. Louis, MO 63110

James L. Carey, MD, MPH
PENN Orthopaedics,
Hospital of the University of Pennsylvania,
Weightman Hall,
33rd and Spruce Streets,
Philadelphia, PA 19104

G. Brian Holloway, MD
Edwin E. Spencer, MD
Shoulder and Elbow Institute Knoxville Orthopaedic Clinic,
260 Fort Sanders West Boulevard,
Knoxville, TN 37922

C. Benjamin Ma, MD
UCSF Sports Medicine,
1500 Owens Street,
San Francisco, CA 94158

Robert G. Marx, MD, MS
Hospital for Special Surgery,
535 East 70th Street,
New York, NY 10021

Eric C. McCarty, MD
CU Sports Medicine,
311 Mapleton Avenue,
Boulder, CO 80304

Sourav K. Poddar, MD
Armando F. Vidal, MD
CU Sports Medicine,
2000 South Colorado Boulevard,
Colorado Center Tower One,
Suite 4500,
Denver, CO 80222

Brian R. Wolf, MD, MS
University of Iowa,
2701 Prairie Meadow Drive,
Iowa City, IA 52242-1088

References

1. Kuroyanagi Y, Nagura T, Matsumoto H, Otani T, Toyama Y, Suda Y. Weight-bearing flexion angle correlates significantly with severity of knee osteoarthritis. *Knee*. 2009 Oct;16(5):371-4. Epub 2009 Feb 11.
2. Neogi T, Felson D, Niu J, Nevitt M, Lewis CE, Aliabadi P, Sack B, Torner J, Bradley L, Zhang Y. Association between radiographic features of knee osteoarthritis and pain: results from two cohort studies. *BMJ*. 2009;339:b2844. Epub 2009 Aug 21.
3. Sadosky AB, Bushmakina AG, Cappelleri JC, Lionberger DR. Relationship between patient-reported disease severity in osteoarthritis and self-reported pain, function and work productivity. *Arthritis Res Ther*. 2010;12(4):R162. Epub 2010 Aug 25.
4. Sher JS, Uribe JW, Posada A, Murphy BJ, Zlatkin MB. Abnormal findings on magnetic resonance images of asymptomatic shoulders. *J Bone Joint Surg Am*. 1995 Jan;77(1):10-5.
5. Yamaguchi K, Tetro AM, Blam O, Evanoff BA, Teeffey SA, Middleton WD. Natural history of asymptomatic rotator cuff tears: a longitudinal analysis of asymptomatic tears detected sonographically. *J Shoulder Elbow Surg*. 2001 May-Jun;10(3):199-203.
6. Reilly P, Macleod I, Macfarlane R, Windley J, Emery RJH. Dead men and radiologists don't lie: a review of cadaveric and radiological studies of rotator cuff tear prevalence. *Ann R Coll Surg Engl*. 2006 Mar;88(2):116-21.
7. Slabaugh MA, Nho SJ, Grumet RC, Wilson JB, Seroyer ST, Frank RM, Romeo AA, Provencher MT, Verma NN. Does the literature confirm superior clinical results in

radiographically healed rotator cuffs after rotator cuff repair? *Arthroscopy*. 2010 Mar;26(3):393-403. Epub 2010 Jan 15.

8. Ainsworth R, Lewis JS. Exercise therapy for the conservative management of full thickness tears of the rotator cuff: a systematic review. *Br J Sports Med*. 2007 Apr;41(4):200-10. Epub 2007 Jan 30.

9. Kijima H, Minagawa H, Nishi T, Kikuchi K, Shimada Y. Long-term follow-up of cases of rotator cuff tear treated conservatively. *J Shoulder Elbow Surg*. 2012. Apr;21(4):491-4. Epub 2012 Jan 21.

10. Kuhn JE, Dunn WR, Sanders R, An Q, Baumgarten KM, Bishop JY, Brophy RH, Carey JL, Holloway BG, Jones GL, Ma CB, Marx RG, McCarty EC, Poddar SK, Smith MV, Spencer EE, Vidal AF, Wolf BR, Wright RW; MOON Shoulder Group. Effectiveness of physical therapy in treating atraumatic full-thickness rotator cuff tears: a multicenter prospective cohort study. *J Shoulder Elbow Surg*. 2013 Oct;22(10):1371-9. Epub 2013 Mar 27.

11. Moosmayer S, Lund G, Seljom U, Svege I, Hennig T, Tariq R, Smith HJ. Comparison between surgery and physiotherapy in the treatment of small and medium-sized tears of the rotator cuff: A randomised controlled study of 103 patients with one-year follow-up. *J Bone Joint Surg Br*. 2010 Jan;92(1):83-91.

12. Oh LS, Wolf BR, Hall MP, Levy BA, Marx RG. Indications for rotator cuff repair: a systematic review. *Clin Orthop Relat Res*. 2007 Feb;455:52-63.

13. Marx RG, Koulouvaris P, Chu SK, Levy BA. Indications for surgery in clinical outcome studies of rotator cuff repair. *Clin Orthop Relat Res*. 2009 Feb;467(2):450-6. Epub 2008 Oct 24.

14. Tashjian RZ. Epidemiology, natural history, and indications for treatment of rotator cuff tears. *Clin Sports Med*. 2012 Oct;31(4):589-604. Epub 2012 Aug 30.

15. Williams GR Jr, Rockwood CA Jr, Bigliani LU, Iannotti JP, Stanwood W. Rotator cuff tears: why do we repair them? *J Bone Joint Surg Am*. 2004 Dec;86-A(12):2764-76.

16. Spencer EE Jr, Dunn WR, Wright RW, Wolf BR, Spindler KP, McCarty E, Ma CB, Jones G, Safran M, Holloway GB, Kuhn JE; Shoulder Multicenter Orthopaedic Outcomes Network. Interobserver agreement in the classification of rotator cuff tears using magnetic resonance imaging. *Am J Sports Med*. 2008 Jan;36(1):99-103. Epub 2007 Oct 11.

17. Kuhn JE, Dunn WR, Ma B, Wright RW, Jones G, Spencer EE, Wolf B, Safran M, Spindler KP, McCarty E, Kelly B, Holloway B; Multicenter Orthopaedic Outcomes Network-Shoulder (MOON Shoulder Group). Interobserver agreement in the classification of rotator cuff tears. *Am J Sports Med*. 2007 Mar;35(3):437-41. Epub 2007 Jan 31.

18. Baumgarten KM, Carey JL, Abboud JA, Jones GL, Kuhn JE, Wolf BR, Brophy RH, Cox CL, Wright RW, Vidal AF, Ma CB, McCarty EC, Holloway GB, Spencer EE Jr, Dunn WR. Reliability of determining and measuring acromial enthesophytes. *HSS J*. 2011 Oct;7(3):218-22. Epub 2011 Jul 13.

19. Baumgarten KM, Vidal AF, Wright RW. Rotator cuff repair rehabilitation: A level I and II systematic review. *Sports Health*. 2009 Mar;1(2):125-30.

20. Wolf BR, Dunn WR, Wright RW. Indications for repair of full-thickness rotator cuff tears. *Am J Sports Med*. 2007 Jun;35(6):1007-16. Epub 2007 Mar 2.

21. Kuhn JE. Exercise in the treatment of rotator cuff impingement: a systematic review and a synthesized evidence-based rehabilitation protocol. *J Shoulder Elbow Surg*. 2009 Jan-Feb;18(1):138-60. Epub 2008 Oct 2.

22. Dunn WR, Schackman BR, Walsh C, Lyman S, Jones EC, Warren RF, Marx RG. Variation in orthopaedic surgeons' perceptions about the indications for rotator cuff surgery. *J Bone Joint Surg Am*. 2005 Sep;87(9):1978-84.

23. Ware JE Jr, Kosinski M, Keller SD. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care*. 1996 Mar;34(3):220-33.

24. Richards RR, An KN, Bigliani LU, Friedman RJ, Gartsman GM, Gristina AG, Iannotti JP, Mow VC, Sidles JA, Zuckerman JD. A standardized method for the as-

essment of shoulder function. *J Shoulder Elbow Surg*. 1994 Nov;3(6):347-52. Epub 2009 Feb 13.

25. Kirkley A, Alvarez C, Griffin S. The development and evaluation of a disease-specific quality-of-life questionnaire for disorders of the rotator cuff: The Western Ontario Rotator Cuff Index. *Clin J Sport Med*. 2003 Mar;13(2):84-92.

26. Williams GN, Gangel TJ, Arciero RA, Uhorchak JM, Taylor DC. Comparison of the Single Assessment Numeric Evaluation method and two shoulder rating scales. Outcomes measures after shoulder surgery. *Am J Sports Med*. 1999 Mar-Apr;27(2):214-21.

27. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum*. 2003 Apr 15;49(2):156-63.

28. Brophy RH, Beauvais RL, Jones EC, Cordasco FA, Marx RG. Measurement of shoulder activity level. *Clin Orthop Relat Res*. 2005 Oct;439(439):101-8.

29. Patte D. Classification of rotator cuff lesions. *Clin Orthop Relat Res*. 1990 May;254:81-6.

30. Zanetti M, Gerber C, Hodler J. Quantitative assessment of the muscles of the rotator cuff with magnetic resonance imaging. *Invest Radiol*. 1998 Mar;33(3):163-70.

31. Baigent C, Harrell FE, Buyse M, Emberson JR, Altman DG. Ensuring trial validity by data quality assurance and diversification of monitoring methods. *Clin Trials*. 2008;5(1):49-55.

32. Tashjian RZ, Deloach J, Porucznik CA, Powell AP. Minimal clinically important differences (MCID) and patient acceptable symptomatic state (PASS) for visual analog scales (VAS) measuring pain in patients treated for rotator cuff disease. *J Shoulder Elbow Surg*. 2009 Nov-Dec;18(6):927-32. Epub 2009 Jun 16.

33. Werner CA. The older population: 2010. US Census Bureau. 2011 Nov. <http://www.census.gov/prod/cen2010/briefs/c2010br-09.pdf>. Accessed 2013 Dec 31.

34. McCormick H. ArthroCare closes Opus Medical acquisition. 2004 Nov 22. Orthopaedic and Dental Industry News. http://www.healthpointcapital.com/research/2004/11/22/arthrocare_closes_opus_medical_acquisition/. Accessed 2013 Dec 31.

35. Yamaguchi K, Tetro AM, Blam O, Evanoff BA, Teefey SA, Middleton WD. Natural history of asymptomatic rotator cuff tears: a longitudinal analysis of asymptomatic tears detected sonographically. *J Shoulder Elbow Surg*. 2001 May-Jun;10(3):199-203.

36. Moosmayer S, Tariq R, Stiris M, Smith HJ. The natural history of asymptomatic rotator cuff tears: a three-year follow-up of fifty cases. *J Bone Joint Surg Am*. 2013 Jul 17;95(14):1249-55.

37. The MOON Shoulder Group: Unruh KP, Kuhn JE, Sanders R, An Q, Baumgarten KM, Bishop JY, Brophy RH, Carey JL, Holloway BG, Jones GL, Ma BC, Marx RG, McCarty EC, Poddar SK, Smith MV, Spencer EE, Vidal AF, Wolf BR, Wright RW, Dunn WR. The duration of symptoms does not correlate with rotator cuff tear severity or other patient-related features. A cross-sectional study of patients with atraumatic, full-thickness rotator cuff tears. *J Shoulder Elbow Surg*. 2014 Jan 8. pii: S1058-2746(13)00527-2. Epub ahead of print.

38. Boileau P, Baqué F, Valerio L, Ahrens P, Chuinard C, Trojani C. Isolated arthroscopic biceps tenotomy or tenodesis improves symptoms in patients with massive irreparable rotator cuff tears. *J Bone Joint Surg Am*. 2007 Apr;89(4):747-57.

39. Szabó I, Boileau P, Walch G. The proximal biceps as a pain generator and results of tenotomy. *Sports Med Arthrosc*. 2008 Sep;16(3):180-6.

40. Vitale MG, Krant JJ, Gelijns AC, Heitjan DF, Arons RR, Bigliani LU, Flatow EL. Geographic variations in the rates of operative procedures involving the shoulder, including total shoulder replacement, humeral head replacement, and rotator cuff repair. *J Bone Joint Surg Am*. 1999 Jun;81(6):763-72.