

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2007

The Par-1/MARK family of protein kinases: From polarity to metabolism

Jonathan Hurov

Beth Israel Deaconess Medical Center

Helen Piwnica-Worms

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Hurov, Jonathan and Piwnica-Worms, Helen, "The Par-1/MARK family of protein kinases: From polarity to metabolism." *Cell Cycle*. 6, 16. 1966-1969. (2007).

https://digitalcommons.wustl.edu/open_access_pubs/3026

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Perspective

The Par-1/MARK Family of Protein Kinases

From Polarity to Metabolism

Jonathan Hurov¹

Helen Piwnica-Worms^{2-4,*}

¹Division of Signal Transduction; Beth Israel Deaconess Medical Center; Boston, Massachusetts USA

²Howard Hughes Medical Institute; ³Department of Cell Biology and Physiology;

⁴Department of Internal Medicine; Washington University School of Medicine; St. Louis, Missouri USA

*Correspondence to: Helen Piwnica-Worms; Howard Hughes Medical Institute; Department of Cell Biology and Physiology; Department of Internal Medicine; Washington University School of Medicine; 660 South Euclid Avenue; Box 8228; St. Louis 63110 Missouri USA; Email: hpiwnica@cellbiology.wustl.edu

Original manuscript submitted: 06/07/07

Manuscript accepted: 06/08/07

Previously published online as a *Cell Cycle* E-publication:
<http://www.landesbioscience.com/journals/cc/article/4576>

KEY WORDS

adiposity, AMPK, aPKC, LKB, metabolism, polarity, Par

ABBREVIATIONS

Par partitioning defective
aPKC atypical protein kinase C
AMPK AMP-activated protein kinase
Glut4 glucose transporter 4

ACKNOWLEDGEMENTS

Helen Piwnica-Worms is an investigator of the Howard Hughes Medical Institute.

ABSTRACT

The Par-1 protein kinases are conserved from yeast to man and belong to a subfamily of kinases that includes the energy sensor and metabolic regulator, AMPK. Par-1 is regulated by LKB1 and atypical PKC and has been shown in multiple organisms and cell types to be critical for regulation of cellular polarity. Recent studies using knockout mice have revealed several surprising physiological functions for Par-1b/MARK2/EMK1. Our recent study shows that Par-1b regulates metabolic rate, adiposity and insulin sensitivity. This is the first study to implicate these kinases in metabolic functions akin to those previously defined for AMPK. Conversely, another series of recent publications now implicate AMPK in regulation of polarity. Here we discuss the metabolic phenotype seen in Par-1b deficient mice and the synthesis of several findings that link Par-1 and AMPK to a degree that has not been previously appreciated.

INTRODUCTION

The prototypical Par-1 kinase was identified in a seminal study published in 1988 designed to identify regulators of early embryonic polarity in *C. elegans*.¹ A series of subsequent studies by numerous groups demonstrated that Par-1, a serine/threonine protein kinase, is one of several evolutionarily conserved proteins (Par-1, Par-3/ASIP, Par-4/LKB1, Par-5/14-3-3, Par-6 and atypical PKC/PKC-3) required for cellular polarity not only in worms but also in flies, frogs and mammals.²⁻¹⁶ This body of work has been reviewed elsewhere.¹⁷⁻²⁰ Studies done in the context of cellular polarity demonstrated that Par-1 mediates at least some of its effects by phosphorylating Par-3 and the microtubule associated protein Tau.²¹⁻²⁵ Numerous other potential Par-1 substrates have been identified including Cdc25C, KSR, Pkp2, Class II HDAC,Dlg and Rab11-FIP.²⁶⁻³² Interestingly, phosphorylation of several Par-1 substrates leads to the generation of phospho-dependent 14-3-3 binding. Par-1 substrates that fall into this category include Par-3, Cdc25C, KSR, Pkp2 and Class II HDAC. Both LKB1 (also known as Par-4) and atypical PKC λ/ι (or PKC-3 in *C. elegans*) regulate Par-1. LKB1 phosphorylates Par-1 and the related AMP activated kinase AMPK on an activation loop Thr residue. This modification is required for Par-1 activity.^{33,34} LKB1 serves as a master regulator of cellular polarity, at least in part, by activating the Par-1 kinases.¹⁶ Atypical PKC (aPKC) phosphorylates Par-1 to regulate its localization and kinase activity.³⁵⁻³⁷ TAO-1/MARKK and GSK-3 β have also been implicated as upstream regulators of Par-1.^{38,39}

METABOLIC FUNCTIONS OF Par-1b/MARK2/EMK

Physiological functions of the mammalian Par-1 kinases have been revealed using targeted gene knockout approaches in mice. The mammalian Par-1 family is comprised of four members that go by several names (Par-1a/MARK3/C-TAK1, Par-1b/MARK2/EMK, Par-1c/MARK1 and Par-1d). Four studies have been published using two independently derived mouse lines null for Par-1b/MARK2/EMK.⁴⁰⁻⁴³ These studies implicate Par-1b in a diverse set of physiological processes, including fertility, immune system homeostasis, learning and memory and growth and metabolism.

Our most recent study identifies a role for Par-1b in the regulation of metabolism.⁴³ Par-1b null mice are growth retarded (~20% reduced body mass relative to wild-type) as early as E13.5 and this growth retardation continues throughout the lifetime of Par-1b null mice.⁴³ Our results and those of Bessone et al. (1999) showing decreased serum IGF-1 levels in Par-1b null mice, provide a reasonable explanation for the observed

pre- and post-natal growth retardation. Although a growth hormone deficiency might explain postnatal growth differences, there is no evidence for a role of growth hormone (GH) in mammalian embryonic growth, suggesting that Par-1b deficiency leads to GH-independent growth defects. Although accurate determination of GH production is technically difficult, neither our study nor that of Bessone et al. (1999) detected altered serum GH levels in Par-1b null mice.

Morphometric analyses of Par-1b null mice revealed that although most tissues are proportionately smaller, knockout mice accumulate disproportionate decreases in adipose tissue (representing ~14% body fat in null mice versus ~23% in wild-type mice at twelve weeks of age). This differential is slightly increased with age (18% body fat in null mice versus 35% fat in wild-type mice at one year of age, $n = 10$ females per genotype, $p = 0.002$) (unpublished data). In addition to having reduced adiposity, Par-1b null mice are resistant to weight gain when placed on a high fat diet and are at the same time hyperphagic, eating twice that of their wild-type littermates. A likely explanation for these observations is the finding that Par-1b null mice are hypermetabolic. Thus, resistance to weight gain due to high fat diet or increased caloric intake is due to increased metabolic rate. Although the molecular mechanism for these metabolic changes is not clear, the data indicates that loss of Par-1b either directly or indirectly increases mitochondrial function in adipose tissue. Intriguingly, these perturbations are also accompanied by insulin hypersensitivity and improved glucose tolerance—possibly the result of compensatory changes that arise due to a chronic hypermetabolic state. Analysis of the relative levels of glucose uptake in muscle and fat indicates that adipose tissue is most dramatically affected in the absence of Par-1b. White and brown fat of Par-1b null mice exhibit increased glucose uptake in both basal and insulin-stimulated states.

Another possible explanation for enhanced insulin sensitivity and glucose uptake in Par-1b null adipose tissue is that Par-1b regulates GLUT4-mediated glucose uptake. Par-3/ASIP studies in 3T3L1 cells indicate that this protein (a downstream target of Par-1) can inhibit insulin induced glucose uptake⁴⁴ aPKC, a negative regulator of Par-1, has been shown to regulate insulin triggered glucose uptake in multiple studies.^{44,45} It is unclear at this point how Par-3/ASIP or aPKC act to regulate glucose uptake. It is conceivable that aPKC influences glucose uptake via regulation of Par-1. Par-1 has also been implicated in the regulation of vesicular trafficking via the exocyst, which in turn has been shown to play a critical role in the transport of insulin-responsive Glut4 vesicles.⁴⁶⁻⁴⁸ Regulation of Rab11-mediated functions might provide a mechanistic explanation for this model.^{32,49} Thus, Par-1 may be involved in GLUT4-mediated glucose uptake via a mechanism(s) that involves Par-3, aPKC and/or the exocyst.

An argument against a direct role for Par-1b in early signaling events that control Glut4-mediated glucose uptake is our finding that insulin receptor proximal signaling (IRS-1 and AKT phosphorylation, phosphatidylinositol 3-kinase recruitment) is slightly decreased in adipose tissue from Par-1b null mice. Furthermore, Zhou et al. (2004) reported that knockdown of Par-1a and/or Par-1b in 3T3L1 cells by siRNA-treatment, has no effect on basal- or insulin-stimulated glucose uptake.⁵⁰ If knockdown was sufficient to block the function of both Par-1a and Par-1b in the 3T3L1 system and if there is not compensation by the other two Par-1 family members, then this data provides an argument against a direct role for Par-1 in Glut4-mediated glucose uptake. Even if Par-1 regulates the activity of Glut4 downstream of AKT, this function would not completely

explain our observation of hypermetabolism and reduced adiposity in the Par-1b null mice because adipose-specific overexpression of Glut4 alone leads to enhanced glucose uptake in combination with increased adiposity (not decreased adiposity as observed in Par-1b deficiency).⁵¹⁻⁵³ Future studies using tissue-specific deletion of Par-1b in adipose tissue will elucidate the contribution made by adipose tissue to these phenotypes.

There is also a significant body of literature describing a role for Par-1 in neuronal cell polarity.^{31,54-56} Several interesting possibilities, including regulation of the hypothalamic-pituitary-adrenergic axis of the neuronal system by Par-1b, might explain the observed metabolic changes in Par-1b null mice.⁵⁷ Again, tissue-specific deletion of Par-1b in the nervous system should clarify how this compartment contributes to the observed phenotypes.

AMPK AND PAR-1: PARTNERS IN POLARITY AND METABOLISM?

Based on sequence similarities (50% identity across their kinase domains), AMPK and Par-1 are closely related members of a subfamily that also includes BRSK1/2, QIK and SIK.⁵⁸ Conservation of the kinase domains of these proteins suggests similar phosphorylation-site preferences. Alignment of several known substrates indicates that Par-1 and AMPK prefer to phosphorylate a serine residue when Leu/Ile/Met, Arg/Lys, and Leu are present in the -5, -3 and +4 positions, respectively (Fig. 1). In addition to sharing a common consensus phosphorylation motif, both AMPK and Par-1 family members are activated by LKB1.³⁴ Interestingly, recent studies indicate that Par-1b and AMPK have overlapping functions in vivo. As mentioned above, Par-1 has historically been associated with regulating polarity, while AMPK has been studied for many years in the context of energy sensing and metabolism. The metabolic

Par-1	
LRRTESVPSD	hKSR (Ser164)
LYRSPSPEN	hCdc25C (Ser216)
LHRTSSVPEY	hPKP2a (Ser82)
MVRRSSDPNL	dPar-3 (Ser151)
MKKSSSLESL	dPar-3 (Ser1085)
LSSAHMSMDL	hRab11-FIP2 (Ser227)
KSKIGSTENL	hTau (Ser262)
AMPK	
IRSSMSGHLH	hACC1 (Ser78)
LNRTSSDSAL	hTORC2 (Ser171)
RMRRNSFTPL	hPFK2 (Ser466)
LSKSSSSPEL	hTSC2 (Ser1387)
RGRSLSVTSL	hGS (Ser8)
RLRSESIIFI	hNOS (Ser1417)
MIHNRSKINL	hHMG-CoA (Ser872)

Figure 1. Par-1 and AMPK substrates share a common phosphorylation consensus sequence. Sequences inclusive of and surrounding the phosphorylation site of several Par-1 and AMPK substrates are aligned. Phosphorylated serine residues are highlighted in blue and preferred residues in the -5, -3 and +4 positions are highlighted in yellow. h, human; d, Drosophila; KSR, kinase suppressor of Ras; PKP2, Plakophilin, ACC1, acetyl CoA carboxylase; TORC2, transducer of regulated cAMP response element-binding protein (CREB); PFK2, phospho-fructo-kinase; TSC2, tuberous sclerosis; GS, glycogen synthase; NOS, nitric oxide synthase; HMG-CoA, 3-hydroxy-3-methylglutaryl-coenzyme A reductase.

functions of AMPK have been well-reviewed elsewhere.^{59,60} These functions include inhibition of fatty acid, glycogen and protein synthesis and activation of glucose uptake (in skeletal muscle) and glycolysis in response to cellular energy stress (increased AMP levels). Our study now elucidates a role for Par-1b in metabolic regulation, potentially via an adipose- and/or a neuronal-specific mechanism(s). At the same time, four recent studies have uncovered a role for AMPK in the regulation of polarity.⁶¹⁻⁶⁴ It is therefore tempting to speculate that Par-1 and AMPK share either a common and/or complementary set of substrates that act to regulate similar biological processes.

CONCLUDING REMARKS

Recent studies using Par-1b null mice demonstrate that mammalian Par-1b/MARK2 is required for multiple physiological processes that could not have been predicted from previous studies conducted in vitro. Our recent work indicates that Par-1b is important for metabolic regulation, and in particular, adiposity. Perhaps not surprising, a combination of studies has now drawn several parallels between substrate specificities, upstream regulators and physiological functions of Par-1 and AMPK. Although their kinase domains are clearly related in sequence, the notion that the functions of Par-1 and AMPK in polarity and metabolism, respectively, might be overlapping, was not predicted. Many exciting questions remain to be answered, including the identification of downstream substrates and upstream regulators of Par-1 and AMPK with respect to their newly defined functions as well as the identification of tissues that are relevant to the observed metabolic defects in Par-1b null mice.

References

- Kemphues KJ, Priess JR, Morton DG, Cheng N. Identification of genes required for cytoplasmic localization in early embryos of *C. elegans*. *Cell* 1988; 52:311-20.
- Guo S, Kemphues KJ. *Par-1*, a gene required for establishing polarity in *C. elegans* embryos, encodes a putative ser/thr kinase that is asymmetrically distributed. *Cell* 1995; 81:611-20.
- Bayraktar J, Zygmunt D, Carthew RW. Par-1 kinase establishes cell polarity and functions in Notch signaling in the *Drosophila* embryo. *J Cell Sci* 2006; 119:711-21.
- Bohm H, Brinkman V, Drab M, Henske A, Kurczak TV. Mammalian Homologues of *C-elegans* polarization gene product *PAR-1* are asymmetrically localized in epithelial cells and may influence their polarity. *Curr Biol* 1997; 7:603-6.
- Cox DN, Lu B, Sun TQ, Williams LT, Jan YN. *Drosophila* par-1 is required for oocyte differentiation and microtubule organization. *Curr Biol* 2001; 11:75-87.
- Kusakabe M, Nishida E. The polarity-inducing kinase Par-1 controls *Xenopus* gastrulation in cooperation with 14-3-3 and aPKC. *EMBO J* 2004; 23:4190-201.
- Lin D, Edwards AS, Fawcett JR, Mbamalu G, Scott JD, Pawson T. A mammalian PAR-3-PAR-6 complex implicated in Cdc42/Rac1 and aPKC signalling and cell polarity. *Nat Cell Biol* 2000; 2:540-7.
- Hurd TW, Gao L, Roh MH, Macara IG, Margolis B. Direct interaction of two polarity complexes implicated in epithelial tight junction assembly. *Nat Cell Biol* 2003; 5:137-42.
- Huynh JR, Petronczki M, Knoblich JA, St Johnston D. Bazooka and PAR-6 are required with PAR-1 for the maintenance of oocyte fate in *Drosophila*. *Curr Biol* 2001; 11:901-6.
- Huynh JR, Shulman JM, Benton R, St Johnston D. PAR-1 is required for the maintenance of oocyte fate in *Drosophila*. *Development* 2001; 128:1201-9.
- Joberty G, Petersen C, Gao L, Macara IG. The cell-polarity protein Par6 links Par3 and atypical protein kinase C to Cdc42. *Nat Cell Biol* 2000; 2:531-9.
- Shulman JM, Benton R, St Johnston D. The *Drosophila* homolog of *C. elegans* PAR-1 organizes the oocyte cytoskeleton and directs oskar mRNA localization to the posterior pole. *Cell* 2000; 101:377-88.
- Tabuse Y, Izumi Y, Piano F, Kemphues KJ, Miwa J, Ohno S. Atypical protein kinase C cooperates with PAR-3 to establish embryonic polarity in *Caenorhabditis elegans*. *Development* 1998; 125:3607-14.
- Tomanek P, Piano F, Riechmann V, Gunsalus KC, Kemphues KJ, Ephrussi A. A *Drosophila melanogaster* homologue of *Caenorhabditis elegans* par-1 acts at an early step in embryonic-axis formation. *Nat Cell Biol* 2000; 2:458-60.
- Martin SG, St Johnston D. A role for *Drosophila* LKB1 in anterior-posterior axis formation and epithelial polarity. *Nature* 2003; 421:379-84.
- Baas AF, Kuipers J, van der Wel NN, Batlle E, Koerten HK, Peters PJ, Clevers HC. Complete polarization of single intestinal epithelial cells upon activation of LKB1 by STRAD. *Cell* 2004; 116:457-66.
- Shin K, Fogg VC, Margolis B. Tight junctions and cell polarity. *Annu Rev Cell Dev Biol* 2006; 22:207-35.
- Arimura N, Kaibuchi K. Neuronal polarity: From extracellular signals to intracellular mechanisms. *Nat Rev Neurosci* 2007; 8:194-205.
- Nance J. PAR proteins and the establishment of cell polarity during *C. elegans* development. *Bioessays* 2005; 27:126-35.
- Krummel MF, Macara I. Maintenance and modulation of T cell polarity. *Nat Immunol* 2006; 7:1143-9.
- Cohen D, Brennwald PJ, Rodriguez-Boulton E, Musch A. Mammalian PAR-1 determines epithelial lumen polarity by organizing the microtubule cytoskeleton. *J Cell Biol* 2004; 164:717-27.
- Doerflinger H, Benton R, Shulman JM, St Johnston D. The role of PAR-1 in regulating the polarised microtubule cytoskeleton in the *Drosophila* follicular epithelium. *Development* 2003; 130:3965-75.
- Betschinger J, Mechtler K, Knoblich JA. The Par complex directs asymmetric cell division by phosphorylating the cytoskeletal protein Lgl. *Nature* 2003; 422:326-30.
- Benton R, St Johnston D. *Drosophila* PAR-1 and 14-3-3 inhibit Bazooka/PAR-3 to establish complementary cortical domains in polarized cells. *Cell* 2003; 115:691-704.
- Drewe G, Ebner A, Preuss U, Mandelkow EM, Mandelkow E. MARK, a novel family of protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule disruption. *Cell* 1997; 89:297-308.
- Ogg S, Gabrielli B, Piwnica-Worms H. Purification of a serine kinase that associates with and phosphorylates human Cdc25C on serine 216. *J Biol Chem* 1994; 269:30461-9.
- Peng CY, Graves PR, Ogg S, Thoma RS, Byrnes MJ, Wu Z, Stephenson M, Piwnica-Worms H. C-TAK1 protein kinase phosphorylates human Cdc25C on serine 216 and promotes 14-3-3 binding. *Cell Growth Differ* 1998; 9:197-208.
- Muller J, Ory S, Copeland T, Piwnica-Worms H, Morrison DK. C-TAK1 regulates Ras signaling by phosphorylating the MAPK scaffold, KSR1. *Mol Cell* 2001; 8:983-93.
- Muller J, Ritt DA, Copeland TD, Morrison DK. Functional analysis of C-TAK1 substrate binding and identification of PKP2 as a new C-TAK1 substrate. *EMBO J* 2003; 22:4431-42.
- Dequiedt F, Martin M, Von Blume J, Vertommen D, Lecomte E, Mari N, Heinen MF, Bachmann M, Twizere JC, Huang MC, Rider MH, Piwnica-Worms H, Seufferlein T, Kettmann R. New role for hPar-1 kinases EMK and C-TAK1 in regulating localization and activity of class IIa histone deacetylases. *Mol Cell Biol* 2006; 26:7086-102.
- Zhang Y, Guo H, Kwan H, Wang JW, Kosek J, Lu B. PAR-1 kinase phosphorylates Dlg and regulates its postsynaptic targeting at the *Drosophila* neuromuscular junction. *Neuron* 2007; 53:201-15.
- Ducharme NA, Hales CM, Lapierre LA, Ham AJ, Oztan A, Apodaca G, Goldenring JR. MARK2/EMK1/Par-1B phosphorylation of Rab11-family interacting protein 2 is necessary for the timely establishment of polarity in Madin-Darby canine kidney cells. *Mol Biol Cell* 2006; 17:3625-37.
- Lizcano JM, Goransson O, Toth R, Deak M, Morrice NA, Boudeau J, Hawley SA, Udd L, Makela TP, Hardie DG, Alessi DR. LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. *EMBO J* 2004; 23:833-43.
- Alessi DR, Sakamoto K, Bayasas JR. Lkb1-dependent signaling pathways. *Annu Rev Biochem* 2006; 75:137-63.
- Hurov JB, Watkins JL, Piwnica-Worms H. Atypical PKC phosphorylates PAR-1 kinases to regulate localization and activity. *Curr Biol* 2004; 14:736-41.
- Suzuki A, Hirata M, Kamimura K, Maniwa R, Yamanaka T, Mizuno K, Kishikawa M, Hirose H, Amano Y, Izumi N, Miwa Y, Ohno S. aPKC acts upstream of PAR-1b in both the establishment and maintenance of mammalian epithelial polarity. *Curr Biol* 2004; 14:1425-35.
- Vaccari T, Rabouille C, Ephrussi A. The *Drosophila* PAR-1 spacer domain is required for lateral membrane association and for polarization of follicular epithelial cells. *Curr Biol* 2005; 15:255-61.
- Timm T, Li XY, Biernat J, Jiao J, Mandelkow E, Vandekerckhove J, Mandelkow EM. MARKK, a Ste20-like kinase, activates the polarity-inducing kinase MARK/PAR-1. *EMBO J* 2003; 22:5090-101.
- Kosuga S, Tashiro E, Kajioka T, Ueki M, Shimizu Y, Imoto M. GSK-3beta directly phosphorylates and activates MARK2/PAR-1. *J Biol Chem* 2005; 280:42715-22.
- Bessone S, Vidal F, Le Bouc Y, Epelbaum J, Bluet-Pajot MT, Darmon M. EMK protein kinase-null mice: Dwarfism and hypofertility associated with alterations in the somatotrophic and prolactin pathways. *Developmental Biology* 1999; 214:87-101.
- Hurov JB, Stappenbeck TS, Zmasek CM, White LS, Ranganath SH, Russell JH, Chan AC, Murphy KM, Piwnica-Worms H. Immune system dysfunction and autoimmune disease in mice lacking *Emk* (*Par-1*) protein kinase. *Mol Cell Biol* 2001; 21:3206-19.
- Segu L, Pascaud A, Costet P, Darmon M, Buhot MC. Impairment of spatial learning and memory in *ELKL* Motif Kinase1 (*EMK1/MARK2*) knockout mice. *Neurobiol Aging* 2006.
- Hurov JB, Huang M, White LS, Lennarz J, Choi CS, Cho YR, Kim HJ, Prior JL, Piwnica-Worms D, Cantley LC, Kim JK, Shulman GI, Piwnica-Worms H. Loss of the Par-1b/MARK2 polarity kinase leads to increased metabolic rate, decreased adiposity, and insulin hypersensitivity in vivo. *Proc Natl Acad Sci USA* 2007; 104:5680-5.
- Kotani K, Ogawa W, Hashimoto M, Onishi T, Ohno S, Kasuga M. Inhibition of insulin-induced glucose uptake by atypical protein kinase C isotype-specific interacting protein in 3T3-L1 adipocytes. *J Biol Chem* 2000; 275:26390-5.
- Faresse RV, Sajjan MB, Standaert ML. Atypical protein kinase C in insulin action and insulin resistance. *Biochem Soc Trans* 2005; 33:350-3.
- Cohen D, Rodriguez-Boulton E, Musch A. Par-1 promotes a hepatic mode of apical protein trafficking in MDCK cells. *Proc Natl Acad Sci USA* 2004; 101:13792-7.

47. Ewart MA, Clarke M, Kane S, Chamberlain LH, Gould GW. Evidence for a role of the exocyst in insulin-stimulated Glut4 trafficking in 3T3-L1 adipocytes. *J Biol Chem* 2005; 280:3812-6.
48. Inoue M, Chang L, Hwang J, Chiang SH, Saltiel AR. The exocyst complex is required for targeting of Glut4 to the plasma membrane by insulin. *Nature* 2003; 422:629-33.
49. Kessler A, Tomas E, Immler D, Meyer HE, Zorzano A, Eckel J. Rab11 is associated with GLUT4-containing vesicles and redistributes in response to insulin. *Diabetologia* 2000; 43:1518-27.
50. Zhou QL, Park JG, Jiang ZY, Holik JJ, Mitra P, Semiz S, Guilherme A, Powelka AM, Tang X, Virbasius J, Czech MP. Analysis of insulin signalling by RNAi-based gene silencing. *Biochem Soc Trans* 2004; 32:817-21.
51. Tozzo E, Shepherd PR, Gnudi L, Kahn BB. Transgenic GLUT-4 overexpression in fat enhances glucose metabolism: Preferential effect on fatty acid synthesis. *Am J Physiol* 1995; 268:E956-64.
52. Carvalho E, Schellhorn SE, Zabolotny JM, Martin S, Tozzo E, Peroni OD, Houseknecht KL, Mundt A, James DE, Kahn BB. GLUT4 overexpression or deficiency in adipocytes of transgenic mice alters the composition of GLUT4 vesicles and the subcellular localization of *GLUT4* and insulin-responsive aminopeptidase. *J Biol Chem* 2004; 279:21598-605.
53. Shepherd PR, Gnudi L, Tozzo E, Yang H, Leach F, Kahn BB. Adipose cell hyperplasia and enhanced glucose disposal in transgenic mice overexpressing GLUT4 selectively in adipose tissue. *J Biol Chem* 1993; 268:22243-6.
54. Biernat J, Wu YZ, Timm T, Zheng-Fischhofer Q, Mandelkow E, Meijer L, Mandelkow EM. Protein kinase MARK/PAR-1 is required for neurite outgrowth and establishment of neuronal polarity. *Mol Biol Cell* 2002; 13:4013-28.
55. Chen YM, Wang QJ, Hu HS, Yu PC, Zhu J, Drewes G, Pownica-Worms H, Luo ZG. Microtubule affinity-regulating kinase 2 functions downstream of the PAR-3/PAR-6/atypical PKC complex in regulating hippocampal neuronal polarity. *Proc Natl Acad Sci USA* 2006; 103:8534-9.
56. Nishimura I, Yang Y, Lu B. PAR-1 kinase plays an initiator role in a temporally ordered phosphorylation process that confers tau toxicity in *Drosophila*. *Cell* 2004; 116:671-82.
57. Kim H, Pennisi PA, Gavrilova O, Pack S, Jou W, Setser-Portas J, East-Palmer J, Tang Y, Manganiello VC, Leroith D. Effect of adipocyte beta3-adrenergic receptor activation on the type 2 diabetic *MDR* mice. *Am J Physiol Endocrinol Metab* 2006; 290:E1227-36.
58. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. The protein kinase complement of the human genome. *Science* 2002; 298:1912-34.
59. Long YC, Zierath JR. AMP-activated protein kinase signaling in metabolic regulation. *J Clin Invest* 2006; 116:1776-83.
60. Towler MC, Hardie DG. AMP-activated protein kinase in metabolic control and insulin signaling. *Circ Res* 2007; 100:328-41.
61. Zhang L, Li J, Young LH, Caplan MJ. AMP-activated protein kinase regulates the assembly of epithelial tight junctions. *Proc Natl Acad Sci USA* 2006; 103:17272-7.
62. Zheng B, Cantley LC. Regulation of epithelial tight junction assembly and disassembly by AMP-activated protein kinase. *Proc Natl Acad Sci USA* 2007; 104:819-22.
63. Lee JH, Koh H, Kim M, Kim Y, Lee SY, Karess RE, Lee SH, Shong M, Kim JM, Kim J, Chung J. Energy-dependent regulation of cell structure by AMP-activated protein kinase. *Nature* 2007.
64. Mirouse V, Swick LL, Kazgan N, St Johnston D, Brenman JE. LKB1 and AMPK maintain epithelial cell polarity under energetic stress. *J Cell Biol* 2007; 177:387-92.