

Washington University School of Medicine

Digital Commons@Becker

---

Open Access Publications

---

2014

## Cross talk between the Cell Wall Integrity and Cyclic AMP/Protein Kinase A pathways in *Cryptococcus neoformans*

Maureen J. Donlin  
*Saint Louis University*

Rajendra Upadhya  
*Washington University School of Medicine in St. Louis*

Kimberly J. Gerik  
*Washington University School of Medicine in St. Louis*

Woei Lam  
*Washington University School of Medicine in St. Louis*

Laura G. VanArendonk  
*Washington University School of Medicine in St. Louis*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Donlin, Maureen J.; Upadhya, Rajendra; Gerik, Kimberly J.; Lam, Woei; VanArendonk, Laura G.; Specht, Charles A.; Sharma, Neil K.; and Lodge, Jennifer K., "Cross talk between the Cell Wall Integrity and Cyclic AMP/Protein Kinase A pathways in *Cryptococcus neoformans*." *mBio*. 5, 4. e01573-14. (2014).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/3144](https://digitalcommons.wustl.edu/open_access_pubs/3144)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

---

## Authors

Maureen J. Donlin, Rajendra Upadhya, Kimberly J. Gerik, Woei Lam, Laura G. VanArendonk, Charles A. Specht, Neil K. Sharma, and Jennifer K. Lodge



## Cross Talk between the Cell Wall Integrity and Cyclic AMP/Protein Kinase A Pathways in *Cryptococcus neoformans*

Maureen J. Donlin, Rajendra Upadhyaya, Kimberly J. Gerik, et al.  
2014. Cross Talk between the Cell Wall Integrity and Cyclic AMP/Protein Kinase A Pathways in *Cryptococcus neoformans*. mBio 5(4): .  
doi:10.1128/mBio.01573-14.

---

Updated information and services can be found at:  
<http://mbio.asm.org/content/5/4/e01573-14.full.html>

---

**SUPPLEMENTAL MATERIAL**

<http://mbio.asm.org/content/5/4/e01573-14.full.html#SUPPLEMENTAL>

**REFERENCES**

This article cites 61 articles, 34 of which can be accessed free at:  
<http://mbio.asm.org/content/5/4/e01573-14.full.html#ref-list-1>

**CONTENT ALERTS**

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more>>](#)

---

---

Information about commercial reprint orders: <http://mbio.asm.org/misc/reprints.xhtml>

Information about Print on Demand and other content delivery options:

<http://mbio.asm.org/misc/contentdelivery.xhtml>

To subscribe to another ASM Journal go to: <http://journals.asm.org/subscriptions/>

---

# Cross Talk between the Cell Wall Integrity and Cyclic AMP/Protein Kinase A Pathways in *Cryptococcus neoformans*

Maureen J. Donlin,<sup>a,b</sup> Rajendra Upadhy,<sup>c</sup> Kimberly J. Gerik,<sup>c</sup> Woei Lam,<sup>c</sup> Laura G. VanArendonk,<sup>d</sup> Charles A. Specht,<sup>e</sup> Neil K. Sharma,<sup>c</sup> Jennifer K. Lodge<sup>c</sup>

Edward A. Doisy Department of Biochemistry and Molecular Biology<sup>a</sup> and Department of Molecular Microbiology and Immunology,<sup>b</sup> Saint Louis University School of Medicine, St. Louis, Missouri, USA; Department of Molecular Microbiology<sup>c</sup> and Center for Genome Sciences and Systems Biology,<sup>d</sup> Washington University School of Medicine, St. Louis, Missouri, USA; Department of Medicine, University of Massachusetts Medical School, Worcester, Massachusetts, USA<sup>e</sup>

**ABSTRACT** *Cryptococcus neoformans* is a fungal pathogen of immunocompromised people that causes fatal meningitis. The fungal cell wall is essential to viability and pathogenesis of *C. neoformans*, and biosynthesis and repair of the wall is primarily controlled by the cell wall integrity (CWI) signaling pathway. Previous work has shown that deletion of genes encoding the four major kinases in the CWI signaling pathway, namely, *PKC1*, *BCK1*, *MKK2*, and *MPK1* results in severe cell wall phenotypes, sensitivity to a variety of cell wall stressors, and for *Mpk1*, reduced virulence in a mouse model. Here, we examined the global transcriptional responses to gene deletions of *BCK1*, *MKK2*, and *MPK1* compared to wild-type cells. We found that over 1,000 genes were differentially expressed in one or more of the deletion strains, with 115 genes differentially expressed in all three strains, many of which have been identified as genes regulated by the cyclic AMP (cAMP)/protein kinase A (PKA) pathway. Biochemical measurements of cAMP levels in the kinase deletion strains revealed significantly less cAMP in all of the deletion strains compared to the wild-type strain. The deletion strains also produced significantly smaller capsules than the wild-type KN99 strain did under capsule-inducing conditions, although the levels of capsule they shed were similar to those shed by the wild type. Finally, addition of exogenous cAMP led to reduced sensitivity to cell wall stress and restored surface capsule to levels near those of wild type. Thus, we have direct evidence of cross talk between the CWI and cAMP/PKA pathways that may have important implications for regulation of cell wall and capsule homeostasis.

**IMPORTANCE** *Cryptococcus neoformans* is a fungal pathogen of immunocompromised people that causes fatal meningitis. The fungal cell wall is essential to viability and pathogenesis of *C. neoformans*, and biosynthesis and repair of the wall are primarily controlled by the cell wall integrity (CWI) signaling pathway. In this study, we demonstrate that deletion of any of three core kinases in the CWI pathway impacts not only the cell wall but also the amount of surface capsule. Deletion of any of the kinases results in significantly reduced cellular cyclic AMP (cAMP) levels, and addition of exogenous cAMP rescues the capsule defect and some cell wall defects, supporting a direct role for the CWI pathway in regulation of capsule in conjunction with the cAMP/protein kinase A pathway.

Received 30 June 2014 Accepted 3 July 2014 Published 12 August 2014

**Citation** Donlin MJ, Upadhy R, Gerik KJ, Lam W, Gibbs VanArendonk L, Specht CA, Sharma NK, Lodge JK. 2014. Cross talk between the cell wall integrity and cyclic AMP/protein kinase A pathways in *Cryptococcus neoformans*. mBio 5(4):e01573-14. doi:10.1128/mBio.01573-14.

**Editor** Joseph Heitman, Duke University

**Copyright** © 2014 Donlin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Noncommercial-ShareAlike 3.0 Unported license](https://creativecommons.org/licenses/by-nc-sa/4.0/), which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Address correspondence to Jennifer K. Lodge, lodgejk@wustl.edu.

This article is a direct contribution from a Fellow of the American Academy of Microbiology.

*Cryptococcus neoformans* is a fungal pathogen of immunocompromised people that causes an estimated 1 million infections each year among HIV-positive patients globally, resulting in 600,000 deaths annually (1). Much of the disease burden occurs in sub-Saharan Africa, where deaths from cryptococcal infections may exceed those from tuberculosis in some areas (1). Cryptococcal infections can be successfully treated with antifungal agents, but the mortality rate remains 15 to 30%, even in the context of antiviral treatments for HIV (2–4). The closely related species *Cryptococcus gattii* can infect immunocompetent people and animals. Although *C. gattii* is more common in Australia, it received global attention in 1999 when it emerged on Vancouver Island in Canada and has since spread into other areas of the Pacific Northwest (5–7).

Infection by *C. neoformans* occurs upon inhalation of spores or desiccated yeast cells, which can then disseminate to the brain and cause fatal meningitis (reviewed in references 8, 9, and 10). *C. neoformans* rapidly establishes an infection in lung alveoli, where low iron availability and increased carbon dioxide levels induce the formation of a polysaccharide capsule on the surface of the cryptococcal cell. The *C. neoformans* capsule has been shown to be protective against phagocytosis and oxidative stress as well as reducing the T-cell immune response (11, 12). The capsule is a key virulence factor, and strains with disruptions or deletions of capsule-related genes are usually avirulent or demonstrate attenuated virulence in murine models of infection (reviewed in reference 13).

*C. neoformans* is a soilborne pathogen found ubiquitously in

**TABLE 1** Number of differentially expressed genes

Strain(s)	No. of genes			No. of unique genes		
	Total	Up <sup>a</sup>	Down <sup>a</sup>	Total	Up	Down
<i>bck1Δ</i> strain	627	340	287	244	171	73
<i>mkk2Δ</i> strain	511	121	390	203	27	176
<i>mpk1Δ</i> strain	453	278	175	206	127	79
<i>bck1Δ</i> and <i>mkk2Δ</i> strains	164	39	123			
<i>bck1Δ</i> and <i>mpk1Δ</i> strains	103	85	13			
<i>mkk2Δ</i> and <i>mpk1Δ</i> strains	28	12	11			
<i>bck1Δ</i> , <i>mkk2Δ</i> , and <i>mpk1Δ</i> strains	116	44	71			

<sup>a</sup> For genes found in more than one strain, they were listed as upregulated (Up) or downregulated (Down) only if the direction of the expression difference was concordance.


the environment of temperate zones that can also infect and persist in mammalian hosts. Thus, *C. neoformans* must be able to grow under a wide range of pHs and temperatures and be able to cope with a variety of exogenous stresses secreted by other soil organisms or mammalian host cells. The ability to survive in such diverse environments requires that the cell be able to sense and process multiple signals from the environment that result in activation of various signaling cascades to allow adaption to and growth under different conditions (reviewed in reference 14).

The cryptococcal cell wall is vital for maintaining cell morphology and forms the scaffold for capsule attachment. Cell wall homeostasis is maintained by the cell wall integrity (CWI) pathway, which has 4 core kinases, protein kinase C (Pkc1), bypass of protein kinase C (Bck1), mitogen-activated kinase kinase (Mkk2), and mitogen-activated protein kinase (Mpk1), all of which are necessary for a fully functional cell wall. The primary kinase for this pathway is Pkc1, a diacylglycerol-activated protein kinase C essential for responding to a variety of cell wall stresses and whose deletion results in severe cell wall defects, reduced melanin, and disrupted capsule (15, 16). Strains with a deletion of the *BCK1* or *MKK2* gene show increased sensitivity to cell wall stressors and high temperature as well as delayed melanin formation (17). Deletion of the gene encoding the putative terminal kinase in the cell wall integrity pathway, *MPK1*, results in attenuated virulence (18) and has been shown to reduce virulence when deleted in *C. gattii* (19). Previous studies of the CWI pathway have used phosphorylation of the terminal kinase, Mpk1, as a marker for activation of the pathway (18). We have demonstrated that in response to nitrosative or oxidative stress, *PKC1* is required for phosphorylation of the terminal kinase, Mpk1 (16). However, in a *pkc1Δ* strain, Mpk1 is still phosphorylated in response to heat shock, but not when the *BCK1* or *MKK2* gene is deleted (20), suggesting that inputs into this signaling cascade can bypass some of the core kinases and still result in activation of the pathway. We hypothesize that this pathway is likely to interact with other signaling cascades, such as the HOG1 and cyclic AMP (cAMP)/protein kinase A (PKA) pathways that respond to various exogenous stresses. To address this hypothesis, we used RNA sequencing to examine the genome-wide expression differences between the wild type and deletions in three of the core CWI kinases, *BCK1*, *MKK2*, and *MPK1*. We find strong evidence of cross talk between the cAMP/PKA and CWI pathways based on the genes differentially expressed in the kinase deletion strains. We demonstrate that deletion of any of the three kinases results in significantly decreased cAMP levels and reduced surface capsule and that exogenous cAMP can partially rescue the capsule- and cell wall-related defects of the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains.

## RESULTS

**Impact of CWI kinases on the transcriptome.** The expression profiles of the three kinase deletion strains (*bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains) were compared to wild-type *C. neoformans* KN99 MATa (KN99a) when grown at 30°C in yeast extract-peptone-dextrose (YPD) medium plus 1 M sorbitol to an optical density (OD) of 1.0. We initially included a deletion of the *PKC1* gene in the experiment, which necessitated the use of sorbitol to act as a stabilizer (16). The RNA expression data revealed that the *pkc1Δ* strain was aneuploid, and thus, it was excluded from further analyses. Overall, 1,064 genes showed statistically significant differential expression with a corrected *P* value cutoff of <0.05 (Table 1; see Table S2 in the supplemental material) (21). More genes were differentially expressed in the *bck1Δ* strain (627 genes) than in the *mkk2Δ* (511 genes) and *mpk1Δ* (453 genes) strains. Of the 1,064 differentially expressed genes, 653 genes were found in only one of the three strains with a fairly even distribution: 244 genes were unique to the *bck1Δ* strain, 203 were unique to the *mkk2Δ* strain, and 206 were unique to the *mpk1Δ* strain (Fig. 1).

There were 116 transcripts differentially expressed in all three of the deletion strains, including genes involved in carbohydrate metabolism, cell wall biogenesis, transport, and signaling (see Table S3 in the supplemental material). Of the 116 genes, 115 were in concordance among the three mutant strains, with 71 upregulated genes and 44 downregulated genes compared to the wild type. The one gene that was not in concordance was a gene encoding a hypothetical protein of unknown function. We identified 10 of the



**FIG 1** Comparison of the distribution and overlap of the differentially expressed genes in the CWI kinase deletion strains, the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains.

TABLE 2 Cell wall- or capsule-related genes

Gene type and Broad ID <sup>a</sup>	Annotation	Log <sub>2</sub> fold change <sup>b</sup>			Reference(s)
		<i>bck1Δ</i> /WT strains	<i>mkk2Δ</i> /WT strains	<i>mpk1Δ</i> /WT strains	
Cell wall-related genes					
CNAG_00261	MPN10 putative mannoprotein			0.43	
CNAG_00546	CHS4 chitin synthase activity	−1.22	−1.33	−1.01	25
CNAG_00914	KRE6 β-glucan synthase	0.42			23
CNAG_01230	CDA2 chitin deacetylase			−0.61	22
CNAG_01234	SPO71 similar to <i>S. cerevisiae</i> spore wall assembly protein	−1.07	−1.81		
CNAG_01341	MPI1 putative mannose-6-phosphate isomerase		−1.17		
CNAG_02071	BNI4 scaffold protein	0.47			50
CNAG_02217	CHS7 chitin synthase	−0.87		−1.15	25
CNAG_02225	EXG1 cellulase	1.11		0.61	51
CNAG_03099	CHS1 chitin synthase	1.03	1.15	0.59	25
CNAG_03223	MPN6 putative mannoprotein	−1.30	−1.25	−1.35	
CNAG_03345	SSD1 translational repressor	−0.63			17
CNAG_03465	LAC1 laccase melanin synthesis	−0.75	−1.18	−0.66	52
CNAG_03782	MPN11 putative mannose protein		−3.34		
CNAG_05663	SCW1 similar to <i>S. cerevisiae</i> cell wall integrity protein	0.37	0.55		
CNAG_05818	CHS5 chitin synthase	−1.46	−1.31	−1.34	25
CNAG_06291	FPD1 polysaccharide deacetylase	−0.68	−1.19	−0.66	22, 51
CNAG_06487	CHS6 chitin synthase	0.51	0.50		25
CNAG_06835	KRE61 β-glucan synthase	−1.33	−2.79	−1.03	23
CNAG_07499	CHS8 chitin synthase			−0.56	25
Capsule-related genes					
CNAG_00600	CAP60 capsule-associated protein	0.81			53
CNAG_00721	CAP59 capsule-associated protein	0.75	0.67	0.31	54
CNAG_01172	PBX1 parallel β-helix repeat protein			0.35	55
CNAG_01371	CRG2 regulator of G-protein signaling	0.46			56, 57
CNAG_01654	CAS34 capsule structure protein	0.61			58
CNAG_02581	CAS33 capsule-associated protein	0.39			59
CNAG_03735	CAP4 glycosyltransferase	0.67	0.56		58
CNAG_05081	PDE1 phosphodiesterase		−0.89		60
CNAG_07554	CAP10 glycosyltransferase	0.59			61

<sup>a</sup> Broad ID, Broad identification or locus tag.<sup>b</sup> The log<sub>2</sub> fold changes in the level of expression for the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* deletion strains compared to that of the wild-type (WT) strain are shown.


115 genes as related to cell wall or capsule homeostasis, reflecting the importance of these kinases working in concert to maintain cell wall integrity (CWI).

**Differential expression of cell wall- and capsule-related genes.** Among all of the differentially expressed transcripts, we identified 29 genes that are known or predicted to be involved in cell wall biogenesis, cell wall regulation, or capsule synthesis (Table 2). Six of the eight chitin synthase genes, *CHS1*, *CHS4*, *CHS5*, *CHS6*, *CHS7*, and *CHS8*, show altered expression in two or more of the kinase deletion strains (Table 2). The *CHS1*, *CHS4*, and *CHS5* genes show altered expression in all three deletion strains, with *CHS1* upregulated and *CHS4* and *CHS5* showing decreased expression. A deacetylase gene, *FPD1*, for which no function has yet been identified (22) was downregulated in all three deletion strains, suggesting a possible adaptive response to the cell wall stress induced by the deletions. The *KRE6* gene, which is involved in β-1,6-glucan synthesis (23), was modestly upregulated in the *bck1Δ* strain, but *KRE61*, a homolog of *KRE6* with no known function, was highly repressed in all three deletions, again suggesting a possible adaptive response or regulation of this gene by the CWI pathway. Of the nine capsule-related genes that are differentially expressed, seven are changed in the *bck1Δ* strain, but only one (*CAP59*) is found in all three deletion strains, precluding a

significant role for the CWI pathway in transcriptional regulation of capsule biosynthesis. Although *CAP59* has been associated mostly with capsule defects, it may have a role in general secretion, rather than a capsule-specific role (24).

**Reduced chitin levels in the CWI kinase deletion strains.** All of the deletion strains had significant cell wall phenotypes, which likely affect cellular chitin and chitosan levels. We determined the chitin and chitosan content of the cell walls, and all three had significantly more chitin and significantly less chitosan than did the wild-type KN99a strain (Fig. 2). The chitin synthase gene, *CHS3*, responsible for the majority of chitosan synthesis during vegetative growth was unchanged in these deletion strains, as were two of the three chitin deacetylase genes. Three chitin synthase genes, *CHS1*, *CHS4*, and *CHS5*, had altered expression in all three of these deletion strains (Table 2). Previous work on the *CHS* genes revealed no phenotypes when *CHS1* was deleted, although *CHS1* transcript could be detected in vegetatively growing cells (25). Strains with deletion of either *CHS4* or *CHS5* have reduced chitin and increased chitosan content, but otherwise had no other *in vitro* cell wall phenotypes (25). These data suggest a possible role for *CHS1*, *CHS4*, and/or *CHS5* in maintaining cell wall morphology but one that is dependent on the CWI kinases. Previous work has demonstrated an important role for chitin oligomers in





**FIG 2** *C. neoformans* BCK1, MKK2, and MPK1 are required for full chitin production. Chitin and chitosan levels of *C. neoformans* KN99a, KN99α, and *bck1Δ*, *mkk2Δ*, and *mpk1Δ* deletion strains. Measurements of strains were taken after 70 h of growth in liquid YPD at 25°C and reported as average nanomoles per milligram (dry weight) of chitin and chitosan as indicated. Each measurement represents the average from three independent cultures for each strain, and error bars indicate standard deviations. The levels of chitin and chitosan for the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains were compared to either KN99a or KN99α using a one-way ANOVA and Dunnett's posthoc test with the KN99a strain as the control. The chitin and chitosan values for all three deletion strains were highly significantly different ( $P < 0.001$ ) from the value for the KN99a strain.

capsule attachment (26, 27). However, the source of the chitin for the oligomers has not been determined. It is plausible that *CHS1*, *CHS4*, and/or *CHS5* may play a role in producing the chitin necessary for capsule attachment.


**Overlap of differentially expressed genes with other stress-responsive signaling pathways.** Because of the cell wall and stress phenotypes associated with the CWI kinase deletion strains, we compared our differentially expressed gene lists with those generated from deletions in genes that regulate the cAMP/PKA pathway (28), the HOG1 stress pathway (29), and capsule synthesis (30–32). There is a substantial overlap in genes identified as differentially expressed in all of these studies, which is expected given that these gene lists include several hundred to thousands of genes. We narrowed the search to examine the overlap of genes differentially expressed in all three of our deletion strains with these other data sets (see Table S3 in the supplemental material). We found the greatest overlap (54 and 56 genes, respectively) with genes dependent on *HOG1*, a mitogen-activated protein (MAP)-type kinase involved in osmoregulation (33) and *SSK1*, a response receiver protein that modulates Hog1 activity (34). We found substantial overlap (38 genes) with *RIM101*-dependent genes (31), with genes dependent on *NRG1* (31 genes) (32) and with genes dependent on *SKN7* (27 genes), another response receiver protein that regulates

sensitivity to Na<sup>+</sup> ions (34). Out of the 115 genes differentially expressed in all three CWI kinase deletion strains were 10 genes that have a known or predicted role in cell wall or capsule biosynthesis or regulation. Of the 10 genes, 6 genes overlap with the genes of a *skn7Δ* strain, 5 overlap with those of a *hog1Δ* strain (29), and 5 overlap with those of a *nrg1Δ* strain (32) (Table S3). Although these high-throughput experiments were conducted under different conditions, the genes differentially expressed in various deletion strains suggest a central role for these genes in response to perturbation of the different signaling pathways and potential mechanisms of cross talk between the pathways.

**Reduced intracellular cAMP in CWI kinase deletion strains.** Recent work has demonstrated the importance of the *RIM101* transcription factor in the cAMP/PKA-dependent regulation of capsule production, defining a novel interaction between the Rim and PKA signaling pathways in *C. neoformans* (30). PKA1 activates the transcription factor NRG1, leading to the transcriptional activation of genes directly involved in capsule assembly (35). The Hog1 kinase is known to be a negative regulator of capsule synthesis in serotype A (33), a process that requires cAMP. In addition, the gene encoding Smg1, identified as a suppressor of Gpa1, the G alpha subunit that functions in cAMP signaling (36), was significantly repressed in all three CWI kinase deletion strains. The overlap between genes differentially expressed in the kinase deletion strains and those differentially expressed in gene deletion strains related to the PKA/cAMP pathway led us to ask if loss of the CWI kinases might negatively impact cellular cAMP levels. We measured cAMP levels in the three deletion strains and wild-type KN99a during log-phase growth in YPD medium. These experiments were done on two separate days, with three replicates for each strain. Since we observed the differential gene expression under log growth in nutrient-rich conditions and not in response to stress, we wanted to measure the cAMP levels under similar conditions. The kinase deletion strains, *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains, all showed significantly less cAMP compared to the KN99a wild-type strain (Fig. 3A). The experiments do not address the effect of cAMP in response to stress; however, the significant decrease in cAMP levels among all three deletion strains provides strong evidence for an impact on basal cAMP activity.

We tested whether the addition of exogenous cAMP could mitigate the sensitivity of the deletions to SDS, calcofluor white, and Congo red. Exogenous cAMP almost completely rescues the sensitivity to SDS for *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains after 5 days (Fig. 3B), but it has a lower effect on sensitivity to calcofluor white and does not rescue sensitivity to Congo red (data not shown). It is possible that the different stresses activate the CWI and cAMP/PKA pathways by different mechanisms, with various degrees of overlap between the activating mechanisms. An alternative explanation may be that cell membranes are more permeable in the presence of SDS. cAMP does not readily cross cell membranes, and the fact that we saw an effect with SDS but not with Congo red may reflect increased permeability in the presence of SDS, which does not happen with Congo red treatment.

**Capsule defect in the CWI kinase deletion strains.** To further investigate the cross talk between the cAMP/PKA and CWI pathways, we examined phenotypes associated with disrupted cAMP levels, specifically, capsule formation when grown in capsule-inducing low-iron medium (LIM) (37). When the deletion strains were grown in LIM, all three had visibly reduced capsule com-



**FIG 3** The downstream kinase deletion strains produce less cAMP than KN99a cells. (A) The cAMP levels were measured in the different deletion strains. Values that are statistically significantly different from the value for KN99a strain as determined by a one-way ANOVA with Dunnett's posthoc test are indicated by an asterisk. Error bars represent the standard deviations. (B) Phenotypes of *bck1Δ*, *mkk2Δ*, and *mpk1Δ* deletion strains grown on YPD, YPD plus 0.01% SDS, or YPD plus 0.01% SDS plus 10 mM cAMP at 3 and 5 days.

pared to wild-type KN99a (Fig. 4A and C), and measurement of the capsule diameters showed significantly reduced capsule diameters for *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains compared to the wild-type KN99a strain (Fig. 4B and C and Table 3). The addition of 10 mM cAMP to the medium increased the size of the capsule in all three deletion strains to wild-type levels while having essentially no effect on the KN99a strain (Fig. 4B and C and Table 3). These data point to an important, and previously unknown, role for the CWI kinases in regulation of cAMP levels. It suggests that production of cAMP is dependent on a functioning CWI pathway, as disruption of the pathway leads to decreased cAMP levels and causes impairment of capsule induction in LIM.

#### Analysis of shed capsule in the CWI kinase deletion strains.

The defect in the increase of capsular polysaccharide in the mutant strains under capsule-inducing conditions suggests that the disruption of the CWI pathway affects the biosynthesis of the capsule, secretion of the capsule material, or attachment of the secreted capsule to the fungal cell wall. To differentiate between these possible explanations, we collected the shed capsular material from the wild-type and mutant strains. Secreted capsular material corresponding to an equal number of yeast cells was resolved on an agarose gel, transferred to a nylon membrane, and probed to determine the amount and relative size of the capsular material with an antiglucuronoxylomannan (anti-GXM) antibody (monoclonal antibody 3C2). We found that the amounts of capsule shed by the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains were nearly equal the amount shed by wild-type cells; however, they exhibited slight differences in the migration pattern on an agarose gel, suggesting potential differences in the physical-chemical properties of the shed capsule material (Fig. 5). Overall, these data suggest that the observed capsule defects in these strains are more likely due to a defect in attachment, not in synthesis or secretion.

#### DISCUSSION


Eukaryotic cellular signaling cascades receive inputs from several proteins simultaneously and then balance the signals to generate

an appropriate response. One of the kinases in the CWI pathway, *PKC1*, plays a central role in the response to various cell wall-perturbing agents and production of virulence factors, capsule, and melanin (15, 16). However, the CWI pathway can be induced by some stresses, such as heat shock, in a *PKC1*-independent manner (20). To explore possible alternative pathway connections or cross talk, we determined the global transcription response to the loss of three other CWI kinases, *BCK1*, *MKK2*, and *MPK1*. We found that over 1,000 genes were differentially expressed in one or more of the deletion strains (see Table S2 in the supplemental material), including 29 cell wall- or capsule-related genes (Table 2) and 115 genes with concordant differential expression in all three strains (Table S3). These data are consistent with a significant adaptive response of the cell to the loss of the CWI kinases.

We hypothesized that disruption of three core kinases in the cell wall integrity pathway affect signaling in other pathways that impact cell wall and capsule homeostasis and looked for overlap between the 115 genes showing differential expression in all three kinase deletion strains (see Table S3 in the supplemental material) with genes dependent on other stress response-related genes as well as those that regulate capsule and cell wall homeostasis. We found a substantial overlap with genes dependent on stress-responsive *HOG1* (29) and *SSK1* (34). We also found >30 genes that overlapped with genes dependent on *RIM101* (31) and *NRG1* (32), which suggested a connection to the cAMP/PKA pathway. We found a significant reduction in the cellular cAMP levels in all three of the CWI kinase deletion strains compared to wild-type cells (Fig. 3A). The addition of exogenous cAMP reduced the sensitivity to some, but not all, cell wall-perturbing agents (Fig. 3B), suggesting multiple inputs for maintaining cell wall homeostasis that are dependent on both the cAMP/PKA and CWI pathways.

We observed a loss of surface capsule with deletion of the CWI kinases and restoration of the surface capsule in those strains with addition of exogenous cAMP (Fig. 4). This is the first experimental evidence of a direct connection between the CWI pathway and





**FIG 4** *C. neoformans* *BCK1*, *MKK2*, and *MPK1* are required for capsule production. (A and B) Cells were incubated under capsule-inducing conditions for 4 days without cAMP (A) or with 10 mM cAMP (B). Capsule size was assessed by staining with India ink and visualizing the zone of exclusion at a magnification of  $\times 60$ . The capsule diameters of the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains were reduced compared to that of the KN99a strain, and capsule size was restored when grown with exogenous cAMP. (C) Comparison of capsule diameters between strain KN99a and *bck1Δ*, *mkk2Δ* and *mpk1Δ* strains grown in low-iron medium (LIM) with and without the addition of 10 mM cAMP. Measurements were taken for a minimum of 100 cells for each strain. Asterisks represent statistically significant differences as determined by a one-way ANOVA with Dunnett's posthoc test. Error bars represent the 95% confidence intervals.

capsule regulation in *C. neoformans*. The reduced capsule does not appear to be due to a defect in synthesis or secretion but is more likely due to a defect in attachment to the cell wall, as we did not observe a significant difference in the amount of shed capsule between the wild-type and kinase deletion strains (Fig. 5). The nature of this defect is not clear, but it is known that  $\alpha$ -glucan and chitin oligomers or chitin-like structures are required for capsule attachment (26, 27, 38). The CWI kinase deletion strains have

reduced chitin/chitosan ratio compared to wild-type cells (Fig. 2), and a similarly reduced chitin/chitosan ratio was also observed in *chs4Δ* and *chs5Δ* strains (25). Expression of both *CHS4* and *CHS5* was downregulated in all three kinase deletion strains (Table 2); however, neither the *chs4Δ* nor *chs5Δ* strain had a measurable capsule defect when grown in LIM (data not shown). A third chitin synthase gene, *CHS1*, was upregulated in all three deletion strains (Table 2), but deletion of *CHS1* has no discernible cell

TABLE 3 Average induced capsule and cell diameters

Strain or genotype	Capsule diam		Capsule diam		Cell diam		<i>P</i> value <sup>d</sup>
	LIM <sup>a</sup>	<i>P</i> value <sup>b</sup>	LIM + cAMP <sup>a</sup>	<i>P</i> value <sup>c</sup>	LIM <sup>a</sup>	LIM + cAMP <sup>a</sup>	
KN99a	2.05 ± 0.61		2.83 ± 0.79	1.00	6.01 ± 1.10	6.16 ± 1.77	0.45
<i>bck1Δ</i>	1.19 ± 0.31	<0.001	2.28 ± 0.39	1.00	4.92 ± 0.61	5.58 ± 0.87	<0.001
<i>mkk2Δ</i>	1.24 ± 0.24	<0.001	1.83 ± 0.51	0.78	4.56 ± 0.72	5.22 ± 0.74	<0.001
<i>mpk1Δ</i>	1.18 ± 0.31	<0.001	1.48 ± 0.53	1.00	4.36 ± 0.69	5.39 ± 1.2	<0.001

<sup>a</sup> Capsule and cell diameters of cells grown in LIM alone or in LIM plus cAMP are reported as averages ± standard deviations of at least 100 measurements.

<sup>b</sup> *P* value comparing the capsule diameters of deletion strains to that of the WT strain (KN99a) in low-iron medium (LIM) only.

<sup>c</sup> *P* value comparing the capsule diameters between deletion strains and WT in LIM plus cAMP.

<sup>d</sup> *P* value comparing the cell diameters for each strain with and without cAMP in the medium.

wall-related phenotype (25) or capsule-related phenotype in LIM (data not shown).

The *RIM101* transcription factor has been shown to play a significant role in the regulation of cell wall morphogenesis (31). Despite the overlap of genes dependent on all three CWI kinases with those dependent on *RIM101*, the *rim101Δ* strain has significantly different phenotypic characteristics compared to the CWI kinase deletion strains. The *rim101Δ* strain is sensitive to LiCl and NaCl but insensitive to SDS, calcofluor white, or Congo red. More interestingly, the *rim101Δ* strain has an increased chitin/chitosan ratio, and two chitin synthase genes, *CHS4* and *CHS5*, are upregulated in the *rim101Δ* strain but downregulated in the CWI kinase deletion strains. This raises the possibility that *RIM101* and the CWI kinases may play antagonistic roles in the regulation of cell wall homeostasis.

Only three genes with a likely role in transcription regulation were differentially expressed in all three CWI deletion strains; the three genes were CNAG\_01103, CNAG\_05290, and CNAG\_07775 (see Table S3 in the supplemental material). Of the three genes, only one, CNAG\_05290 which encodes SPT3, a sub-

unit of the SAGA transcriptional regulatory complex seems likely to have a role in cell wall or capsule regulation (32, 39). The CNAG\_05290 transcript for the putative transcription factor, SPT3, is upregulated in an *ada2Δ* strain at 37°C with CO<sub>2</sub> compared to the wild type (32), but no further characterization of this putative transcription factor has been published.

The regulation of capsule production in *C. neoformans* is very complex. Capsule synthesis is responsive to iron and CO<sub>2</sub> levels via the cAMP/PKA signaling pathway (reviewed in reference 13), and the downstream changes in capsule biosynthetic and trafficking gene expression are controlled by at least five different transcription factors, which are also responsive to a number of other proteins (30, 32, 35, 40–42). Therefore, the hypothesis that the CWI pathway might have a role in the regulation of capsule production is plausible. One possible mechanism of capsule regulation is that the CWI kinases and some subset of chitin synthases interact to regulate the synthesis of the chitin oligomers that are necessary for capsule attachment (26, 27). However, experimentally discerning the mechanism of that regulation is beyond the scope of this study.

In previous studies, deletions of *C. neoformans* genes with homology to regulators of the *Saccharomyces cerevisiae* CWI pathway, *ROM2*, *LRG1*, and *SIT4* (reviewed in reference 43) resulted in moderate to severe cell wall-related phenotypes and increased sensitivity to cell wall stressors (17). However, we saw no differential expression of these genes in any of the CWI kinase deletion strains, nor did we observe differential expression of the regulators of the HOG pathway, *SKN7*, *SSK1*, and *TCO1* or *TCO2* (33, 34) in the kinase deletion strains. However, we did observe considerable overlap between genes differentially expressed in the CWI kinase deletion strains with genes that are differentially expressed in response to the deletion of the stress-responsive genes *HOG1*, *SSK1*, and *SKN7* (29, 33, 34). The lack of differences in expression of genes previously identified as critical for stress response may reflect the fact that regulation of their activity occurs primarily via protein modifications, not by transcriptional changes. We also conducted our expression analysis of the CWI kinase deletion strains under nutrient-rich conditions in the absence of exogenous stress. It is possible that if the CWI kinase deletion strains were exposed to an external stress, we would observe transcriptional changes in some of these other pathway regulators.

In summary, we have demonstrated a direct connection between the CWI and cAMP/PKA pathways in the regulation of intracellular cAMP levels. These data support a role for the CWI pathway in the regulation of surface capsule. The mechanisms and transcription factors, which mediate the cAMP and capsule response to the CWI pathway, remain to be elucidated.




FIG 5 The levels of capsule shed by the *bck1Δ*, *mkk2Δ*, and *mpk1Δ* strains were similar to the level shed by the wild-type KN99a strain. The wild-type, *bck1Δ*, *mkk2Δ*, *mpk1Δ*, and *cap59Δ* strains were grown in YPD medium for 5 days, and the number of cells was determined by absorption at 650 nm. The supernatants were collected, and the cell counts were normalized before loading. The gel was transferred to a blot and probed with the 3C2 anti-GXM monoclonal antibody.

## MATERIALS AND METHODS

**Strains and media.** Wild-type (WT) *C. neoformans* var. *grubii* strain KN99 (serotype A, MAT $\alpha$ ) was used in this study (see Table S1 in the supplemental material). Strains containing deletions in *BCK1* or *MKK2* in the KN99 MAT $\alpha$  (KN99a) background were generated by overlap PCR (44) using the primers listed in Table S1. A strain deleted for *MPK1* in the KN99 MAT $\alpha$  background was generated by amplification of the *mpk1* $\Delta$  cassette (pNAT-STM150) from strain KK3, containing *mpk1* $\Delta$  in the H99 MAT $\alpha$  background (generously provided by Joseph Heitman, Duke University) (45). Each fragment was biolistically transformed into strain KN99a (46), and isolates were selected and verified by PCR screening as described previously (17) utilizing the primers listed in Table S1. Southern blotting was performed, using the selectable marker as a probe to rule out ectopic integrations.

**RNA preparation and sequencing.** The KN99a, *bck1* $\Delta$ , *mkk2* $\Delta$ , and *mpk1* $\Delta$  cells were grown for 3 days on plates containing yeast extract-peptone-dextrose (YPD) medium supplemented with 1 M D-sorbitol. A 50-ml flask of YPD medium supplemented with 1 M sorbitol was inoculated and grown overnight at 30°C with shaking (300 rpm). Cells were diluted to an optical density at 650 nm (OD<sub>650</sub>) of 0.004 and grown overnight to an OD<sub>650</sub> of 0.75 to 1.0. A strain with a deletion in the *PKC1* gene, which required sorbitol to grow, was originally part of this study, but it was later determined to have an aneuploid chromosome. It was excluded from further analyses (data not shown). RNA was prepared from three biological replicates using the Qiagen RNeasy plant minikit (Qiagen, Valencia, CA) with minor modifications (details of methods available upon request). RNA quality was assessed on a formaldehyde agarose gel, and the ratio of 25S/18S rRNA was quantified using ImageQuant software (GE Healthcare Biosciences, Pittsburgh, PA). Total RNA (10  $\mu$ g) from each sample was used as starting material to prepare sequencing libraries with the Illumina mRNA sequencing sample preparation kit (Illumina, San Diego, CA) following the manufacturer's instructions. Samples were bar-coded and sequenced on a single lane of an Illumina HiSeq sequencing system to generate nondirectional, single-ended 50-bp reads. The library preparation and RNA sequencing were performed by the Genome Access Technology Center at Washington University (<http://gtac.wustl.edu>). The bar code sequences were trimmed to 42 bp, and quality-filtered reads were mapped to the H99 reference genome sequence prepared by the Broad Institute ([http://www.broadinstitute.org/annotation/genome/cryptococcus\\_neoformans/MultiHome.html](http://www.broadinstitute.org/annotation/genome/cryptococcus_neoformans/MultiHome.html)) using Bow tie 2.1.0 as implemented in the TopHat-Cufflinks suite (21, 47).

**Differential gene expression.** The *C. neoformans* H99 transcript file prepared by the Broad Institute was used as the reference. Transcript files for each biological replicate were generated from each alignment file using Cufflinks, merged into a single transcript file using Cuffmerge, and differentially expressed transcripts were identified using Cuffdiff (21). Genes were considered differentially expressed if *P* values were <0.05 after Benjamini-Hochberg correction for multiple testing.

**Capsule measurements.** Strains were grown in low-iron medium [LIM plus ethylenediamine-*N,N'*-bis(2-hydroxyphenylacetic acid) (ED-DHA)] (37) with and without exogenous 10 mM cAMP for 4 days at 30°C. All mutant strains were compared to the wild-type KN99a strain. Capsule-induced strains were suspended in a 1:4 India ink-H<sub>2</sub>O solution and photographed on an Olympus BX61 microscope at a magnification of  $\times 60$ . Cell diameters (one including the capsule and one excluding it) were measured for a minimum of 100 cells per strain using SlideBook, version 5.0. Capsule length was calculated as the difference between the radii obtained from the diameters. All values were averaged, and comparisons were considered significant if the *P* value was <0.05 using a one-way analysis of variance (ANOVA) with a Dunnet's posthoc test (one-tailed comparison; < control).

**Chitin and chitosan assays.** The cellular chitin and chitosan levels were determined as previously described (22). Samples were divided into two aliquots. One aliquot was treated with acetic anhydride to measure chitin plus chitosan. The other aliquot was left untreated to measure total

chitin. Chitosan was estimated from the difference between the two measurements.

**Measurement of cAMP.** Overnight cultures of the WT and the deletion strains were diluted in 50 ml YPD medium to an OD<sub>650</sub> of 0.05 and grown for 20 h at 30°C with shaking (300 rpm). Cells were harvested, washed, and resuspended in MES-EDTA buffer (10 mM morpholineethanesulfonic acid [MES], 0.1 mM EDTA, pH 6.0). Cells were counted using a hemocytometer, centrifuged, and resuspended at a concentration of  $5 \times 10^8$  cells/ml. Cells (0.5 ml) were lysed by bead beating in 2.5% trichloroacetic acid at 4°C with 0.5-mm glass beads for 1 min, followed by a 2-min rest, for a total of 4 cycles. Crude lysates were centrifuged at  $20,800 \times g$  for 15 min. Lysates were extracted four times with water-saturated diethyl ether and centrifuged for  $20,800 \times g$  at 4°C for 2 min between extractions, frozen in liquid nitrogen, and lyophilized overnight. The lyophilized samples were suspended in 0.5 ml of 0.1 M HCl, and cAMP assays were performed according to Sigma cAMP enzyme immunoassay kit (catalog no. CA200; Sigma, St. Louis, MO). Assays were performed twice with three technical replicates. cAMP concentrations were calculated for individual samples, and statistical significance was determined using a one-way ANOVA with Dunnet's posthoc test (two-tailed test) using IBM SPSS Statistics (version 20).

**SDS viability assays.** Overnight cultures of wild-type strain KN99a and *bck1* $\Delta$ , *mkk2* $\Delta$ , and *mpk1* $\Delta$  strains in YPD medium were diluted to an OD<sub>650</sub> of 1.0 with phosphate-buffered saline (PBS). Tenfold serial dilutions were plated on YPD medium plus 0.01% SDS and YPD medium plus 0.01% SDS and 10 mM cAMP. The plates were incubated at 30°C for up to 5 days and photographed. The SDS concentration used was based on levels determined to be deleterious, but not lethal, to the mutant strains.

**Immunoblotting of the secreted capsule.** Immunoblotting of the shed capsular material was performed as described previously (48). Briefly, conditioned medium for each strain corresponding to an equal number of yeast cells grown in YPD medium was passed through a 0.22- $\mu$ m filter, mixed with DNA loading dye, and separated on a 0.6% agarose gel using Tris-acetic acid-EDTA buffer (pH 8.0). The gel was transferred onto a positively charged nylon membrane by capillary transfer and immunoblotted with 1  $\mu$ g/ml of antiglycuroxylomannan (anti-GXM) 3C2 antibody (generously provided by Thomas Kozel, University of Nevada) (49).

**Microarray data accession number.** RNA sequences/microarray data were deposited in the NCBI GEO database and assigned accession number GSE57217.

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.01573-14/-/DCSupplemental>.

Table S1, DOCX file, 0.1 MB.

Table S2, PDF file, 0.3 MB.

Table S3, PDF file, 0.1 MB.

## ACKNOWLEDGMENTS

We thank Ping Wang for technical assistance with cAMP assays, David Gohara for technical assistance with analysis of the RNA sequence data, and Nicole Gilbert for helpful discussions and manuscript review. We thank the Genome Technology Access Center in the Department of Genetics at Washington University School of Medicine for help with genomic analysis.

This work was supported by the National Institutes of Health grant R01 AI50184 to J.K.L. The Genome Technology Access Center is partially supported by NCI Cancer Center support grant P30 CA91842 to the Siteman Cancer Center and by ICTS/CTSA grant UL1TR000448 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH) and NIH Roadmap for Medical Research.

This publication is solely the responsibility of the authors and does not necessarily represent the official view of NCRR or NIH.



## REFERENCES

- Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM. 2009. Estimation of the current global burden of cryptococcal meningitis among persons living with HIV/AIDS. *AIDS* 23:525–530. <http://dx.doi.org/10.1097/QAD.0b013e328322ffac>.
- Lortholary O. 2007. Management of cryptococcal meningitis in AIDS: the need for specific studies in developing countries. *Clin. Infect. Dis.* 45: 81–83. <http://dx.doi.org/10.1086/518583>.
- Mdodo R, Brown K, Omonge E, Jaoko W, Baddley J, Pappas P, Kempf MC, Aban I, Odera S, Suleh A, Jolly PE. 2010. The prevalence, clinical features, risk factors and outcome associated with cryptococcal meningitis in HIV positive patients in Kenya. *East Afr. Med. J.* 87:481–487.
- Bratton EW, El Hussein N, Chastain CA, Lee MS, Poole C, Stürmer T, Juliano JJ, Weber DJ, Perfect JR. 2012. Comparison and temporal trends of three groups with cryptococcosis: HIV-infected, solid organ transplant, and HIV-negative/non-transplant. *PLoS One* 7:e43582. <http://dx.doi.org/10.1371/journal.pone.0043582>.
- Kidd SE, Hagen F, Tschärke RL, Huynh M, Bartlett KH, Fyfe M, Macdougall L, Boekhout T, Kwon-Chung KJ, Meyer W. 2004. A rare genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on Vancouver Island (British Columbia, Canada). *Proc. Natl. Acad. Sci. U. S. A.* 101:17258–17263. <http://dx.doi.org/10.1073/pnas.0402981101>.
- Datta K, Bartlett KH, Baer R, Byrnes E, Galanis E, Heitman J, Hoang L, Leslie MJ, MacDougall L, Magill SS, Morshed MG, Marr KA, *Cryptococcus gattii* Working Group of the Pacific Northwest. 2009. Spread of *Cryptococcus gattii* into Pacific Northwest region of the United States. *Emerg. Infect. Dis.* 15:1185–1191. <http://dx.doi.org/10.3201/eid1508.081384>.
- Byrnes EJ, III, Li W, Lewit Y, Ma H, Voelz K, Ren P, Carter DA, Chaturvedi V, Bildfell RJ, May RC, Heitman J. 2010. Emergence and pathogenicity of highly virulent *Cryptococcus gattii* genotypes in the northwest United States. *PLoS Pathog.* 6:e1000850. <http://dx.doi.org/10.1371/journal.ppat.1000850>.
- Kronstad JW, Attarian R, Cadieux B, Choi J, D'Souza CA, Griffiths EJ, Geddes JM, Hu G, Jung WH, Kretschmer M, Saikia S, Wang J. 2011. Expanding fungal pathogenesis: *Cryptococcus* breaks out of the opportunistic box. *Nat. Rev. Microbiol.* 9:193–203. <http://dx.doi.org/10.1038/nrmicro2522>.
- Kozubowski L, Heitman J. 2012. Profiling a killer, the development of *Cryptococcus neoformans*. *FEMS Microbiol. Rev.* 36:78–94. <http://dx.doi.org/10.1111/j.1574-6976.2011.00286.x>.
- Sabiiti W, May RC. 2012. Mechanisms of infection by the human fungal pathogen *Cryptococcus neoformans*. *Future Microbiol.* 7:1297–1313. <http://dx.doi.org/10.2217/fmb.12.102>.
- Zaragoza O, Rodrigues ML, De Jesus M, Frases S, Dadachova E, Casadevall A. 2009. The capsule of the fungal pathogen *Cryptococcus neoformans*. *Adv. Appl. Microbiol.* 68:133–216. [http://dx.doi.org/10.1016/S0065-2164\(09\)01204-0](http://dx.doi.org/10.1016/S0065-2164(09)01204-0).
- Syme RM, Bruno TF, Kozel TR, Mody CH. 1999. The capsule of *Cryptococcus neoformans* reduces T-lymphocyte proliferation by reducing phagocytosis, which can be restored with anticapsular antibody. *Infect. Immun.* 67:4620–4627.
- O'Meara TR, Alspaugh JA. 2012. The *Cryptococcus neoformans* capsule: a sword and a shield. *Clin. Microbiol. Rev.* 25:387–408. <http://dx.doi.org/10.1128/CMR.00001-12>.
- Kozubowski L, Lee SC, Heitman J. 2009. Signalling pathways in the pathogenesis of *Cryptococcus*. *Cell. Microbiol.* 11:370–380. <http://dx.doi.org/10.1111/j.1462-5822.2008.01273.x>.
- Heung LJ, Kaiser AE, Luberto C, Del Poeta M. 2005. The role and mechanism of diacylglycerol-protein kinase C1 signaling in melanogenesis by *Cryptococcus neoformans*. *J. Biol. Chem.* 280:28547–28555. <http://dx.doi.org/10.1074/jbc.M503404200>.
- Gerik KJ, Bhimireddy SR, Ryerse JS, Specht CA, Lodge JK. 2008. PKC1 is essential for protection against both oxidative and nitrosative stresses, cell integrity, and normal manifestation of virulence factors in the pathogenic fungus *Cryptococcus neoformans*. *Eukaryot. Cell* 7:1685–1698. <http://dx.doi.org/10.1128/EC.00146-08>.
- Gerik KJ, Donlin MJ, Soto CE, Banks AM, Banks IR, Maligie MA, Selitrennikoff CP, Lodge JK. 2005. Cell wall integrity is dependent on the PKC1 signal transduction pathway in *Cryptococcus neoformans*. *Mol. Microbiol.* 58:393–408. <http://dx.doi.org/10.1111/j.1365-2958.2005.04843.x>.
- Kraus PR, Fox DS, Cox GM, Heitman J. 2003. The *Cryptococcus neoformans* MAP kinase Mpk1 regulates cell integrity in response to antifungal drugs and loss of calcineurin function. *Mol. Microbiol.* 48:1377–1387. <http://dx.doi.org/10.1046/j.1365-2958.2003.03508.x>.
- Ngamskulrungron P, Price J, Sorrell T, Perfect JR, Meyer W. 2011. *Cryptococcus gattii* virulence composite: candidate genes revealed by microarray analysis of high and less virulent Vancouver Island outbreak strains. *PLoS One* 6:e16076. <http://dx.doi.org/10.1371/journal.pone.0016076>.
- Lam WC, Gerik KJ, Lodge JK. 2013. Role of *Cryptococcus neoformans* Rho1 GTPases in the PKC1 signaling pathway in response to thermal stress. *Eukaryot. Cell* 12:118–131. <http://dx.doi.org/10.1128/EC.05305-11>.
- Trapnell C, Hendrickson DG, Sauvageau M, Goff L, Rinn JL, Pachter L. 2013. Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat. Biotechnol.* 31:46–53.
- Baker LG, Specht CA, Donlin MJ, Lodge JK. 2007. Chitosan, the deacetylated form of chitin, is necessary for cell wall integrity in *Cryptococcus neoformans*. *Eukaryot. Cell* 6:855–867. <http://dx.doi.org/10.1128/EC.00399-06>.
- Gilbert NM, Donlin MJ, Gerik KJ, Specht CA, Djordjevic JT, Wilson CF, Sorrell TC, Lodge JK. 2010. KRE genes are required for beta-1,6-glucan synthesis, maintenance of capsule architecture and cell wall protein anchoring in *Cryptococcus neoformans*. *Mol. Microbiol.* 76:517–534. <http://dx.doi.org/10.1111/j.1365-2958.2010.07119.x>.
- Garcia-Rivera J, Chang YC, Kwon-Chung KJ, Casadevall A. 2004. *Cryptococcus neoformans* CAP59 (or Cap59p) is involved in the extracellular trafficking of capsular glucuronoxylomannan. *Eukaryot. Cell* 3:385–392. <http://dx.doi.org/10.1128/EC.3.2.385-392.2004>.
- Banks IR, Specht CA, Donlin MJ, Gerik KJ, Levitz SM, Lodge JK. 2005. A chitin synthase and its regulator protein are critical for chitosan production and growth of the fungal pathogen *Cryptococcus neoformans*. *Eukaryot. Cell* 4:1902–1912. <http://dx.doi.org/10.1128/EC.4.11.1902-1912.2005>.
- Fonseca FL, Nimrichter L, Cordero RJ, Frases S, Rodrigues J, Goldman DL, Andruszkiewicz R, Milewski S, Travassos LR, Casadevall A, Rodrigues ML. 2009. Role for chitin and chitooligomers in the capsular architecture of *Cryptococcus neoformans*. *Eukaryot. Cell* 8:1543–1553. <http://dx.doi.org/10.1128/EC.00142-09>.
- Rodrigues ML, Alvarez M, Fonseca FL, Casadevall A. 2008. Binding of the wheat germ lectin to *Cryptococcus neoformans* suggests an association of chitinlike structures with yeast budding and capsular glucuronoxylomannan. *Eukaryot. Cell* 7:602–609. <http://dx.doi.org/10.1128/EC.00307-07>.
- Maeng S, Ko YJ, Kim GB, Jung KW, Floyd A, Heitman J, Bahn YS. 2010. Comparative transcriptome analysis reveals novel roles of the Ras and cyclic AMP signaling pathways in environmental stress response and antifungal drug sensitivity in *Cryptococcus neoformans*. *Eukaryot. Cell* 9:360–378. <http://dx.doi.org/10.1128/EC.00309-09>.
- Ko YJ, Yu YM, Kim GB, Lee GW, Maeng PJ, Kim S, Floyd A, Heitman J, Bahn YS. 2009. Remodeling of global transcription patterns of *Cryptococcus neoformans* genes mediated by the stress-activated HOG signaling pathways. *Eukaryot. Cell* 8:1197–1217. <http://dx.doi.org/10.1128/EC.00120-09>.
- O'Meara TR, Norton D, Price MS, Hay C, Clements MF, Nichols CB, Alspaugh JA. 2010. Interaction of *Cryptococcus neoformans* Rim101 and protein kinase A regulates capsule. *PLoS Pathog.* 6:e1000776. <http://dx.doi.org/10.1371/journal.ppat.1000776>.
- O'Meara TR, Holmer SM, Selvig K, Dietrich F, Alspaugh JA. 2013. *Cryptococcus neoformans* Rim101 is associated with cell wall remodeling and evasion of the host immune responses. *mBio* 4(1):e00522-12. <http://dx.doi.org/10.1128/mBio.00522-12>.
- Haynes BC, Skowrya ML, Spencer SJ, Gish SR, Williams M, Held EP, Brent MR, Doering TL. 2011. Toward an integrated model of capsule regulation in *Cryptococcus neoformans*. *PLoS Pathog.* 7:e1002411. doi: 10.1371/journal.ppat.1002411.
- Bahn YS, Kojima K, Cox GM, Heitman J. 2005. Specialization of the HOG pathway and its impact on differentiation and virulence of *Cryptococcus neoformans*. *Mol. Biol. Cell* 16:2285–2300. <http://dx.doi.org/10.1091/mbc.E04-11-0987>.
- Bahn YS, Kojima K, Cox GM, Heitman J. 2006. A unique fungal two-component system regulates stress responses, drug sensitivity, sexual de-

- velopment, and virulence of *Cryptococcus neoformans*. *Mol. Biol. Cell* 17: 3122–3135. <http://dx.doi.org/10.1091/mbc.E06-02-0113>.
35. Cramer KL, Gerrald QD, Nichols CB, Price MS, Alspaugh JA. 2006. Transcription factor Nrg1 mediates capsule formation, stress response, and pathogenesis in *Cryptococcus neoformans*. *Eukaryot. Cell* 5:1147–1156. <http://dx.doi.org/10.1128/EC.00145-06>.
  36. Palmer DA, Thompson JK, Li L, Prat A, Wang P. 2006. Gb2, a novel Gbeta-like/RACK1 homolog, functions as a Gbeta subunit in cAMP signaling and is essential in *Cryptococcus neoformans*. *J. Biol. Chem.* 281: 32596–32605. <http://dx.doi.org/10.1074/jbc.M602768200>.
  37. Vartivarian SE, Anaissie EJ, Cowart RE, Sprigg HA, Tingler MJ, Jacobson ES. 1993. Regulation of cryptococcal capsular polysaccharide by iron. *J. Infect. Dis.* 167:186–190. <http://dx.doi.org/10.1093/infdis/167.1.186>.
  38. Reese AJ, Doering TL. 2003. Cell wall alpha-1,3-glucan is required to anchor the *Cryptococcus neoformans* capsule. *Mol. Microbiol.* 50: 1401–1409. <http://dx.doi.org/10.1046/j.1365-2958.2003.03780.x>.
  39. O'Meara TR, Hay C, Price MS, Giles S, Alspaugh JA. 2010. *Cryptococcus neoformans* histone acetyltransferase Gcn5 regulates fungal adaptation to the host. *Eukaryot. Cell* 9:1193–1202. <http://dx.doi.org/10.1128/EC.00098-10>.
  40. Jung WH, Sham A, White R, Kronstad JW. 2006. Iron regulation of the major virulence factors in the AIDS-associated pathogen *Cryptococcus neoformans*. *PLoS Biol.* 4:e410. <http://dx.doi.org/10.1371/journal.pbio.0040410>.
  41. Jung WH, Saikia S, Hu G, Wang J, Fung CK, D'Souza C, White R, Kronstad JW. 2010. HapX positively and negatively regulates the transcriptional response to iron deprivation in *Cryptococcus neoformans*. *PLoS Pathog.* 6:e1001209. <http://dx.doi.org/10.1371/journal.ppat.1001209>.
  42. Chun CD, Brown JC, Madhani HD. 2011. A major role for capsule-independent phagocytosis-inhibitory mechanisms in mammalian infection by *Cryptococcus neoformans*. *Cell Host Microbe* 9:243–251. <http://dx.doi.org/10.1016/j.chom.2011.02.003>.
  43. Levin DE. 2011. Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the cell wall integrity signaling pathway. *Genetics* 189: 1145–1175. <http://dx.doi.org/10.1534/genetics.111.128264>.
  44. Davidson RC, Blankenship JR, Kraus PR, de Jesus Berrios M, Hull CM, D'Souza C, Wang P, Heitman J. 2002. A PCR-based strategy to generate integrative targeting alleles with large regions of homology. *Microbiology* 148:2607–2615.
  45. Kojima K, Bahn YS, Heitman J. 2006. Calcineurin, Mpk1 and Hog1 MAPK pathways independently control fludioxonil antifungal sensitivity in *Cryptococcus neoformans*. *Microbiology* 152:591–604. <http://dx.doi.org/10.1099/mic.0.28571-0>.
  46. Toffaletti DL, Rude TH, Johnston SA, Durack DT, Perfect JR. 1993. Gene transfer in *Cryptococcus neoformans* by use of biolistic delivery of DNA. *J. Bacteriol.* 175:1405–1411.
  47. Jendretzki A, Wittland J, Wilk S, Straede A, Heinisch JJ, Fuchs BB, Mylonakis E. 2011. How do I begin? Sensing extracellular stress to maintain yeast cell wall integrity. *Eur. J. Cell Biol.* 90:740–744. <http://dx.doi.org/10.1016/j.ejcb.2011.04.006>.
  48. Yoneda A, Doering TL. 2008. Regulation of *Cryptococcus neoformans* capsule size is mediated at the polymer level. *Eukaryot. Cell* 7:546–549. <http://dx.doi.org/10.1128/EC.00437-07>.
  49. Gates-Hollingsworth MA, Kozel TR. 2009. Phenotypic heterogeneity in expression of epitopes in the *Cryptococcus neoformans* capsule. *Mol. Microbiol.* 74:126–138. <http://dx.doi.org/10.1111/j.1365-2958.2009.06855.x>.
  50. Liu OW, Chun CD, Chow ED, Chen C, Madhani HD, Noble SM. 2008. Systematic genetic analysis of virulence in the human fungal pathogen *Cryptococcus neoformans*. *Cell* 135:174–188. <http://dx.doi.org/10.1016/j.cell.2008.07.046>.
  51. Biondo C, Mancuso G, Midiri A, Bombaci M, Messina L, Beninati C, Teti G. 2006. Identification of major proteins secreted by *Cryptococcus neoformans*. *FEMS Yeast Res.* 6:645–651. <http://dx.doi.org/10.1111/j.1567-1364.2006.00043.x>.
  52. Williamson PR. 1994. Biochemical and molecular characterization of the diphenol oxidase of *Cryptococcus neoformans*: identification as a laccase. *J. Bacteriol.* 176:656–664.
  53. Chang YC, Kwon-Chung KJ. 1998. Isolation of the third capsule-associated gene, CAP60, required for virulence in *Cryptococcus neoformans*. *Infect. Immun.* 66:2230–2236.
  54. Chang YC, Kwon-Chung KJ. 1994. Complementation of a capsule-deficient mutation of *Cryptococcus neoformans* restores its virulence. *Mol. Cell. Biol.* 14:4912–4919.
  55. Liu OW, Kelly MJ, Chow ED, Madhani HD. 2007. Parallel beta-helix proteins required for accurate capsule polysaccharide synthesis and virulence in the yeast *Cryptococcus neoformans*. *Eukaryot. Cell* 6:630–640. <http://dx.doi.org/10.1128/EC.00398-06>.
  56. Shen G, Wang YL, Whittington A, Li L, Wang P. 2008. The RGS protein Crg2 regulates pheromone and cyclic AMP signaling in *Cryptococcus neoformans*. *Eukaryot. Cell* 7:1540–1548. <http://dx.doi.org/10.1128/EC.00154-08>.
  57. Xue C, Hsueh YP, Chen L, Heitman J. 2008. The RGS protein Crg2 regulates both pheromone and cAMP signalling in *Cryptococcus neoformans*. *Mol. Microbiol.* 70:379–395. <http://dx.doi.org/10.1111/j.1365-2958.2008.06417.x>.
  58. Moyrand F, Fontaine T, Janbon G. 2007. Systematic capsule gene disruption reveals the central role of galactose metabolism in *Cryptococcus neoformans* virulence. *Mol. Microbiol.* 64:771–781. <http://dx.doi.org/10.1111/j.1365-2958.2007.05695.x>.
  59. Moyrand F, Chang YC, Himmelreich U, Kwon-Chung KJ, Janbon G. 2004. CaS3p belongs to a seven-member family of capsule structure designer proteins. *Eukaryot. Cell* 3:1513–1524.
  60. Hicks JK, Bahn YS, Heitman J. 2005. Pde1 phosphodiesterase modulates cyclic AMP levels through a protein kinase A-mediated negative feedback loop in *Cryptococcus neoformans*. *Eukaryot. Cell* 4:1971–1981. <http://dx.doi.org/10.1128/EC.4.12.1971-1981.2005>.
  61. Chang YC, Kwon-Chung KJ. 1999. Isolation, characterization, and localization of a capsule-associated gene, CAP10, of *Cryptococcus neoformans*. *J. Bacteriol.* 181:5636–5643.