

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2014

Pharmacokinetic changes and dosing modification of aminoglycosides in critically ill obese patients: A literature review

Dimitrios Velissaris
University Hospital of Patras

Vasilios Karamouzos
University Hospital of Patras

Markos Marangos
University Hospital of Patras

Charalampos Pierrakos
Brugmann University Hospital

Menelaos Karanikolas
Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Velissaris, Dimitrios; Karamouzos, Vasilios; Marangos, Markos; Pierrakos, Charalampos; and Karanikolas, Menelaos, "Pharmacokinetic changes and dosing modification of aminoglycosides in critically ill obese patients: A literature review." *Journal of Clinical Medicine Research*. 6, 4. 227-233. (2014).
https://digitalcommons.wustl.edu/open_access_pubs/3260

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Pharmacokinetic Changes and Dosing Modification of Aminoglycosides in Critically Ill Obese Patients: A Literature Review

Dimitrios Velissaris^a, Vasilios Karamouzos^a, Markos Marangos^a,
Charalampos Pierrakos^b, Menelaos Karanikolas^{c, d}

Abstract

The objective of the paper is to review the literature and provide recommendations for use of aminoglycoside antibiotics in critically ill obese patients. Literature search in PubMed for all articles on the use of aminoglycosides in critically ill obese patients was conducted, and all articles related to pharmacokinetics in obesity were reviewed. Bibliographies of all searched manuscripts were also reviewed in an attempt to find additional references. Although aminoglycoside pharmacokinetics have been described in detail, data on aminoglycoside use and appropriate dose modification in critically ill obese patients are very limited. Knowledge on aminoglycoside pharmacokinetics and use in critically ill obese patients is incomplete. Pathophysiologic changes in obesity can result in sub- or supra-therapeutic aminoglycoside plasma concentrations, especially in the presence of sepsis. Rigorous clinical studies are needed to establish aminoglycoside dosing guidelines in critically ill obese patients with sepsis.

Keyword: Aminoglycoside; Gentamicin; Amikacin; Tobramycin; Obesity; Sepsis; Critical illness; Pharmacokinetics

Introduction

Obesity is a major worldwide health problem, and is associated with serious diseases and increased morbidity and mortality. Physiologic changes in obesity significantly influence

antibiotic pharmacokinetic characteristics, including distribution, protein binding, metabolism and renal excretion. In order to reach therapeutic levels, some medications require dosing adjustments in obese patients, especially in the presence of critical illness. Aminoglycosides act synergistically with other antimicrobial agents, and are very important for treatment of serious infections. However, because aminoglycoside plasma levels can be significantly affected by obesity, due to aminoglycoside distribution into adipose tissue, dosing modification is needed in order to reach therapeutic plasma levels, and therapeutic drug monitoring (TDM) has been recommended. The aim of this study is to review data on aminoglycoside pharmacokinetics in obesity and summarize published data regarding the use of aminoglycosides and the need for dose modification in critically ill obese patients with sepsis.

Literature Search Methods

We searched for publications relevant to this review, using the PubMed database. The initial PubMed search was conducted in September 2012 and was updated in October 2013. The search was conducted using the terms “aminoglycosides dosing and obesity”, “aminoglycosides pharmacokinetics and obesity”, “aminoglycosides in sepsis and obesity” and “aminoglycosides pharmacokinetics in sepsis” as keywords. We excluded animal studies, and studies published in languages other than English. There was no exclusion based on type of aminoglycoside or publication date. The search included all types of articles, including case reports and review articles, and the bibliography of all extracted manuscripts was reviewed in attempt to identify additional references. All articles published in English related to aminoglycoside use in obesity were reviewed. When we identified publications of overlapping data, we only used data from the newer or more detailed publication. Two authors reviewed the abstract and text of all articles that seemed relevant to this review.

Literature Review

Our literature search revealed 20 articles in the “aminogly-

Manuscript accepted for publication May 6, 2014

^aInternal Medicine Department, University Hospital of Patras, Rion 26500, Greece

^bIntensive Care Department, Brugmann University Hospital, Brussels 1030, Belgium

^cDepartment of Anesthesiology, Washington University School of Medicine, St. Louis, Missouri, USA

^dCorresponding Author: Menelaos Karanikolas, Department of Anesthesiology, Washington University School of Medicine, Campus Box 8054, 660 S. Euclid Ave, St Louis, Missouri, 63110, USA.
Email: menelaos.karanikolas@gmail.com; karanikm@wustl.edu

doi: <http://dx.doi.org/10.14740/jocmr1858w>

Table 1. Summary, in Chronological Order, of Studies on Aminoglycoside Pharmacokinetics and Dose Adjustment in Obesity

Author, year	Antibiotic studied	Main findings
Schwartz et al, 1978 [1]	Gentamicin, tobramycin	Vd in obese subjects closely approximates Vd in normal subjects when using the formula: $ABW = IBW + 0.4 \times (TBW - IBW)$
Blouin et al, 1979 [2]	Tobramycin	Tobramycin loading dose based on $ABW = IBW + 0.58 \times (TBW - IBW)$
Korsager, 1980 [4]	Gentamicin	Gentamicin uptake in adipose tissue = 43.7% of uptake in total body mass of normal-weight patients
Bauer et al, 1980 [3]	Amikacin	Amikacin loading dose based on $ABW = IBW + 0.38 \times (TBW - IBW)$
Bauer et al, 1983 [5]	Gentamicin, tobramycin, amikacin	Loading dose based on $ABW = IBW + 0.4 \times (TBW - IBW)$
Leader et al, 1994 [11]	Gentamicin	Initial dose based on calculation of $ClCr$ by Cockcroft equation with $IBW + 0.4 \times (TBW - IBW)$
Traynor et al, 1995 [8]	Tobramycin	The TBW/IBW ratio predicts Vd. For patients with TBW/IBW ratio ≥ 1.25 , doses based on $ABW = IBW + 0.43 \times (TBW - IBW)$
Wurtz et al, 1997 [10]	Aminoglycosides	$DWCF = 0.4$, therefore $ABW = IBW + 0.4 \times (TBW - IBW)$
Pai et al, 2007 [12]	Aminoglycosides	Dosing of aminoglycosides should be based on ABW.

ABW: adjusted body weight; $ClCr$: creatinine clearance; $DWCF$: dosing weight correction factor; IBW: ideal body weight; TBW: total body weight; Vd: volume of distribution.

cosides dosing and obesity” term, 42 articles in the “aminoglycosides pharmacokinetics and obesity”, 165 articles in the “aminoglycosides pharmacokinetics in sepsis” and seven articles in the “aminoglycosides in sepsis and obesity”. We then searched for publications with text combining most keywords. The combination of the above searches revealed a total of 33 articles, case series and letters which were included in this review.

Overall, we found very limited data on aminoglycoside pharmacokinetics and dose modification in obesity. Four early studies on morbidly obese patients, by Schwartz referring to gentamicin and tobramycin in 1978 [1], by Blouin referring to tobramycin in 1979 [2], by Bauer referring to amikacin in 1980 [3] and by Korsager referring to gentamicin in 1980 [4] concluded that morbidly obese patients have larger volume of distribution (Vd) compared to normal weight patients.

Schwartz et al evaluated the pharmacokinetics of gentamicin and tobramycin and calculated elimination constants and volumes of distribution in 13 obese vs. 13 normal weight subjects. Their results showed that gentamicin and tobramycin are distributed less to adipose tissue compared to other

tissues, and mean Vd in obese subjects closely approximates the Vd seen in normal subjects, when using normalized body mass +40% of adipose mass as total weight in obese subjects [1]. Bauer et al studied aminoglycoside pharmacokinetics in two matched groups of 30 normal weight vs. 30 obese patients with documented Gram negative infection, and concluded that pharmacokinetics are altered in obesity, therefore morbidly obese patients require significantly higher gentamicin, tobramycin or amikacin doses compared to normal weight patients [5]. To compensate for increased Vd in obesity, aminoglycoside loading doses can be based on adjusted body weight (ABW), which is calculated using the formula: $ABW = IBW + 0.4 \times (TBW - IBW)$, where IBW is ideal body weight, and TBW is total body weight [5]. However, these studies were conducted at a time when once-daily antibiotic regimens (5 - 7 mg/kg for tobramycin or gentamicin), which are popular today, were not widely used. Analysis of pharmacokinetic data from 2073 patients (497 receiving tobramycin and 1,576 receiving gentamicin) using various formulas for Vd, showed that Vd was normalized for both tobramycin and gentamicin at all weight categories with use of lean body weight (LBW) estimated based as described by

Janmahasatian et al [6, 7].

In 1995, Traynor et al investigated aminoglycoside pharmacokinetics in 1,708 patients who received gentamicin and tobramycin. Regression analysis revealed that the TBW to IBW ratio (TBW/IBW) predicted Vd. Furthermore, there were no large differences between dosing weight correcting factors (DWCFs) derived from IBW and body mass index (BMI)-based classification systems. This study concluded that, when calculating aminoglycoside doses, BMI values were not better than IBW, and proposed that DWCF for aminoglycoside doses should be $IBW + 0.43 \times EBW$ (where EBW is the excess body weight ($EBW = TBW - IBW$)) for patients with TBW/IBW ratio ≥ 1.25 [8].

In 1997, Wurtz et al published a review on antimicrobial pharmacokinetics and dosing in obesity, and proposed that, because 30% of adipose tissue is water, a DWCF of 0.3 should be used. Therefore the weight used to calculate doses of hydrophilic antibiotics is derived from the equation: $IBW + 0.3 \times (TBW - IBW)$, where IBW is derived from the Devine formula [9] and TBW is total body weight. However, other clinical data suggest that the DWCF is 0.4 for aminoglycosides [10]. Published data on aminoglycoside DWCFs are summarized in Table 1 [1-5, 8, 10-12].

Our search revealed several reports describing physiologic changes in obesity [2, 10, 13, 14]. Because aminoglycoside dosing in critically ill obese patients has not been extensively studied, we could only find limited data, including a literature review by Erstad, which showed that circulatory changes in sepsis affect dosing and pharmacokinetics [15]. Data published by Cheymol support modification of loading and maintenance doses in critically ill obese patients due to physiologic alterations related to sepsis [16]. Although dose modification is described in studies by Leader, Corcoran and Blouin, appropriate aminoglycoside doses in critically ill obese patients are still debated [2, 11, 13, 17].

Regarding existing methods for measuring creatinine clearance (CICr), published data provide accurate estimates for patients with normal weight, but not for obese ones. Compared to measured CICr in obese patients, when the Cockcroft-Gault equation is used with IBW, CICr is underestimated, whereas CICr is overestimated when TBW is used [18]. The Salazar-Corcoran equation, which shows strong correlation between CICr and fat free body mass is as accurate as the Cockcroft-Gault equation for the normal weight patients, but seems superior when applied to obese patients [19]. Therefore, when CICr is not measured, estimation of CICr using the Salazar-Corcoran equation should improve the ability to select appropriate doses for drugs cleared principally by renal filtration.

With regard to administration of aminoglycoside maintenance doses, most meta-analyses showed that once-daily regimens are equal or superior to multiple dose regimens, with regard to clinical efficacy, bacteriologic efficacy and nephrotoxicity [20-24].

Discussion

Pathophysiologic changes in obesity affect most hydrophilic medications [10, 14, 25]. Because pharmacokinetic alterations in obesity affect absorption, Vd and clearance of many antibiotics, estimation of the correct antibiotic dose is of critical importance, especially in septic patients with multiple organ failure, where morbidity and mortality are high. However, despite obesity becoming a worldwide problem, antibiotic dosing in obese patients has not been studied adequately, and published data are very limited [26].

Morbid obesity alters aminoglycosides pharmacokinetics, as both Vd and clearance are increased in morbidly obese patients compared to patients with normal weight [16], but most published data in obesity refer to drug distribution. Pathophysiologic changes in obesity result in altered cardiac structure and function with increased cardiac output, altered tissue blood flow and increased gut perfusion. However, drug absorption does not seem to be altered, and several studies evaluating drug absorption did not find differences in obese vs. non-obese patients [10, 15]. At the present time, although Vd of many drugs, particularly lipophilic ones, is altered in obesity, there is no universally accepted factor to estimate Vd in obese individuals [27]. Although plasma proteins such as α_1 acid-glycoprotein and lipoproteins are highly concentrated in obesity, binding of drugs to albumin does not seem to be altered. Vd of drugs equally soluble in water and oil (such as aminoglycosides) is slightly increased in obesity. Also, because distribution of a drug between adipose tissue and other tissues influences pharmacokinetics in obese patients, loading dose should be adjusted for IBW. Maintenance dosage also needs adjustment, depending on changes in drug clearance, as described in several studies on obese individuals [16]. Hepatic clearance of many drugs is also altered in obesity, but the metabolic capacity of the liver must be reduced $> 90\%$ before drug metabolism is significantly affected [25, 28].

In conclusion, Vd seems to be the pharmacokinetic variable most affected in obesity, and can be estimated using an ABW that includes a fraction of the excess body weight (TBW-IBW) [29].

CICr, a useful measure for approximating glomerular filtration rate (GFR), is generally overestimated when it is calculated based on TBW, but is generally underestimated when based on IBW [2, 10]. The Cockcroft-Gault equation can be used to estimate GFR in lean patients, but its use in obesity is questionable due to disparity between muscle mass and body weight ratio [18]. The Salazar-Corcoran equation is an attempt to more accurately estimate CICr in obesity, by taking into account serum creatinine, gender, TBW, age and height [19]. A comparison of the Cockcroft-Gault vs. the Salazar-Corcoran equation showed that prediction of gentamicin elimination rate constant, clearance and elimination half time was best when CICr was estimated using the Cock-

roft-Gault equation [11]. However, according to retrospective studies, the Salazar-Corcoran formula is more precise for obese patients [12, 30].

Aminoglycosides (gentamicin, amikacin, tobramycin, netilmicin, kanamycin and streptomycin) are cornerstones of therapy for serious Gram negative and aerobic bacilli infections [31], therefore appropriate aminoglycoside loading and maintenance dosing is important for good outcome. When initiating aminoglycoside therapy, a loading dose is needed to achieve therapeutic levels quickly. The loading dose is very important with aminoglycosides, because a high initial dose is needed to maximize their anti-bacterial effect and lower mortality [32, 33]. In addition, because aminoglycoside effect is concentration-dependent, administration of maintenance doses using extended interval regimens is preferred. Loading dose depends on Vd and target plasma concentration (Cp), both of which are affected by critical illness: Vd is altered in sepsis, due to alterations in microvascular permeability and abnormalities of extracellular body water. In one prospective study, Vd was estimated to be 0.43 ± 0.12 L/kg in the beginning of gentamicin therapy, but, as patient condition stabilized Vd was reduced to 0.29 ± 0.17 L/kg on the seventh day of treatment [34]. Because Vd and drug metabolism change with patient condition, optimization of aminoglycoside dosing is difficult in critical illness: hepatic and renal failure can affect antibiotic clearance, and critically ill patients have increased Vd and variable clearance compared to patients with less severe disease [35]. As Vd increases, aminoglycoside loading doses need to be increased in sepsis [15], and plasma level monitoring (also known as TDM) has been recommended to improve safety and effectiveness [34, 36, 37].

Aminoglycoside dosing in obese critically ill patients is even more challenging because drug distribution between fat and lean tissue influences pharmacokinetics, therefore aminoglycosides, being hydrophilic agents, need dosing adjustment. At the present time, there are limited data on the effects of fluid changes in critically ill obese ICU patients with sepsis [15], but the need for dose adjustment has been established for gentamicin, tobramycin and amikacin [2, 11, 17].

In healthy adults, aminoglycoside Vd is approximately 0.26 L/kg (range: 0.2 - 0.3). However, aminoglycoside dosing needs modification in obesity using a correcting factor of 0.4, because an estimated 40% of the dose is distributed into adipose tissue. [11, 15, 17]. Therefore, the equation for calculating ABW reads as follows: $ABW = IBW + 0.4 \times (TBW - IBW)$.

The above dosing modification is very important, because calculation of aminoglycoside dosing based on TBW may result in high serum concentrations, thereby increasing the risk for nephrotoxicity and ototoxicity [2, 16] in obesity. However, data regarding nephrotoxicity are conflicting, with studies on aminoglycoside kinetics showing no significant

difference between obese patients and patients with normal weight with regard to ClCr for gentamicin [11].

Obesity is related to glomerular hyperfiltration, rather than to effects on tubular secretion [7]. The Cockcroft-Gault [18], the Salazar-Corcoran [19] and the modification of diet in renal disease (MDRD) [38] equation are the equations most commonly used for estimation of clearance. However, although these formulas are widely used, they are not reliable estimators of function, particularly in patients with high BUN/SCr ratio [39]. Furthermore, the predictive performance of newer formulas such as the MDRD and chronic kidney disease-epidemiology (CKD-EPI) [40] compared to the Cockcroft-Gault equation for assessment of kidney function and estimation of aminoglycoside clearance in obesity is unknown [41]. A study by Pai suggested that the CKD-EPI equation best predicts aminoglycoside clearance [7]. Overall, MDRD provides more reliable estimations of renal function compared to the Cockcroft-Gault formula, but both formulas lack precision. MDRD is not superior to Cockcroft-Gault for drug dosing, and MDRD estimated GFR (eGFR) needs adjustment to patient's body surface area for drug dosing. Although Cockcroft-Gault has the advantage of simplicity and longer use [42], drug dosing guidelines are often based on ClCr estimated using the Cockcroft-Gault equation as surrogate for GFR.

Controversy exists with regard to the use of these formulas in obesity: the Cockcroft-Gault equation relies on TBW, and therefore overestimates GFR in obese patients. Similarly, the MDRD equation, which indexes GFR based on a normalized body surface area of mL/min/1.73 m² also overestimates GFR in obese patients.

Aminoglycoside dosing is based on weight, according to either of two strategies: traditional, more frequent dosing, vs. extended-interval dosing, but dosing interval should be adjusted in patients with renal impairment. According to the American Society of Health Systems Pharmacists, in patients with normal renal function, traditional dosing consists of 1 - 2 mg/kg doses given multiple times per day, whereas the newer strategy utilizes 5 - 10 mg/kg every 24 h. Although aminoglycosides have been administered using multiple daily dosing regimens for decades, newer clinical and laboratory studies suggest that once-daily dosing confers advantages with regard to efficacy [43, 44]. With regard to extended interval dosing in obesity, a retrospective analysis by Ross suggests that use of DWCF of 40% is accurate [45].

In ICU patients, however, other factors such as sepsis and renal dysfunction can significantly alter aminoglycoside pharmacokinetics, and individual TDM is warranted [46]. Details regarding the effects of fluid shifts on aminoglycoside pharmacokinetics have not been well studied in critically ill obese patients. Finally, the advantages of large once-daily aminoglycoside doses have been questioned, because of apparent pyrogen-mediated toxicity [47]. As relevant published data are very limited [47], there is great need for well-

conducted clinical trials on obese critically ill patients [15].

Understanding of the pathophysiologic changes in obesity and knowledge of the literature on aminoglycosides are valuable for rational determination of dosages in obese patients. As aminoglycosides are first-line drugs in sepsis, there is clearly a need for more studies on the appropriate use and dosing of aminoglycosides in critically ill obese patients. It is important to realize that studies on pharmacokinetics in obesity have limitations, such as patient population heterogeneity, small patient numbers, the fact that many data are referring to single doses rather than ongoing use, and limited knowledge on the effects of obesity on pharmacokinetics. Failure to appropriately modify doses in obesity can result in therapeutic failure or increased toxicity, therefore TDM, when feasible, has been recommended in order to optimize the safety and effectiveness of aminoglycoside antibiotic therapy [37, 48].

Conclusion

Modification of drug dosage in obese patients is very important, particularly when using medications with narrow therapeutic index. Alterations of aminoglycoside pharmacokinetic parameters in obesity may necessitate deviation from dosages commonly recommended for non-obese individuals, but knowledge on the influence of obesity on pharmacokinetics is limited. This review shows that currently available literature supports the need for aminoglycoside dosage modification in critically ill obese patients. Physicians need to take every possible step to ensure that aminoglycoside doses are adequate and safe. Because pathophysiologic changes in obesity can result in sub- or supra-therapeutic concentrations, aminoglycoside doses should be re-evaluated daily in critically ill obese patients, and TDM can help optimize therapy. Clinicians prescribing antibiotics for critically ill obese patients should be familiar with drug and patient factors influencing the effectiveness of antibiotic therapy.

Financial Disclosures

This manuscript was supported solely by Department Funds.

Conflicts of Interest

All authors state that they do not have any conflicts of interest to report.

References

- Schwartz SN, Pazin GJ, Lyon JA, Ho M, Pasculle AW. A controlled investigation of the pharmacokinetics of gentamicin and tobramycin in obese subjects. *J Infect Dis.* 1978;138(4):499-505.
- Blouin RA, Mann HJ, Griffen WO, Jr., Bauer LA, Record KE. Tobramycin pharmacokinetics in morbidly obese patients. *Clin Pharmacol Ther.* 1979;26(4):508-512.
- Bauer LA, Blouin RA, Griffen WO, Jr., Record KE, Bell RM. Amikacin pharmacokinetics in morbidly obese patients. *Am J Hosp Pharm.* 1980;37(4):519-522.
- Korsager S. Administration of gentamicin to obese patients. *Int J Clin Pharmacol Ther Toxicol.* 1980;18(12):549-553.
- Bauer LA, Edwards WA, Dellinger EP, Simonowitz DA. Influence of weight on aminoglycoside pharmacokinetics in normal weight and morbidly obese patients. *Eur J Clin Pharmacol.* 1983;24(5):643-647.
- Janmahasatian S, Duffull SB, Ash S, Ward LC, Byrne NM, Green B. Quantification of lean bodyweight. *Clin Pharmacokinet.* 2005;44(10):1051-1065.
- Pai MP, Nafziger AN, Bertino JS, Jr. Simplified estimation of aminoglycoside pharmacokinetics in underweight and obese adult patients. *Antimicrob Agents Chemother.* 2011;55(9):4006-4011.
- Traynor AM, Nafziger AN, Bertino JS, Jr. Aminoglycoside dosing weight correction factors for patients of various body sizes. *Antimicrob Agents Chemother.* 1995;39(2):545-548.
- Devine BJ. Gentamicin therapy. *Drug Intell Clin Pharm.* 1974;8:650-655.
- Wurtz R, Itokazu G, Rodvold K. Antimicrobial dosing in obese patients. *Clin Infect Dis.* 1997;25(1):112-118.
- Leader WG, Tsubaki T, Chandler MH. Creatinine-clearance estimates for predicting gentamicin pharmacokinetic values in obese patients. *Am J Hosp Pharm.* 1994;51(17):2125-2130.
- Pai MP, Bearden DT. Antimicrobial dosing considerations in obese adult patients. *Pharmacotherapy.* 2007;27(8):1081-1091.
- Blouin RA, Brouwer KL, Record KE, Griffen WO, Jr., Plezia PM, John W. Amikacin pharmacokinetics in morbidly obese patients undergoing gastric-bypass surgery. *Clin Pharm.* 1985;4(1):70-72.
- Casati A, Putzu M. Anesthesia in the obese patient: pharmacokinetic considerations. *J Clin Anesth.* 2005;17(2):134-145.
- Erstad BL. Dosing of medications in morbidly obese patients in the intensive care unit setting. *Intensive Care Med.* 2004;30(1):18-32.
- Cheyamol G. Effects of obesity on pharmacokinetics implications for drug therapy. *Clin Pharmacokinet.* 2000;39(3):215-231.
- Corcoran GB, Salazar DE, Schentag JJ. Excessive aminoglycoside nephrotoxicity in obese patients. *Am J Med.*

- 1988;85(2):279.
18. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron*. 1976;16(1):31-41.
 19. Salazar DE, Corcoran GB. Predicting creatinine clearance and renal drug clearance in obese patients from estimated fat-free body mass. *Am J Med*. 1988;84(6):1053-1060.
 20. Ali MZ, Goetz MB. A meta-analysis of the relative efficacy and toxicity of single daily dosing versus multiple daily dosing of aminoglycosides. *Clin Infect Dis*. 1997;24(5):796-809.
 21. Bailey TC, Little JR, Littenberg B, Reichley RM, Dunagan WC. A meta-analysis of extended-interval dosing versus multiple daily dosing of aminoglycosides. *Clin Infect Dis*. 1997;24(5):786-795.
 22. Barza M, Ioannidis JP, Cappelleri JC, Lau J. Single or multiple daily doses of aminoglycosides: a meta-analysis. *BMJ*. 1996;312(7027):338-345.
 23. Ferriols-Lisart R, Alos-Alminana M. Effectiveness and safety of once-daily aminoglycosides: a meta-analysis. *Am J Health Syst Pharm*. 1996;53(10):1141-1150.
 24. Munchhof WJ, Grayson ML, Turnidge JD. A meta-analysis of studies on the safety and efficacy of aminoglycosides given either once daily or as divided doses. *J Antimicrob Chemother*. 1996;37(4):645-663.
 25. Blouin RA, Chandler MH. Special pharmacokinetic considerations in the obese. In: Evans WE, Schentag JJ, Jusko WJ, editors. *Applied pharmacokinetics: principles of therapeutic drug monitoring*. Vancouver, BC: Applied Therapeutics Inc, 1992:3-20.
 26. Medico CJ, Walsh P. Pharmacotherapy in the critically ill obese patient. *Crit Care Clin*. 2010;26(4):679-688.
 27. Hanley MJ, Abernethy DR, Greenblatt DJ. Effect of obesity on the pharmacokinetics of drugs in humans. *Clin Pharmacokinet*. 2010;49(2):71-87.
 28. Park GR. Molecular mechanisms of drug metabolism in the critically ill. *Br J Anaesth*. 1996;77(1):32-49.
 29. Bearden DT, Rodvold KA. Dosage adjustments for antibacterials in obese patients: applying clinical pharmacokinetics. *Clin Pharmacokinet*. 2000;38(5):415-426.
 30. Vincent F, El-Khoury N, Rondeau E. Renal function in critically ill, morbidly obese patients. *Am J Respir Crit Care Med*. 2004;169(12):1332; author reply 1333.
 31. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, Reinhart K, et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med*. 2008;36(1):296-327.
 32. McKenzie C. Antibiotic dosing in critical illness. *J Antimicrob Chemother*. 2011;66:(Suppl 2)ii25-31.
 33. Moore RD, Lietman PS, Smith CR. Clinical response to aminoglycoside therapy: importance of the ratio of peak concentration to minimal inhibitory concentration. *J Infect Dis*. 1987;155(1):93-99.
 34. Triginer C, Izquierdo I, Fernandez R, Rello J, Torrent J, Benito S, Net A. Gentamicin volume of distribution in critically ill septic patients. *Intensive Care Med*. 1990;16(5):303-306.
 35. Rea RS, Capitano B. Optimizing use of aminoglycosides in the critically ill. *Semin Respir Crit Care Med*. 2007;28(6):596-603.
 36. Taccone FS, Laterre PF, Spapen H, Dugernier T, Delattre I, Layeux B, De Backer D, et al. Revisiting the loading dose of amikacin for patients with severe sepsis and septic shock. *Crit Care*. 2010;14(2):R53.
 37. Pai MP, Rodvold KA. Aminoglycoside dosing in patients by kidney function and area under the curve: the Sawchuk-Zaske dosing method revisited in the era of obesity. *Diagn Microbiol Infect Dis*. 2014;78(2):178-187.
 38. Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med*. 1999;130(6):461-470.
 39. Poggio ED, Nef PC, Wang X, Greene T, Van Lente F, Dennis VW, Hall PM. Performance of the Cockcroft-Gault and modification of diet in renal disease equations in estimating GFR in ill hospitalized patients. *Am J Kidney Dis*. 2005;46(2):242-252.
 40. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF, 3rd, Feldman HI, Kusek JW, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009;150(9):604-612.
 41. Stevens LA, Schmid CH, Greene T, Zhang YL, Beck GJ, Froissart M, Hamm LL, et al. Comparative performance of the CKD Epidemiology Collaboration (CKD-EPI) and the Modification of Diet in Renal Disease (MDRD) Study equations for estimating GFR levels above 60 mL/min/1.73 m². *Am J Kidney Dis*. 2010;56(3):486-495.
 42. Livio F, Biollaz J, Burnier M. [Renal function estimation by MDRD equation: interest and limitations for drug dosing]. *Rev Med Suisse*. 2008;4(181):2596-2600.
 43. Blaser J. [Dosing aminoglycosides once a day]. *Schweiz Med Wochenschr Suppl*. 1996;76:39S-48S.
 44. Freeman CD, Nicolau DP, Belliveau PP, Nightingale CH. Once-daily dosing of aminoglycosides: review and recommendations for clinical practice. *J Antimicrob Chemother*. 1997;39(6):677-686.
 45. Ross AL, Tharp JL, Hobbs GR, McKnight R, Cumpston A. Evaluation of extended interval dosing aminoglycosides in the morbidly obese population. *Adv Pharmacol Sci*. 2013;2013:194389.
 46. Buijk SE, Mouton JW, Gyssens IC, Verbrugh HA, Bruining HA. Experience with a once-daily dosing program of aminoglycosides in critically ill patients. *Intensive Care Med*. 2002;28(7):936-942.

47. Buchholz U, Richards C, Murthy R, Arduino M, Pon D, Schwartz W, Fontanilla E, et al. Pyrogenic reactions associated with single daily dosing of intravenous gentamicin. *Infect Control Hosp Epidemiol.* 2000;21(12):771-774.
48. Begg EJ, Barclay ML, Kirkpatrick CM. The therapeutic monitoring of antimicrobial agents. *Br J Clin Pharmacol.* 2001;52:(Suppl 1)35S-43S.