

2014

## Common exposure to STL polyomavirus during childhood

Efrem S. Lim

*Washington University School of Medicine in St. Louis*

Natalie M. Meinerz

*University of Colorado Boulder*

Blake Primi

*University of Colorado Boulder*

David Wang

*Washington University School of Medicine in St. Louis*

Robert L. Garcea

*University of Colorado Boulder*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Lim, Efrem S.; Meinerz, Natalie M.; Primi, Blake; Wang, David; and Garcea, Robert L., "Common exposure to STL polyomavirus during childhood." *Emerging Infectious Diseases*. 20, 9. 1559-61. (2014).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/3541](https://digitalcommons.wustl.edu/open_access_pubs/3541)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

# Common Exposure to STL Polyomavirus During Childhood

Efrem S. Lim, Natalie M. Meinerz, Blake Primi, David Wang, and Robert L. Garcea

STL polyomavirus (STLPyV) was recently identified in human specimens. To determine seropositivity for STLPyV, we developed an ELISA and screened patient samples from 2 US cities (Denver, Colorado [500]; St. Louis, Missouri [419]). Overall seropositivity was 68%–70%. The age-stratified data suggest that STLPyV infection is widespread and commonly acquired during childhood.

Polyomaviruses are nonenveloped double-stranded circular DNA viruses that infect a wide range of hosts, including humans. The capsid of the virus comprises primarily 72 pentamers of the major coat protein, VP1. Human polyomaviruses have been associated with several diseases (1). BK polyomavirus (BKPyV) has been associated with nephropathy in renal transplant recipients and JC polyomavirus (JCPyV) with progressive multifocal leukoencephalopathy in immunocompromised persons (2,3). Trichodysplasia spinulosa-associated polyomavirus (TSPyV) infection is linked to a rare skin disease in immunocompromised patients called trichodysplasia spinulosa (4). Furthermore, infection with Merkel cell polyomavirus (MCPyV) in rare instances leads to Merkel cell carcinomas, an aggressive form of skin cancer (5). Other polyomaviruses, including WU polyomavirus (WUPyV), KI polyomavirus (KIPyV), human polyomavirus 6, human polyomavirus 7, human polyomavirus 9, MW polyomavirus (MWPyV), STL polyomavirus (STLPyV), and human polyomavirus 12, have been identified in specimens from humans, but their role in disease remains to be defined (1).

Seroepidemiology has played an important role in establishing the link between human polyomaviruses and disease and in understanding infection dynamics. The seroprevalences of BKPyV and JCPyV range from 70% to 90% and 40% to 60%, respectively, with an age profile indicating high frequency of early-age infections and lifelong persistence (6–8). Seropositivity for MCPyV in healthy

persons ranges from 25% to 64%; all patients with Merkel cell carcinoma are seropositive (6,9).

STLPyV was recently identified from fecal specimens from a child in Malawi (10). Viral DNA also was detected in fecal specimens from the United States and The Gambia, and STLPyV has been found in a surface-sanitized skin wart surgically removed from the buttocks of a patient with a primary immunodeficiency called WHIM (warts, hypogammaglobulinemia, infections, and myelokathexis) syndrome (11). These observations suggest that STLPyV might infect humans. We defined the seropositivity rate of STLPyV in humans using serum from 2 independent US sites (Denver, Colorado, and St. Louis, Missouri).

## The Study

To determine the seropositivity for STLPyV, we developed a capture ELISA using recombinant glutathione S-transferase-tagged STLPyV VP1 capsomeres (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/20/9/14-0561-Techapp1.pdf>). Electron microscopy of the STLPyV capsomeres showed 10-nm pentamers characteristic of polyomaviruses (Figure 1, panel A). We assessed the specificity of the STLPyV ELISA by preincubating 24 serum samples in the presence and absence of soluble STLPyV VP1 pentamers before addition to the immobilized STLPyV glutathione S-transferase-tagged VP1. The ELISA signal intensity was markedly reduced when serum was preincubated with STLPyV VP1 pentamers (Figure 1, panel B; compare white bars to gray bars). This result indicates the ELISA seroreactivity could be self-competed with soluble STLPyV pentamers. STLPyV shares 55% aa identity in the VP1 region with its next most closely related polyomavirus, MWPyV (10). Therefore, we examined whether cross-reactivity existed between STLPyV and MWPyV VP1 capsomeres. Competition assays with soluble MWPyV VP1 pentamers showed limited interference with the ELISA seroreactivity (Figure 1, panel C; compare white bars to gray bars). This result indicates that there was no significant cross-reactivity between STLPyV and MWPyV VP1 capsomeres. Taken together, these data demonstrate that the ELISA was specific to STLPyV VP1.

We screened 500 serum specimens collected from children and adults in Denver for antibodies against STLPyV. The overall seropositivity for STLPyV was 68.0% (Figure 2, panel A). Children 1–3 years of age had the lowest seropositivity rate (23.8%). In contrast, seroprevalence of children 4–20 years of age ranged from 61.1% to 70.8%. Similar seropositivity rates persisted in adults (>21 years of age), ranging from 68.8% to 74.2%.

We next examined a panel of 419 serum specimens from St. Louis that had a higher resolution of age-stratification in young infants. The overall seropositivity for STLPyV was 70.0% (Figure 2, panel B), similar to that

Author affiliations: Washington University, St. Louis, Missouri, USA (E.S. Lim, D. Wang); and University of Colorado, Boulder, Colorado, USA (N.M. Meinerz, B. Primi, R.L. Garcea)

DOI: <http://dx.doi.org/10.3201/eid2009.140561>




Figure 1. STL polyomavirus (STLPyV) ELISA. A) Electron microscopy image shows purified STLPyV VP1 capsomeres. Scale bar = 100 nm. B) Serum samples were pre-incubated in the absence (white bars) or presence of soluble STLPyV VP1 pentamers (gray bars), followed by the STLPyV-capture ELISA. Serum was tested in triplicate, and average absorbance values are shown. Error bars indicate SD. Representative data are shown. C) Seroreactivity to STLPyV in the absence (white bars) or presence of competition with MW polyomavirus (MWPyV) VP1 pentamers (gray bars) are shown. Serum was tested in triplicate, and average absorbance values are shown. Error bars indicate SD. Representative data are shown.

in the specimens from Denver (68.0%). Seropositivity for STLPyV in infants dropped from 53.3% in the <0.5-year age group to 37.9% in the 0.5–1-year age group, with the lowest seropositivity of 22.6% in the 1–2-year group. In contrast, seropositivity rates for children >2 years of age were higher, ranging from 60.0% to 85.3%. Finally, seropositivity in adults (>21 years of age) ranged from 91.2% to 95.2%. Thus, these data indicate that exposure to STLPyV occurs during early childhood and that immune responses to STLPyV are maintained in adults.




Figure 2. Age-specific seropositivity for STL polyomavirus (STLPyV) from serum specimens collected in Denver, Colorado, USA (A), and St. Louis, Missouri, USA (B). A total of 500 serum specimens from Denver and 417 from St. Louis were tested for seroreactivity to STLPyV VP1 proteins. Overall seropositivity in Denver was 68% and in St. Louis, 70%. Error bars indicate SD.

## Conclusions

In our analysis of the seroepidemiology of STLPyV in 2 areas of the United States, we found that prevalence of the virus was similar (68.0%–70.0%). This prevalence is slightly higher than the 41.8% for MWPyV, the polyomavirus most closely related to STLPyV (12). However, the seropositivity of STLPyV is comparable to other human polyomaviruses (>60%), such as BKPyV, KIPyV, WUPyV, MCPyV, human polyomavirus 6, and TSPyV (6,8). We found no cross-reactivity with MWPyV VP1, the most closely-related polyomavirus that shares 55% aa identity in the VP1 region. Thus, the seroepidemiology strongly supports the notion that STLPyV is a bona fide infectious agent of humans.

Age stratification of the seropositive specimens suggested an initial waning of immune response followed by rapid seroconversion during childhood. In the St. Louis specimens, seropositivity was higher for the <0.5-year and 0.5–1-year age groups (53.3% and 37.9%, respectively) than for the 1–2-year group (22.6%). This observation was followed by an increase in STLPyV seropositivity in the 2–3-year group and older age groups (60.0%). Specimens from Denver were too few to reliably stratify the data to

the same resolution. Nonetheless, we observed a similar trend in specimens from Denver where the seropositivity for the 1–3-year age group (23.8%) was lower than for the 4–6-year group (61.1%). These data indicate that immune responses to STLPyV decreased in the first 2 years of age, which suggests waning maternal antibodies. However, seropositivity was rapidly acquired thereafter, indicating high exposure of STLPyV in children. Because the specimens were selected from hospital-associated blood draws of children of unknown health status, seropositivity rates for healthy children might have varied somewhat from the results we obtained. Nonetheless, the trends we observed with STLPyV are similar to profiles that have been reported for JCPyV, BKPyV, TSPyV, MCPyV, WUPyV, and KIPyV (6,13–15).

STLPyV has not been clearly associated with any disease. However, the high overall seropositivity rate suggests widespread infection in the population at large. None of the pathogenic human polyomaviruses (JCPyV, BKPyV, MCPyV, TSPyV) have been clearly associated with acute disease at the time of initial infection. Rather, immunosuppression is a critical co-factor that is coupled to the ability of polyomaviruses to persist throughout life or to integrate into the genome, as in the case of MCPyV, ultimately leading to disease. Our study demonstrates that a large segment of the general population has been infected by STLPyV and might harbor persistent STLPyV infection, assuming the persistence paradigms of JCPyV and BKPyV hold true for STLPyV. A recent report identified STLPyV DNA in a skin wart specimen from an adult with primary immunodeficiency (11). Thus, it is critical to determine whether human diseases exist that are caused by STLPyV, especially in immunocompromised persons.

R.L.G. was supported by National Institutes of Health grant CA37667. D.W. holds an investigator in the pathogenesis of infectious disease award from the Burroughs Wellcome Fund. E.S.L. is an Eli and Edythe Broad Fellow of the Life Sciences Research Foundation.

Dr Lim is a postdoctoral research associate at Washington University in St. Louis. His research focuses on the discovery and characterization of novel viruses and human pathogens.

## References

- DeCaprio JA, Garcea RL. A cornucopia of human polyomaviruses. *Nat Rev Microbiol.* 2013;11:264–76. <http://dx.doi.org/10.1038/nrmicro2992>
- Ferency MW, Marshall LJ, Nelson CD, Atwood WJ, Nath A, Khalili K, et al. Molecular biology, epidemiology, and pathogenesis of progressive multifocal leukoencephalopathy, the JC virus-induced demyelinating disease of the human brain. *Clin Microbiol Rev.* 2012;25:471–506. <http://dx.doi.org/10.1128/CMR.05031-11>
- Kuypers DR. Management of polyomavirus-associated nephropathy in renal transplant recipients. *Nat Rev Nephrol.* 2012;8:390–402. <http://dx.doi.org/10.1038/nrneph.2012.64>
- van der Meijden E, Janssens RW, Lauber C, Bouwes Bavinck JN, Gorbalenya AE, Feltkamp MC. Discovery of a new human polyomavirus associated with trichodysplasia spinulosa in an immunocompromised patient. *PLoS Pathog.* 2010;6:e1001024. <http://dx.doi.org/10.1371/journal.ppat.1001024>
- Feng H, Shuda M, Chang Y, Moore PS. Clonal integration of a polyomavirus in human Merkel cell carcinoma. *Science.* 2008;319:1096–100. <http://dx.doi.org/10.1126/science.1152586>
- Kean JM, Rao S, Wang M, Garcea RL. Seroepidemiology of human polyomaviruses. *PLoS Pathog.* 2009;5:e1000363. <http://dx.doi.org/10.1371/journal.ppat.1000363>
- Viscidi RP, Rollison DE, Sondak VK, Silver B, Messina JL, Giuliano AR, et al. Age-specific seroprevalence of Merkel cell polyomavirus, BK virus, and JC virus. *Clin Vaccine Immunol.* 2011;18:1737–43. <http://dx.doi.org/10.1128/CVI.05175-11>
- Moens U, Van Ghelue M, Song XB, Ehlers B. Serological cross-reactivity between human polyomaviruses. *Rev Med Virol.* 2013;23:250–64. <http://dx.doi.org/10.1002/rmv.1747>
- Tolstov YL, Pastrana DV, Feng HC, Becker JC, Jenkins FJ, Moschos S, et al. Human Merkel cell polyomavirus infection II. MCV is a common human infection that can be detected by conformational capsid epitope immunoassays. *Int J Cancer.* 2009;125:1250–6. <http://dx.doi.org/10.1002/ijc.24509>
- Lim ES, Reyes A, Antonio M, Saha D, Ikumapayi UN, Adeyemi M, et al. Discovery of STL polyomavirus, a polyomavirus of ancestral recombinant origin that encodes a unique T antigen by alternative splicing. *Virology.* 2013;436:295–303. <http://dx.doi.org/10.1016/j.virol.2012.12.005>
- Pastrana DV, Fitzgerald PC, Phan GQ, Raiji MT, Murphy PM, McDermott DH, et al. A divergent variant of the eleventh human polyomavirus species, St. louis polyomavirus. *Genome Announc.* 2013;1:e00812–3. <http://dx.doi.org/10.1128/genomeA.00812-13>
- Nicol JT, Leblond V, Arnold F, Guerra G, Mazzoni E, Tognon M, et al. Seroprevalence of human Malawi polyomavirus. *J Clin Microbiol.* 2014;52:321–3. <http://dx.doi.org/10.1128/JCM.02730-13>
- Nguyen NL, Le BM, Wang D. Serologic evidence of frequent human infection with WU and KI polyomaviruses. *Emerg Infect Dis.* 2009;15:1199–205. <http://dx.doi.org/10.3201/eid1508.090270>
- Martel-Jantin C, Pedergnana V, Nicol JT, Leblond V, Trégouët DA, Tortevoye P, et al. Merkel cell polyomavirus infection occurs during early childhood and is transmitted between siblings. *J Clin Virol.* 2013;58:288–91. <http://dx.doi.org/10.1016/j.jcv.2013.06.004>
- van der Meijden E, Bialasiewicz S, Rockett RJ, Tozer SJ, Sloots TP, Feltkamp MC. Different serologic behavior of MCPyV, TSPyV, HPyV6, HPyV7 and HPyV9 polyomaviruses found on the skin. *PLoS ONE.* 2013;8:e81078. <http://dx.doi.org/10.1371/journal.pone.0081078>

Address for correspondence: Robert L. Garcea, University of Colorado, School of Medicine, Molecular, Cellular, & Developmental Biology, and The BioFrontiers Institute, Jennie Smoly Caruthers Biotechnology Building, Campus Box 596, Boulder, CO 80309, USA; email: Robert.Garcea@colorado.edu



## GovDelivery

Manage your email alerts so you only receive content of interest to you.

Sign up for an online subscription:  
<http://wwwnc.cdc.gov/eid/subscribe.htm>