

2015

## A twin study of breastfeeding with a preliminary genome-wide association scan

Lucia Colodro-Conde  
*University of Murcia*

Gu Zhu  
*Berghofer Medical Research Institute*

Robert A. Power  
*King's College London - Institute of Psychiatry*

Anjali Henders  
*Berghofer Medical Research Institute*

Andrew C. Heath  
*Washington University School of Medicine in St. Louis*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Colodro-Conde, Lucia; Zhu, Gu; Power, Robert A.; Henders, Anjali; Heath, Andrew C.; Madden, Pamela A F; Montgomery, Grant W.; Medland, Sarah; Ordoñana, Juan R.; and Martin, Nicholas G., "A twin study of breastfeeding with a preliminary genome-wide association scan." *Twin Research and Human Genetics*. 18, 1. 61-72. (2015).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/3737](https://digitalcommons.wustl.edu/open_access_pubs/3737)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

---

**Authors**

Lucia Colodro-Conde, Gu Zhu, Robert A. Power, Anjali Henders, Andrew C. Heath, Pamela A F Madden, Grant W. Montgomery, Sarah Medland, Juan R. Ordoñana, and Nicholas G. Martin

# A Twin Study of Breastfeeding With a Preliminary Genome-Wide Association Scan

Lucia Colodro-Conde,<sup>1,2</sup> Gu Zhu,<sup>1</sup> Robert A. Power,<sup>3</sup> Anjali Henders,<sup>1</sup> Andrew C. Heath,<sup>4</sup> Pamela A. F. Madden,<sup>4</sup> Grant W. Montgomery,<sup>1</sup> Sarah Medland,<sup>1</sup> Juan R. Ordoñana,<sup>2</sup> and Nicholas G. Martin<sup>1</sup>

<sup>1</sup>QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia

<sup>2</sup>Murcia Twin Registry, Department of Human Anatomy and Psychobiology, University of Murcia, IMIB-Arrixaca, Murcia, Spain

<sup>3</sup>MRC Social Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, King's College London, DeCrespigny Park, Denmark Hill, London, UK

<sup>4</sup>School of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA

Breastfeeding has been an important survival trait during human history, though it has long been recognized that individuals differ in their exact breastfeeding behavior. Here our aims were, first, to explore to what extent genetic and environmental influences contributed to the individual differences in breastfeeding behavior; second, to detect possible genetic variants related to breastfeeding; and lastly, to test if the genetic variants associated with breastfeeding have been previously found to be related with breast size. Data were collected from a large community-based cohort of Australian twins, with 3,364 women participating in the twin modelling analyses and 1,521 of them included in the genome-wide association study (GWAS). Monozygotic (MZ) twin correlations ( $r_{MZ} = 0.52$ , 95% CI 0.46–0.57) were larger than dizygotic (DZ) twin correlations ( $r_{DZ} = 0.35$ , 95% CI 0.25–0.43) and the best-fitting model was the one composed by additive genetics and unique environmental factors, explaining 53% and 47% of the variance in breastfeeding behavior, respectively. No breastfeeding-related genetic variants reached genome-wide significance. The polygenic risk score analyses showed no significant results, suggesting breast size does not influence breastfeeding. This study confers a replication of a previous one exploring the sources of variance of breastfeeding and, to our knowledge, is the first one to conduct a GWAS on breastfeeding and look at the overlap with variants for breast size.

■ **Keywords:** breastfeeding, twin study, genome-wide association study

Breast milk, either from an infant's own mother or from another woman, can be assumed to have been the sole source of nourishment for infants during most of human evolution, having evolved as an adaptation to transfer immune factors to offspring and to space births (Sellen, 2007). It has been suggested that breastfeeding duration in prehistoric times was of two–three years and that from the Middle Ages to the 19th century, most infants in Europe were typically weaned between the ages of one and two years (Schön & Silvé, 2007). It was not until the end of the 19th century, that is, in our recent history, that artificial feeding became a safe alternative to breast milk, and breastfeeding became dependent on the mother's choice in most milieus. Consequently, nowadays a large variability in this behavior can be observed both within and between countries, with

global statistics showing that 37% of infants were exclusively breastfed until six months age in 2007 (World Health Organization, 2013), which is the current recommended minimum weaning age (World Health Organization & United Nations Children's Fund, 2003). This percentage varied considerably across countries; for example, 18.8% in the United States in 2011 (Centers for Disease Control and Prevention, 2014), 17.6% in Australia in 2011–2012 (Australian Bureau of Statistics, 2013) and 28.5% in Spain in 2011–2012 (Min-

RECEIVED 14 July 2014; ACCEPTED 29 September 2014.

ADDRESS FOR CORRESPONDENCE: Lucia Colodro-Conde, QIMR Berghofer Medical Research Institute, 300 Herston Rd, Herston QLD 4006, Australia. E-mail: [lucia.c.c@um.es](mailto:lucia.c.c@um.es)

isterio de Sanidad, Servicios Sociales e Igualdad, 2012). The instruments for data collection vary between countries, for example, in being explicit about the inclusion of expressed milk in exclusive breastfeeding, so some rates could be higher.

Given the positive impact of breastfeeding on infants' and mothers' health (Ip et al., 2009; World Health Organization & United Nations Children's Fund, 2003), there is a current interest in determining which factors are related to women's decisions regarding the method of feeding their infants, that is, the predispositions or situations that make them more likely to initiate breastfeeding or bottle feeding, and the duration of breastfeeding. Breastfeeding is a biocultural behavior (Holman & Grimes, 2003) and is influenced by multiple factors (Thulier & Mercer, 2009), such as the action of prolactin and oxytocin (Heinig & Dewey, 1997), pain from nipple trauma, mastitis, maternal exhaustion and perceived poor milk supply (Almqvist-Tangen et al., 2012; Spencer, 2008), babies' sucking dynamics (Sakalidis et al., 2013), body mass index (Wojcicki, 2011), level of education (Colodro-Conde et al., 2011), working conditions (Calnen, 2010), psychological factors, personality, self-efficacy, anxiety (Brown, 2014; Colodro-Conde et al., in press (a); Colodro-Conde et al., in press (b); De Jager et al., 2012; Li et al., 2008; McFadden & Toole, 2006; Wagner et al., 2006), support from partner, family and peers (Thulier & Mercer, 2009), social norms (Swanson & Power, 2005), and advice from health professionals (Brown et al., 2011).

The magnitude of the genetic and environmental sources of variance in breastfeeding behavior were explored in a previous study in a cohort of Spanish twins from the Murcia Twin Registry (Ordoñana et al., 2006, 2013). The best-fitting models explained the observed variance through additive genetic and non-shared environmental factors for initiation and duration in the first-born child and the average for the complete offspring, with heritability ranging between 44% and 54% (Colodro-Conde et al., 2013). It was hypothesized that among the genetic factors influencing breastfeeding behavior, those related to the regulation of hormone production could play a part in breastfeeding outcomes. In support of this hypothesis, Jonas et al. (2013) found that the single nucleotide polymorphism (SNP) rs2740210 in the oxytocin peptide gene was associated with exclusive breastfeeding at three and six months postpartum. Other authors, however, did not find a relationship between selected oxytocin receptor SNPs and breastfeeding (Tharner et al., 2012). As breastfeeding has been suggested to be significantly influenced by genetic factors, it would be interesting to explore the genetic variants underlying this trait.

Several studies have concluded that body mass index is associated with breastfeeding behavior, so mothers who are overweight or obese are less likely to initiate lactation, and are prone to early cessation of breastfeeding (Jevitt et al., 2007; Wojcicki, 2011). A study with health professionals showed that from their experience, initiating was more dif-

ficult than continuing breastfeeding for women with large breasts and/or obesity (Katz et al., 2010). The link between overweight, elevated serum testosterone concentration and pathologies such as the polycystic ovary syndrome could be underlying this relationship (Balen et al., 1995; Barber et al., 2006). Heritability of breast cup size has been estimated to be 56%, and one third of this variance was shared with body mass index (Wade et al., 2010). Some genetic variants associated with breast size also influence breast cancer risk (Eriksson et al., 2012).

This article has three main objectives. First, it aims to replicate previous findings related to heritability of breastfeeding by exploring the magnitude of the genetic and environmental influences in the variation of breastfeeding behavior in a sample of Australian twins. For that aim, we used a twin design, calculating twin correlations for breastfeeding and fitting an ACE variance components model. Second, in order to detect possible genetic variants related to breastfeeding behavior, we conducted a GWAS of this phenotype, using data collected from this large community-based cohort of Australian twins. Lastly, we report the results of a polygenic risk score analysis examining whether an individual's number of genetic variants predisposing to breast size are associated with breastfeeding.

## Materials and Methods

### Participants

Participants in this study were 3,364 female twin mothers from the QIMR health and lifestyle studies Cohorts I and II with data on breastfeeding. Women in Cohort I were born between 1892 and 1963 ( $n = 3,205$ ) and women in Cohort II were born between 1,964 and 1971 ( $n = 419$ ). At the time of the survey (1988–93), the mean age was 43.41 years ( $SD = 12.32$ , range = 25–86 years), for women in Cohort I and 24.81 years ( $SD = 1.87$ , range = 19–29 years) for women in Cohort II. Further details of the sample, data collection and zygosity determination are described elsewhere for cohort I (Heath et al., 1997) and for cohort II (Knopik et al., 2004).

The sample comprised 992 complete twin pairs: 629 MZ and 363 DZ, and 1,380 individual twins from incomplete pairs (411 MZ and 347 DZ from female-female pairs, 621 from female-male pairs and 5 of unknown zygosity). Genome-wide genotypic data were available for 1,521 of the 3,364 individuals.

### Procedure

In 1988–1993, participants of both cohorts completed a mailed questionnaire including information about their childbearing. Some of these who did not complete the mailed questionnaire were interviewed over the telephone in 1989–1992. A small subsample of participants of Cohort I ( $n = 341$ ) was re-surveyed two years after their initial

contact to establish the test-retest reliability of the interview measures.

As part of follow-up studies, blood samples were collected from the majority of participants in Cohorts I and II. DNA samples were genotyped using the Illumina 317, 370 and 610 SNP chips. Following quality control, the data were imputed using the subset of ~281,000 markers, which were available across all chips. The genotypes were phased using MACH, then imputed (including the X chromosome) using Minimac based on the 283 European reference individuals in the March 2013 release of phased haplotype from the 1000 Genomes project. In the analyses presented here, we used data from 6.59 M imputed SNPs that passed quality control ( $R^2 > 0.3$  and minor allele frequency  $> 1\%$ ) for association analysis. Details of SNP typing, quality control, data cleaning (including ancestry exclusion) and analysis are given in detail elsewhere (Medland et al., 2009).

This study was approved by the Queensland Institute of Medical Research Human Research Ethics Committee and the storage of the data follows national regulations regarding personal data protection. All of the participants provided informed consent.

### Measures

Data were based on retrospective self-reports. All the participants were asked about details of the birth of up to eight children, including birth complications, time of birth, birth weight, hours of labor, feelings of depression and type of feeding in the first month (see Appendix). The variable we analyzed was the mean number of months they breastfed each child, averaged across all live births and standardized to a Z-score.

### Data Analysis

**Twin modelling.** Data preparation and descriptive and preliminary analyses were performed in SPSS v.19 (SPSS, 2010). Assumptions of the twin design were checked, including the homogeneity of the means and variances of first- and second-born twins and across zygosity groups. Further details of the twin design, including checking at assumptions, can be found elsewhere (Neale & Cardon, 1992; Posthuma et al., 2003).

Statistical analyses employed full information maximum-likelihood modelling (FIML) procedures using the statistical package Mx (Neale et al., 2006). In FIML, both complete and incomplete pairs of twins can be used in the analyses and the goodness-of-fit of a model to the observed data is distributed as chi-square ( $\chi^2$ ). By testing the change in chi-square ( $\Delta\chi^2$ ) against the change in degrees of freedom ( $\Delta df$ ), we can test whether dropping or equating specific model parameters significantly worsens the model fit. The best-fitting model was chosen in each case by deducting the residual deviance of the compared

models and by comparing Akaike's information criterion (AIC).

In a first step, we determined twin pair correlations per zygosity group for breastfeeding. We tested whether MZ correlations were higher than those of DZ twin pairs, which would suggest a genetic influence on individual differences in this trait. Then, variance component model-fitting was conducted to partition the variation in breastfeeding into genetic, shared environmental and non-shared environmental influences. Observed MZ and DZ twin correlations generally reflect a combination of A, C, D, and E influences, and structural equation modeling determines the combination that best matches the observed data (Posthuma et al., 2003). Age at the survey time was modelled as covariate in all the analyses.

**SNP-based genome-wide association study.** One thousand genome imputed dosage data were tested for association with breastfeeding using the additive family-based association test in Merlin-off line (Chen & Abecasis, 2007). Correction for age at the survey time was performed by fitting the covariate in the regression model. We adopted a genome-wide significance level for the association between SNP and phenotype of  $5 \times 10^{-8}$  or smaller to correct for the total number of independent tests (Dudbridge & Gusnanto, 2008).

**Gene-based analysis.** A gene-based test, Versatile Gene-based Association Study (VEGAS), designed for use with GWAS data with related individuals (Liu et al., 2010), was conducted to determine the level of association across the gene while correcting for linkage disequilibrium (LD) and gene size. Details of this procedure are summarized elsewhere (Verweij et al., 2010). In brief, this test explores association of each gene (including 50 Kb up and down of the coding region) taking into account the  $p$  values of all SNPs (after first pruning out those in high LD) ( $r^2 > 0.20$ ), and the LD between them. A  $p$  value below  $\alpha = 2.8 \times 10^{-6}$  was considered to be genome-wide significant as the gene-based association test included 17,585 genes (0.05/17,585 genes).

**Polygenic study.** To examine a potential shared genetic etiology between breastfeeding and breast size, we tested the effects of SNPs previously associated with breast size through a polygenic risk scoring approach. This took the top 2,079 SNPs associated with breast size in a previous GWAS in an independent sample (Eriksson et al., 2012) at a  $p$  value of  $\leq .0001$ . These were then clumped within PLINK (Purcell et al., 2007) to correct for LD within the QIMR sample ( $R^2 < 0.2$  in 250 kb windows), identifying 66 overlapping independent SNPs associated with breast size. A polygenic score was generated using PLINK weighting on the beta for each SNP. This score was then tested for association with breastfeeding phenotypes, correcting

**TABLE 1**  
Breastfeeding Duration (Months)  
According to Childbirth Order

Child order	n	Mean	SD	Range
1st child	3,295	5.32	4.74	0–48
2nd child	2,568	5.19	5.17	0–48
3rd child	1,362	5.24	5.51	0–36
4th child	547	5.67	6.63	0–60
5th child	209	4.93	5.99	0–36
6th child	76	3.93	4.44	0–20
7th child	25	4.76	8.07	0–40
8th child	8	4.12	3.44	0–9

for covariates of five principal components. Variance explained by this score was derived as that of a model including the polygenic score and covariates minus the variance explained by a model including only covariates as predictors of breastfeeding.

## Results

### Descriptive Results

As stated before, our sample is entirely comprised of mothers who reported data about breastfeeding. The mean number of children was 2.52 (*SD* = 1.26, range = 1–12). Three out of four (75.2%) women breastfed for at least 1 month for each one of their births. The mean duration of breastfeeding was 5.31 months (*SD* = 4.69, range: 0–48), with slight changes according to child order (Table 1). Table 2 presents the correlation matrix for the duration of breastfeeding for the first five children. Breastfeeding reports showed a high test–retest reliability ( $r = 0.96, p < .001$ ) and internal consistency among breastfeeding durations (Cronbach’s  $\alpha = 0.98$ ; ICC = 0.85).

**TABLE 2**  
Correlations (N) for Breastfeeding Duration (Months) in the First Five Children

Child order	1st child	2nd child	3rd child	4th child	5th child
1st child	1 (3,295)				
2nd child	0.67* (2,508)	1 (2,568)			
3rd child	0.51* (1,317)	0.71* (1,336)	1 (1,362)		
4th child	0.47* (525)	0.68* (531)	0.77* (531)	1 (547)	
5th child	0.49* (199)	0.61* (205)	0.68* (203)	0.82* (206)	1 (209)

Note: Spearman’s rho (N). \* $p < .01$ .

**TABLE 3**  
Model-Fitting Results for Univariate Models for Breastfeeding Mean Duration and Proportions of Variance Explained By Additive Genetic Influences (A), Common Environment (C) and Unique Environment (E)

Model	Parameter estimates (CI = 95%)			Goodness-of-fit index					
	A	C	E	-2LL	df	AIC	$\Delta\chi^2$	$\Delta df$	p
ACE	0.34 (0.14, 0.55)	0.17 (0, 0.35)	0.48 (0.43, 0.54)	8862.88	3359	2144.88	—	—	—
<b>AE</b>	<b>0.53 (0.47, 0.58)</b>	—	<b>0.47 (0.42, 0.53)</b>	<b>8866.20</b>	<b>3360</b>	<b>2146.20</b>	<b>3.11</b>	<b>1</b>	<b>0.08</b>
CE	—	0.45 (0.40, 0.50)	0.55 (0.50, 0.60)	8874.71	3360	2154.71	11.83	1	0.001
E	—	—	1 (1,1)	9080.60	3361	2358.60	217.72	2	<.001

Note: A: additive genetic factors, AIC: Akaike’s information criterion, BF: breastfeeding mean duration across all births, C: common environmental factors, CI: confidence interval, df: degrees of freedom, E: unique environmental factors, -2LL: twice negative log-likelihood,  $\Delta\chi^2$ : difference in  $\chi^2$  to saturated model,  $\Delta df$ : difference in degrees of freedom to saturated model. Bold type indicates best-fitting model.


### Twin Correlations and Variance Component Model-Fitting

The MZ twin correlation for the mean breastfeeding duration ( $r_{MZ} = 0.52, 95\% \text{ CI } 0.46\text{--}0.57$ ) was larger than for DZ twins ( $r_{DZ} = 0.35, 95\% \text{ CI } 0.25\text{--}0.43$ ), which suggests the presence of genetic influences (see Figure 1). Lower correlations were found for the breastfeeding duration of the first-born child ( $r_{MZ} = 0.41, 95\% \text{ CI } 0.34\text{--}0.47$ ;  $r_{DZ} = 0.27, 95\% \text{ CI } .17\text{--}0.37$ ).


A univariate model was fit to disentangle the sources of variance of the mean breastfeeding duration (see Table 3). Since the DZ twin correlation was more than half the MZ twin correlation, C was estimated instead of D. The best-fitting model was the one that included additive genetic and non-shared environmental sources of variation — shared environmental factors did not account for a significant portion of variation. Additive genetic factors accounted for 53% in breastfeeding (95% CI 47%–58%) and the remaining 47% was due to unique environmental factors (95% CI 42%–53%). The reduced model fit well, as it did not differ significantly from the full ACE model. The model-fitting analysis revealed a similar structure of the underlying variance for the breastfeeding duration in the first-born child, though with lower heritability (0.42 vs. 0.53 for average duration).

### Genome-Wide Association Study

We conducted a GWAS of breastfeeding in 1,521 individuals from 1,073 Australian families. The average age of the genotyped sample was 46.46 years (*SD* = 11.38). Breastfeeding mean duration in this subsample was 5.52 months (*SD* = 4.75). We tested 6.59 M SNPs for association with breastfeeding, correcting for age at the time of survey.

**FIGURE 1**

Scatter plot of twin correlations with 95% confidence intervals (CI) for breastfeeding duration.  
Note. t1: twin 1, t2: twin 2


**FIGURE 2**

(Colour online) Manhattan plot showing the results of the genome-wide association analyses for breastfeeding. Genes at or nearby best SNPs are indicated. The vertical axis shows the  $-\log_{10}$  of the associated  $p$  values and the horizontal axis shows the chromosome numbers divided into 22 autosomes and the X chromosome.

The Manhattan plot of association  $p$  values for 6.59 M SNPs is shown in Figure 2. While no SNP achieved genome-wide significance ( $p < 6.6 \times 10^{-8}$ ), regions of suggestive association signals, with the smallest  $p$  value of  $1.2 \times 10^{-7}$  obtained for a SNP (rs6950451) were observed on

chromosome 7. Suggestive association signals were also detected in chromosomes 2 (SNP rs930421) and 18 (rs9807759), with  $p$  values of  $1.2 \times 10^{-6}$  in both cases.

The quantile-quantile plot (Q-Q plot) of the observed versus expected (under the null-hypothesis of no

**FIGURE 3**

(Colour online) Quantile–quantile plot for breastfeeding mean duration. The horizontal axis shows the  $-\log_{10}$  of expected  $p$  values of association from a 1 degree of freedom chi-square distribution and the vertical axis shows the  $-\log_{10}$  of  $p$  values from the observed chi-square distribution. The colored dots represent the top hit SNPs. Genes at or nearby best SNPs are indicated.

association)  $\log_{10}(p)$  value from the association analysis is presented in Figure 3. The genomic control  $\lambda$  (1.007) was close to 1.0, indicating that there was no evidence for inflation of the test statistics or a bias because of possible population stratification in the results (Bacanu et al., 2000) and that the family-based association model had correctly accounted for relatedness.

The top SNPs and gene regions from GWAS analysis with the strongest association with breastfeeding are shown in Table 4.

Figure 4 shows the locus zoom plots of the unconditioned analyses for the top five SNPs, rs6950451, rs930451, rs9807759, rs17381960, rs7446359, and also for rs498793, since there are genes at or nearby this latter SNP that have

been emphasized for their role in the composition of fatty acids in mothers' milk (Glaser et al., 2011; Standl et al., 2012). Notably, although LD structure surrounding the chromosome 7 locus is rather sparse, rs6950451 was well imputed ( $R^2 = 0.84$ ). Conditional analyses on the top SNP were also performed, and there were no associated SNPs with  $p < .001$ .

#### Gene-Based Analysis

Gene-based test results are shown in Table 5. Although none of the genes reached genome-wide significance, we list the top 20 genes. Two of these (MRAs and OXER1) have already been noted in the top SNP analysis (Table 4), but the other genes listed are all novel.



**TABLE 4**

Top Ten SNPs and Potential Candidates and Their Gene Regions From GWAS Analysis Showing the Strongest Associations With Breastfeeding

SNP	CHR	bp	Alleles	AL1%	B	SE B	p value	At or near gene(s)
rs6950451	7	22,979,808	A/G	77.1	-0.259	0.049	$1.2 \times 10^{-7}$	SNORD93, FAM126A
rs930451	2	42,981,239	A/G	68.4	-0.204	0.042	$1.2 \times 10^{-6}$	OXER1, HAAO, MTA3
rs9807759	18	60,269,356	C/T	84.4	-0.401	0.083	$1.2 \times 10^{-6}$	ZCCHC2
rs17381960	13	50,095,682	G/A	96.0	-0.778	0.164	$2.2 \times 10^{-6}$	PHF11, RCBTB1
rs7446359	5	127,890,720	G/A	98.3	-1.185	0.256	$3.6 \times 10^{-6}$	FBN2
rs7635879	3	193,736,373	T/A	55.2	0.176	0.039	$5.8 \times 10^{-6}$	DPPA2P3, LOC647323
rs7699884	4	181,295,238	T/C	60.2	0.179	0.040	$7.7 \times 10^{-6}$	(No Gene in $\pm 400$ kb)
rs2277212	10	11,299,735	T/A	74.7	-0.298	0.067	$8.7 \times 10^{-6}$	CELF2, CELF-AS2
rs218271	4	55,411,591	C/G	79.4	-0.238	0.054	$8.9 \times 10^{-6}$	KIT
rs4724103	7	42,339,894	C/T	55.3	-0.167	0.038	$9.8 \times 10^{-6}$	GLI3
rs498793	11	61,624,705	C/T	61.3	-0.185	0.043	$1.7 \times 10^{-5}$	FADS2, FADS3, FADS1

**TABLE 5**

Genes ( $p < 5 \times 10^{-3}$ ) From VEGAS Gene-Based Analysis Showing the Strongest Associations With Breastfeeding

Gene	Chr	Start position	End position	N SNPs	p value	Best SNP	SNP p value
MRAS	3	138,066,489	138,124,377	100	$1.1 \times 10^{-5}$	rs4678408	$1.8 \times 10^{-5}$
EED	11	85,955,805	85,989,785	129	$4.1 \times 10^{-5}$	rs7951030	$2.0 \times 10^{-5}$
PAPD4	5	78,908,242	78,982,471	135	$1.9 \times 10^{-4}$	rs4704567	$5.8 \times 10^{-5}$
DMAP1	1	44,679,124	44,686,351	43	$3.2 \times 10^{-4}$	rs1291169	$1.9 \times 10^{-4}$
INPP5B	1	38,326,368	38,412,729	69	$3.2 \times 10^{-4}$	rs12120737	$2.8 \times 10^{-4}$
SEL1L2	20	13,830,049	13,971,262	244	$4.1 \times 10^{-4}$	rs6042425	$2.1 \times 10^{-5}$
PAPOLB	7	4,897,368	4,901,625	68	$6.5 \times 10^{-4}$	rs6966725	$3.7 \times 10^{-4}$
TPRG1	3	188,889,762	189,041,271	303	$7.0 \times 10^{-4}$	rs1562758	$7.9 \times 10^{-5}$
MTF1	1	38,275,238	38,325,292	53	$7.0 \times 10^{-4}$	rs4329476	$4.1 \times 10^{-4}$
CMYA5	5	78,985,658	79,096,049	245	$1.1 \times 10^{-3}$	rs6859704	$5.8 \times 10^{-5}$
OXER1	2	42,989,638	42,991,401	121	$1.2 \times 10^{-3}$	rs930421	$1.2 \times 10^{-6}$
ARID3B	15	74,833,547	74,890,472	70	$1.3 \times 10^{-3}$	rs11072492	$1.5 \times 10^{-4}$
MYCT1	6	153,019,029	153,045,715	193	$1.7 \times 10^{-3}$	rs6934838	$1.2 \times 10^{-4}$
C20orf7	20	13,765,671	13,799,067	149	$2.0 \times 10^{-3}$	rs6033833	$1.2 \times 10^{-4}$
CHRNA4	15	78,916,635	78,933,587	95	$2.3 \times 10^{-3}$	rs8043123	$6.2 \times 10^{-5}$
ATP9B	18	76,829,396	77,138,282	358	$2.9 \times 10^{-3}$	rs9954562	$1.0 \times 10^{-4}$
MMP7	11	102,391,238	102,401,478	197	$3.4 \times 10^{-3}$	rs7951520	$3.5 \times 10^{-5}$
CPNE2	16	57,126,454	57,181,878	137	$3.7 \times 10^{-3}$	rs2216758	$7.8 \times 10^{-5}$
ADAMT5	15	79,051,544	79,103,773	63	$4.9 \times 10^{-3}$	rs12899940	$3.2 \times 10^{-5}$

### Polygenic Study

Next, we tested an association between the polygenic score for breast size and we found no significant associations between a polygenic score of genetic predisposition to breast size and breastfeeding among the 66 SNPs with  $p < 10^{-4}$  available ( $\beta = 0.011$ ,  $r^2 = 0.003$ ,  $p = 0.67$ ).


### Discussion

In this investigation, we used a broad behavioral genetics approach to analyze individual differences in breastfeeding. The first objective was to explore the proportion of genetic and environmental sources in the variation of breastfeeding. Correlations were higher for MZ twins than for DZ twins, suggesting the presence of genetic factors implicated in this behavior. The results of the twin analysis showed that the best-fitting model was the one including additive genetics effects and unique environmental factors, explaining 53% (CI 95%: 0.47–0.58) and 47% (CI 95%: 0.42–0.53) of the variance in breastfeeding behavior, respectively. The results are also compatible with the full model, in which 17% (CI 95%: 0–0.35) of variance was accounted for by

common environmental factors. The values obtained here are very similar to those found recently in a Spanish sample (Colodro-Conde et al., 2013), where 54% of the variance of breastfeeding duration of all the offspring was due to additive genetic factors and the remaining 46% to unique environmental factors. Despite the samples coming from two distinct cultural backgrounds (Australia vs. Spain) and measures being different (quantitative vs. ordinal), the similar results obtained confirm that genetic factors are an important source of variation between women's breastfeeding behavior. While we have not been able to detect any significant effect of shared environmental factors, this may be because of a lack of power due to sample size and our results, which are compatible with a  $C^2$  as high as 35%.

Confirmation that genetic factors influence breastfeeding behavior justified our exploring the genetic variants underlying this trait by GWAS, as is discussed below.

The present study is, to our knowledge, the first to perform a GWAS on breastfeeding behaviors. No genome-wide significant SNPs were identified. However, although not significant, we found a promising genetic region on chromosome 7, around rs6950451. Likewise, the VEGAS



**FIGURE 4**

(Colour online) Regional association plots for breastfeeding. (A), showing the top associated SNP rs6950451 ( $p = 1.2 \times 10^{-7}$ ) on chromosome 7; (B-F), showing the possible associated regions; on chromosomes 2, 5, 11, 13 and 18.

gene-based analysis did not find any significant results. In addition, we detected suggestive associations of some genetic regions with plausible links to breastfeeding. Among them, the oxoeicosanoid (OXE) receptor 1 gene ( $p = 1.2 \times 10^{-3}$ ) in chromosome 2 is a receptor for eicosanoids and polyunsaturated fatty acids (Rebhan et al., 1997), which are present in breast milk (Koletzko et al., 2008).

It is well known that breast milk provides a unique supply of long chain polyunsaturated fatty acids and that their

synthesis is controlled by key enzymes encoded by the FADS gene cluster (Marquardt et al., 2000; Martin et al., 2011). In this work, we found a suggestive association of the FADS1 ( $p = 2.3 \times 10^{-2}$ ), FADS2 ( $p = 3.9 \times 10^{-2}$ ) and FADS3 ( $p = 4 \times 10^{-2}$ ) genes, on chromosome 11 (Figure 4, panel F), with breastfeeding. Several studies have shown strong associations between the FADS gene cluster and fatty acid levels in breast milk (Glaser et al., 2011; Standl et al., 2012). Maternal genetic variants in the FADS gene cluster have previously

been associated not only with higher colostrum levels of long chain polyunsaturated fatty acids but also with higher cognitive scores in their children (Morales et al., 2011).

Accordingly, changes in these gene regions may have an effect on breastfeeding duration. As no other study has explored the molecular genetic basis of breastfeeding, we cannot yet replicate our findings. The fact that we did not find genome-wide significant results is to be expected, considering the low power.

This article has some limitations that need to be taken into consideration. First, measures of breastfeeding were based on self-report and much of the data used was reported a long time after the breastfeeding took place. A larger sample is also required for the GWAS as we expect that breastfeeding is influenced by a large number of SNPs with very small effect size. Ideally, we seek to conduct a meta-analysis combining different samples.

In summary, we have confirmed that genetic factors are significant in the explanation of breastfeeding behavior variability. However, we could neither identify any breastfeeding-related SNP reaching conventional levels of genome-wide significance, nor any overlap with an individual's genetic predisposition towards breast size. This study provides replication of a previous one that explored the sources of variance of breastfeeding in a Spanish sample and, to our knowledge, is the first one to conduct a preliminary GWAS on breastfeeding.

## Acknowledgments

We are grateful to the twins for their participation and to the data collectors and data managers Anthony Conciatore, Richard Parker and Scott Gordon. We also thank Liz Ritschel, Jana Strohmaier, Brendan Zietsch, Ada Johansson, Patrick Jern, and Miguel Renteria for their comments on earlier drafts of this paper. Genome-wide genotyping in the Australian sample was funded by NIH Grants AA013320, AA013321, AA013326, AA011998 and AA017688. LCC was supported for a research visit at Genetic Epidemiology Laboratory, QIMR, and for a postdoctoral fellowship by Seneca Foundation — Regional Agency for Science and Technology, Murcia, Spain (12431/FPI/09 and 19151/PD/13).

## References

Almqvist-Tangen, G., Bergman, S., Dahlgren, J., Roswall, J., & Alm, B. (2012). Factors associated with discontinuation of breastfeeding before 1 month of age. *Acta Paediatrica*, *101*, 55–60.

Australian Bureau of Statistics. (2013). *Australian Health Survey: Health Service Usage and Health Related Actions, 2011–12*. Retrieved from <http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/6664B939E49FD9C1CA257B39000F2E4B>

Bacanu, S.-A., Devlin, B., & Roeder, K. (2000). The power of genomic control. *The American Journal of Human Genetics*, *66*, 1933–1944.

Balen, A. H., Conway, G. S., Kaltsas, G., Techatrasak, K., Manning, P. J., West, C., . . . Jacobs, H. S. (1995). Andrology: Polycystic ovary syndrome: The spectrum of the disorder in 1741 patients. *Human Reproduction*, *10*, 2107–2111.

Barber, T., McCarthy, M., Wass, J., & Franks, S. (2006). Obesity and polycystic ovary syndrome. *Clinical Endocrinology*, *65*, 137–145.

Brown, A. (2014). Maternal trait personality and breastfeeding duration: The importance of confidence and social support. *Journal of Advanced Nursing*, *70*, 587–598.

Brown, A., Raynor, P., & Lee, M. (2011). Healthcare professionals' and mothers' perceptions of factors that influence decisions to breastfeed or formula feed infants: A comparative study. *Journal of Advanced Nursing*, *67*, 1993–2003.

Calnen, G. (2010). A strategy for the future. *Breastfeeding Medicine*, *5*, 95–95.

Centers for Disease Control and Prevention. (2014). *Breastfeeding among US children born 2001–2011, CDC National Immunization Survey*. Retrieved from <http://www.cdc.gov/breastfeeding/data/nis/data/>

Chen, W.-M., & Abecasis, G. R. (2007). Family-based association tests for genomewide association scans. *The American Journal of Human Genetics*, *81*, 913–926.

Colodro-Conde, L., Limiñana Gras, R. M., & Ordoñana, J.R. (in press (a)). Estilos de Personalidad y Adaptación Psicológica en la toma de decisiones sobre Lactancia en mujeres primíparas [Personality Styles and Psychological Adjustment when making decisions about Breastfeeding in primiparous women]. *Revista Iberoamericana de Diagnóstico y Evaluación Psicológica*.

Colodro-Conde, L., Limiñana Gras, R. M., Sánchez-López, M.P., & Ordoñana, J.R. (in press (b)). Gender, Health and Initiation of Breastfeeding. *Women & Health*.

Colodro-Conde, L., Sánchez-Romera, J. F., & Ordoñana, J. R. (2013). Heritability of initiation and duration of breastfeeding behavior. *Twin Research and Human Genetics*, *16*, 575–580.

Colodro-Conde, L., Sánchez-Romera, J. F., Tornero-Gómez, M. J., Pérez-Riquelme, F., Polo-Tomás, M., & Ordoñana, J. R. (2011). Relationship between level of education and breastfeeding duration depends on social context breastfeeding trends over a 40-year period in Spain. *Journal of Human Lactation*, *27*, 272–278.

De Jager, M., Hartley, K., Terrazas, J., & Merrill, J. (2012). Barriers to breastfeeding – A global survey on why women start and stop breastfeeding. *European Obstetrics & Gynaecology*, *7*, 25–30.

Dudbridge, F., & Gusnanto, A. (2008). Estimation of significance thresholds for genomewide association scans. *Genetic Epidemiology*, *32*, 227–234.

Eriksson, N., Benton, G. M., Do, C. B., Kiefer, A. K., Mountain, J. L., Hinds, D. A., . . . Tung, J. Y. (2012). Genetic variants associated with breast size also influence breast cancer risk. *BMC Medical Genetics*, *13*, 53.

- Glaser, C., Lattka, E., Rzehak, P., Steer, C., & Koletzko, B. (2011). Genetic variation in polyunsaturated fatty acid metabolism and its potential relevance for human development and health. *Maternal & Child Nutrition*, 7, 27–40.
- Heath, A. C., Bucholz, K., Madden, P., Dinwiddie, S., Slutske, W., Bierut, L., . . . Martin, N. (1997). Genetic and environmental contributions to alcohol dependence risk in a national twin sample: Consistency of findings in women and men. *Psychological Medicine*, 27, 1381–1396.
- Heinig, M. J., & Dewey, K. G. (1997). Health effects of breast feeding for mothers: A critical review. *Nutrition Research Reviews*, 10, 35–56.
- Holman, D. J., & Grimes, M. A. (2003). Patterns for the initiation of breastfeeding in humans. *American Journal of Human Biology*, 15, 765–780.
- Ip, S., Chung, M., Raman, G., Chew, P., Magula, N., DeVine, D., . . . Lau, J. (2009). *Breastfeeding and maternal and infant health outcomes in developed countries* (Evidence Report/Technology Assessment No. 153). Rockville, MD: Agency for Healthcare Research and Quality.
- Jevitt, C., Hernandez, I., & Groër, M. (2007). Lactation complicated by overweight and obesity: Supporting the mother and newborn. *Journal of Midwifery & Women's Health*, 52, 606–613.
- Jonas, W., Mileva-Seitz, V., Girard, A., Bisceglia, R., Kennedy, J., Sokolowski, M., . . . Steiner, M. (2013). Genetic variation in oxytocin rs2740210 and early adversity associated with postpartum depression and breastfeeding duration. *Genes, Brain and Behavior*, 12, 681–694.
- Katz, K. A., Nilsson, I., & Rasmussen, K. M. (2010). Danish health care providers' perception of breastfeeding difficulty experienced by women who are obese, have large breasts, or both. *Journal of Human Lactation*, 26, 138–147.
- Knopik, V. S., Heath, A. C., Madden, P. A., Bucholz, K. K., Slutske, W. S., Nelson, E. C., . . . Martin, N. G. (2004). Genetic effects on alcohol dependence risk: Re-evaluating the importance of psychiatric and other heritable risk factors. *Psychological Medicine*, 34, 1519–1530.
- Koletzko, B., Lien, E., Agostoni, C., Böhles, H., Campoy, C., Cetin, I., . . . Forsyth, S. (2008). The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: Review of current knowledge and consensus recommendations. *Journal of Perinatal Medicine*, 36, 5–14.
- Li, J., Kendall, G., Henderson, S., Downie, J., Landsborough, L., & Oddy, W. (2008). Maternal psychosocial well-being in pregnancy and breastfeeding duration. *Acta Paediatrica*, 97, 221–225.
- Liu, J. Z., Mcrae, A. F., Nyholt, D. R., Medland, S. E., Wray, N. R., Brown, K. M., . . . Martin, N. G. (2010). A versatile gene-based test for genome-wide association studies. *The American Journal of Human Genetics*, 87, 139–145.
- Marquardt, A., Stöhr, H., White, K., & Weber, B. H. (2000). cDNA cloning, genomic structure, and chromosomal localization of three members of the human fatty acid desaturase family. *Genomics*, 66, 175–183.
- Martin, N. W., Benyamin, B., Hansell, N. K., Montgomery, G. W., Martin, N. G., Wright, M. J., . . . Bates, T. C. (2011). Cognitive function in adolescence: Testing for interactions between breast-feeding and FADS2 polymorphisms. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50, 55–62.
- McFadden, A., & Toole, G. (2006). Exploring women's views of breastfeeding: A focus group study within an area with high levels of socio-economic deprivation. *Maternal & Child Nutrition*, 2, 156–168.
- Medland, S. E., Nyholt, D. R., Painter, J. N., McEvoy, B. P., McRae, A. F., Zhu, G., . . . Henders, A. K. (2009). Common variants in the trichohyalin gene are associated with straight hair in Europeans. *The American Journal of Human Genetics*, 85, 750–755.
- Ministerio de Sanidad, Servicios Sociales e Igualdad (MSSSI). Encuesta Nacional de Salud de España [Spanish National Health Survey]. (2012). Retrieved from <http://www.msssi.gob.es/en/estadEstudios/estadisticas/sisInfSanSNS/home.htm>.
- Morales, E., Bustamante, M., Gonzalez, J. R., Guxens, M., Torrent, M., Mendez, M., . . . Vrijheid, M. (2011). Genetic variants of the FADS gene cluster and ELOVL gene family, colostrums LC-PUFA levels, breastfeeding, and child cognition. *PLoS One*, 6, e17181.
- Neale, M., Boker, S., Xie, G., & Maes, H. (2006). *Mx: Statistical modeling* (7th ed.). Richmond, VA: Virginia Institute for Psychiatric and Behavioral Genetics. Department of Psychiatry.
- Neale, M. C., & Cardon, L. R. (1992). *Methodology for genetic studies of twins and families*. Dordrecht, The Netherlands: Kluwer Academic Publishers.
- Ordoñana, J. R., Perez-Riquelme, F., Gonzalez-Javier, F., Carrillo, E., Gomez-Amor, J., & Martinez-Selva, J. M. (2006). An initiative in Spain for the study of women's health: The Murcia twin registry. *Twin Research and Human Genetics*, 9, 865–867.
- Ordoñana, J. R., Rebollo-Mesa, I., Carrillo, E., Colodro-Conde, L., García-Palomo, F. J., González-Javier, F., . . . Pérez-Riquelme, F. (2013). The Murcia twin registry: A population-based registry of adult multiples in Spain. *Twin Research and Human Genetics*, 16, 302–306.
- Posthuma, D., Beem, A. L., de Geus, E. J. C., van Baal, G. C. M., von Hjelmberg, J. B., Iachine, I., . . . Boomsma, D. I. (2003). Theory and practice in quantitative genetics. *Twin Research and Human Genetics*, 6, 361–376.
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., . . . Daly, M. J. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. *The American Journal of Human Genetics*, 81, 559–575.
- Rebhan, M., Chalifa-Caspi, V., Prilusky, J., & Lancet, D. (1997). *GeneCards: Encyclopedia for genes, proteins and diseases*. Retrieved from <http://www.genecards.org/>.
- Sakalidis, V. S., Williams, T. M., Garbin, C. P., Hepworth, A. R., Hartmann, P. E., Paech, M. J., . . . Geddes, D. T. (2013). Ultrasound imaging of infant sucking dynamics during the establishment of lactation. *Journal of Human Lactation*, 29, 205–213.
- Schön, R. A., & Silvé, M. (2007). Natural parenting: Back to basics in infant care. *Evolutionary Psychology*, 5, 102–183.

- Sellen, D. W. (2007). Evolution of infant and young child feeding: Implications for contemporary public health. *Annual Review of Nutrition*, 27, 123–148.
- Spencer, J. P. (2008). Management of mastitis in breastfeeding women. *American Family Physician*, 78, 727–731.
- SPSS. (2010). *IBM SPSS Statistics Base 19*. Chicago, IL: SPSS Inc.
- Standl, M., Sausenthaler, S., Lattka, E., Koletzko, S., Bauer, C., Wichmann, H., . . . Schaaf, B. (2012). FADS gene cluster modulates the effect of breastfeeding on asthma. Results from the GINIplus and LISApplus studies. *Allergy*, 67, 83–90.
- Swanson, V., & Power, K. G. (2005). Initiation and continuation of breastfeeding: Theory of planned behaviour. *Journal of Advanced Nursing*, 50, 272–282.
- Tharner, A., Luijk, M. P., Raat, H., IJzendoorn, M. H., Bakermans-Kranenburg, M. J., Moll, H. A., . . . Tiemeier, H. (2012). Breastfeeding and its relation to maternal sensitivity and infant attachment. *Journal of Developmental & Behavioral Pediatrics*, 33, 396–404.
- Thulier, D., & Mercer, J. (2009). Variables associated with breastfeeding duration. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*, 38, 259–268.
- Verweij, K. J., Zietsch, B. P., Medland, S. E., Gordon, S. D., Benjamin, B., Nyholt, D. R., . . . Madden, P. A. (2010). A genome-wide association study of Cloninger's temperament scales: Implications for the evolutionary genetics of personality. *Biological Psychology*, 85, 306–317.
- Wade, T. D., Zhu, G., & Martin, N. G. (2010). Body mass index and breast size in women: Same or different genes? *Twin Research and Human Genetics*, 13, 450–454.
- Wagner, C. L., Wagner, M. T., Ebeling, M., Chatman, K. G., Cohen, M., & Hulseley, T. C. (2006). The role of personality and other factors in a mother's decision to initiate breastfeeding. *Journal of Human Lactation*, 22, 16–26.
- Wojcicki, J. M. (2011). Maternal prepregnancy body mass index and initiation and duration of breastfeeding: A review of the literature. *Journal of Women's Health*, 20, 341–347.
- World Health Organization. (2013). Global health observatory data repository [database].
- World Health Organization, K. G. (1997). *Growth patterns of breastfed infants and the current status of growth charts for infants*. (No. Joint WHO/UNICEF Meeting on Infant and Young Child Feeding: Statement and Recommendations). Geneva, Switzerland: World Health Organization.
- World Health Organization, & United Nations Children's Fund. (2003). *Global strategy for infant and young child feeding*. Geneva, Switzerland: World Health Organization.

---

**Appendix**

**Births**

---

1. With each pregnancy resulting in birth have you had any of the following?

---

	Birth							
	1	2	3	4	5	6	7	8+
PLEASE TICK IF ANSWER IS YES								
1. High blood pressure, toxemia								
2. Premature baby (over 2 weeks early)								
3. Epidural anesthetic (block)								
4. Induced labor								
5. Stitches (episiotomy or tear)								
6. Stillborn child								
7. Forceps delivery								
8. Cesarean section								
2. What time was the baby born? (e.g., 2 pm)								
3. How many hours was the labor? (e.g., 11 hours)								
4. What was the birth weight of the baby?								
5. Was each labor painful/difficult? (1, 2 or 3)								
(1) Extremely (2) Quite (3) Not really								
6. Did you feel depressed after the birth of any of your children? TICK IF YES								
IF YES:								
How many weeks did this go on for?								
Did you need to seek help for the depression?								
7. How many months did you breastfeed for?								
8. How many months old was the child when you introduced formula or artificial feeds?								
9. How many months old was the child when you introduced solids?								

---