

2015

## Evaluation of $\alpha,\beta$ -unsaturated ketones as antileishmanial agents

Miguel A. Vasquez  
*University of Texas at El Paso*

Eva Iniguez  
*University of Texas at El Paso*

Umashankar Das  
*University of Saskatchewan*

Stephen M. Beverley  
*Washington University School of Medicine in St. Louis*

Linda J. Herrera  
*University of Texas at El Paso*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Vasquez, Miguel A.; Iniguez, Eva; Das, Umashankar; Beverley, Stephen M.; Herrera, Linda J.; Dimmock, Jonathan R.; and Maldonado, Rosa A., "Evaluation of  $\alpha,\beta$ -unsaturated ketones as antileishmanial agents." *Antimicrobial agents and chemotherapy*. 59, 3598-3601. (2015).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/3929](https://digitalcommons.wustl.edu/open_access_pubs/3929)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

---

**Authors**

Miguel A. Vasquez, Eva Iniguez, Umashankar Das, Stephen M. Beverley, Linda J. Herrera, Jonathan R. Dimmock, and Rosa A. Maldonado

# Evaluation of $\alpha,\beta$ -Unsaturated Ketones as Antileishmanial Agents

Miguel A. Vasquez,<sup>a</sup> Eva Iniguez,<sup>a</sup> Umashankar Das,<sup>b</sup> Stephen M. Beverley,<sup>c</sup> Linda J. Herrera,<sup>a</sup> Jonathan R. Dimmock,<sup>b</sup>  
Rosa A. Maldonado<sup>a</sup>

Border Biomedical Research Center, Department of Biological Sciences, The University of Texas at El Paso, El Paso, Texas, USA<sup>a</sup>; College of Pharmacy and Nutrition, University of Saskatchewan, Saskatoon, Saskatchewan, Canada<sup>b</sup>; Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA<sup>c</sup>

**In this study, we assessed the antileishmanial activity of 126  $\alpha,\beta$ -unsaturated ketones. The compounds NC901, NC884, and NC2459 showed high leishmanicidal activity for both the extracellular (50% effective concentration [EC<sub>50</sub>], 456 nM, 1,122 nM, and 20 nM, respectively) and intracellular (EC<sub>50</sub>, 1,870 nM, 937 nM, and 625 nM, respectively) forms of *Leishmania major* propagated in macrophages, with little or no toxicity to mammalian cells. Bioluminescent imaging of parasite replication showed that all three compounds reduced the parasite burden in the murine model, with no apparent toxicity.**

*Leishmania major* is the causative agent of cutaneous leishmaniasis (CL) disease (1). Worldwide, the incidence of CL is estimated to be 0.7 to 1.2 million new cases per year (2). Other than the drug miltefosine, which was approved by the FDA in 2014, most of the currently used antileishmanial drugs were developed in the 1940s (3, 4). However, drug-resistant strains of the parasite have emerged. Furthermore, toxicity to the host and the high costs of these drugs limit their wider application and use (4, 5). Therefore, the burden of leishmaniasis and limited effective treatments clearly point to the need for new drugs.

In view of the current interest in examining different antineoplastic agents for their antiprotozoal properties (6–8), we assessed the antileishmanial activity of  $\alpha,\beta$ -unsaturated ketones (enones) provided by J. R. Dimmock from the University of Saskatchewan in Canada (1, 9). Enones react preferentially with cellular thiols, in contrast to amino or hydroxyl functional groups present in protein and DNA (10, 11); therefore, interactions with nucleic acids, which can lead to adverse genotoxic effects, should be absent in enones (12). Since thiol-dependent metabolism is the main detoxifying mechanism in trypanosomatids, we hypothesized that enones are attractive candidates for examination as potential antileishmanial agents.

Hence, in this study, we screened the enone library for antiparasitic and cytotoxic activity. The compounds were dissolved in dimethyl sulfoxide (DMSO) and tested at concentrations ranging from 500  $\mu$ M to 1 nM (1, 9). The parasites tested were a firefly luciferase-expressing line of *L. major* promastigotes described previously (Lmj-FV1-LUC-TK [*L. major* strain Friedlin {MHOM/JL/80/Friedlin}], clone V1) and cultured as previously described (13, 14). The 126 enones were incubated with 10<sup>6</sup> *L. major* promastigotes or rhesus monkey kidney epithelial cells (LLC-MK<sub>2</sub>)/well for 96 h and were analyzed for toxicity and parasite survival as measured by luciferase activity with the substrate 5'-fluoroluciferin (ONE-Glo luciferase assay system; Promega) using a luminometer (Luminoskan; Thermo Scientific). Sixty-four compounds inhibited the survival of *L. major* promastigotes at >75%. At the same concentrations tested, 20 out of 126 screened compounds displayed minimal to no toxicity (>75% survival) against LLC-MK<sub>2</sub> cells. Only six compounds met the criteria for both antiparasitic activity (<25%) and low cytotoxicity to LLC-MK<sub>2</sub> cells (>75%) (see Fig. S1 in the supplemental material). These six compounds were further tested in the following mammalian cell lines: Hs27

human fibroblasts, RAW 264.7 murine macrophages (American Type Culture Collection [ATCC], Manassas, VA), and peritoneal BALB/c mice macrophages, obtained as described previously (15). Only NC901, NC884, and NC2459 showed effective activity against *L. major*. NC2459 showed a 100-fold difference between the 50% effective concentration (EC<sub>50</sub>) of the parasite and that for all three mammalian cell lines (Table 1).

The *in vitro* infectivity experiments were carried out to determine the activity of NC901, NC884, and NC2459 against *L. major* intracellular amastigotes. Peritoneal macrophages isolated from BALB/c mice were infected with *L. major* metacyclic promastigotes for 24 h, followed by treatment with the NC lead compounds for an additional 48 h (Table 1). The antileishmanial activity exhibited by the compounds was evaluated using BD Pathway Bio-imager high-content imaging assay (HCIA) analysis (16–18). In comparison to the 1% DMSO control, all three compounds showed a significant decrease in the percentage of infected cells. NC2459 was the most effective at a concentration of 1.25  $\mu$ M (Table 1).

Additionally, the activity of the compounds against *L. major* was investigated in a murine model of CL. The first set of experiments (Fig. 1A) consisted of 12 female BALB/c mice organized in four groups of three, each infected with 10<sup>5</sup> *L. major* metacyclic promastigotes (Table 2). As shown in Fig. 1A, at 46 days postinfection, all experimental groups exhibited a lesion size smaller

Received 14 August 2014 Returned for modification 22 October 2014

Accepted 15 March 2015

Accepted manuscript posted online 23 March 2015

Citation Vasquez MA, Iniguez E, Das U, Beverley SM, Herrera LJ, Dimmock JR, Maldonado RA. 2015. Evaluation of  $\alpha,\beta$ -unsaturated ketones as antileishmanial agents. *Antimicrob Agents Chemother* 59:3598–3601. doi:10.1128/AAC.04056-14.




Address correspondence to Rosa A. Maldonado, ramaldonado@utep.edu.

M.A.V. and E.I. contributed equally to this article.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.04056-14>.

Copyright © 2015, American Society for Microbiology. All Rights Reserved. doi:10.1128/AAC.04056-14

TABLE 1 Summary of EC<sub>50</sub> and IC<sub>50</sub> for NC901, NC884, and NC2459 tested against *L. major* and mammalian cells<sup>a</sup>

Compound	Mammalian cell EC <sub>50</sub> (μM) ± SD for:			<i>L. major</i> cells (μM) (TI)	
	Peritoneal murine macrophages	LLC-MK <sub>2</sub>	Hs27 fibroblasts	Promastigote EC <sub>50</sub>	Intracellular amastigote IC <sub>50</sub>
 NC901 2HCl	7.03 ± 0.42	8.02 ± 0.25	16.0 ± 0.12	0.45 (15.62)	1.87 ± 0.20 (3.76)
 NC884 2HCl	7.67 ± 0.31	15.1 ± 0.33	16.3 ± 0.17	1.12 (6.55)	0.937 ± 0.13 (8.18)
 NC2459	5.45 ± 0.36	2.0 ± 0.093	10.01 ± 0.09	0.020 (272.5)	0.625 ± 0.11 (8.72)

<sup>a</sup> EC<sub>50</sub>, half-maximal effective concentration calculated with 95% confidence; IC<sub>50</sub>, 50% inhibitory concentration; TI, therapeutic index, calculated as (EC<sub>50</sub> peritoneal murine macrophages)/(EC<sub>50</sub>/IC<sub>50</sub> parasites). The EC<sub>50</sub> was obtained as the exponent of the negative ratio of the  $\gamma$ -intercept and the slope of the fitted regression line (version 9.2; SAS Software). *P* value, <0.0001 for all concentrations. The Z-factor calculated for the HCIA is in the range of 0.5 to 0.91, indicating that the quality of the assay is excellent.


than or equal to that of the group treated with amphotericin B. Further support was obtained by determining the relative amount of luminescence emitted from the luciferase-expressing parasites in the infected footpad at the endpoint of the study, namely, 46 days postinfection (Fig. 1C). All mice in the NC2459 group did not develop measurable lesions until 2 to 3 weeks after the last treatment (total of 6 to 7 weeks postinfection). Furthermore, one out of the three mice did not develop a lesion up to 12 weeks postinfection.

To determine the relative toxicity of the compounds, the weights of the mice were measured twice a week (data not shown). There was no significant weight loss for any of the groups except for the mice receiving amphotericin B. Additionally, no mice died as a result of the toxicity of the compounds.

Hyperinfection with *L. major* was tested. Twenty-six female BALB/c mice organized in four groups of five and one group of six mice were infected with 10<sup>6</sup> *L. major* metacyclic promastigotes (Fig. 1B). All experimental groups were given higher doses than the groups receiving 10<sup>5</sup> *L. major* metacyclic promastigotes (Table 2). At 4 weeks postinfection, the lesion size in all three NC compounds significantly reduced in comparison to that of the diluent control group (*P* ≤ 0.05). However, at 29 days postinfection, the mice in the diluent control group had to be euthanized to avoid physical distress caused by a large lesion; all other groups were monitored for an additional 4 days. For this reason, the luminescence from the infected footpads for each group was analyzed at 29 days postinfection and not at the endpoint of the experiment (Fig. 1D). After 1 week of treatment with NC2459, three of the mice

from this group were given double the daily dose of drug (8 mg/kg of body weight/day), while the other three remained on the same daily dose (4 mg/kg/day). In addition, the three mice given the higher dose were treated with four additional doses. Those given NC2459 (extended treatment) at 33 days postinfection showed a smaller lesion than that of all the other groups (*P* ≤ 0.05). The increase in dosage for NC2459 (extended treatment) did not result in any additional weight loss or toxic side effects in comparison to those of the control group. The procedures were performed to minimize distress and pain for the animals according to NIH guidance and the animal protocol approved by The University of Texas at El Paso (UTEP) Institutional Animal Care and Use Committee (IACUC).

Our findings demonstrate the ability of these compounds to reduce *L. major* replication *in vivo* without any obvious toxic side effects. NC2459 evidently exhibited a much smaller amount of swelling than that in all other groups by decreasing the footpad lesion by 99.99% (Fig. 1A) in comparison with that of the diluent-treated group. In summary, the data obtained in this study suggest that our lead compound, NC2459, is an excellent antiparasitic agent and might be useful as a potential alternative to the current clinical drugs used to treat leishmaniasis. Clearly, NC2459 is an important lead molecule, which is structurally divergent from contemporary medications employed to treat parasitic diseases. It is a prototypic molecule, and further molecular modifications are required, followed by the relevant bioassays, to increase potency and especially to increase the differential in toxicities between parasites and normal cells.



**FIG 1** *In vivo* activity of compounds NC901, NC884, and NC2459 in the murine model of CL. (A and B) Footpad size of BALB/c mice infected with  $10^5$  and  $10^6$  *L. major* metacyclic promastigotes, respectively. The Rx1 line refers to the time period in which treatment was administered daily, and the Rx2 line indicates that the daily dosage of NC2459 was doubled for that time period. (C) Luminescence of BALB/c mice footpads at 46 days postinfection with  $10^5$  metacyclic promastigotes. \*,  $P < 0.05$ . (D) Luminescence of BALB/c mice footpads at 29 days postinfection with  $10^6$  metacyclic promastigotes. C, group treated with vehicle control; CD, group treated with amphotericin B. A statistical analysis of the net intensity (luminescence) of both the  $10^5$  and  $10^6$  experiments was carried out using the two-sided unpaired *t* test. \*,  $P \leq 0.05$ ; \*\*,  $P \leq 0.01$ .

**TABLE 2** Treatment regimens of the *in vivo* experiments

Compound	Treatment (mg/kg/day) after infection with:	
	$10^5$ promastigotes <sup>a</sup>	$10^6$ promastigotes <sup>b</sup>
DMSO (100 $\mu$ l)		
AmpB <sup>c</sup>	8	8
NC901	2	4
NC884	4	8
NC2459	1	4 (3 mice/14 days)/ 8 (3 mice/days 8–17)

<sup>a</sup> The mice infected with  $10^5$  *L. major* metacyclic promastigotes were treated for 14 consecutive days at the stated dosages.

<sup>b</sup> The mice infected with  $10^6$  *L. major* metacyclic promastigotes (hyperinfection experiment) were split into two groups of three mice after 7 days of treatment with NC2459. One group remained on the same treatment, 4 mg/kg/day, for seven more days (14 total days of treatment), and the other group was given double the daily dosage, 8 mg/kg/day, for 10 more days (17 total days of treatment).

<sup>c</sup> AmpB, amphotericin B.

## ACKNOWLEDGMENTS

This study was supported by grant 2S06GM00812-37 from the NIH/MBRS/NIGMS/SCORE program and NIH grant AI29646 to S.M.B. M.A.V. was supported by the RISE (grant 5R25GM069621-06) Graduate Research Program.

We thank the Biomolecule Analysis Core Facility (BACF), High-Throughput Core Facility (HTSCF) (Renato Aguilera, Carolina Lema, and Armando Varela), and the Statistical Consulting Laboratory (SCL) at the Border Biomedical Research Center (BBRC), UTEP, supported by NIH-NIMHD-RCMI grant 2G12MD007592. We thank Kelly Robinson for providing the luciferase-expressing *L. major* parasites. We also thank Stephen M. Beverley and F. Matthew Kuhlman at Washington University in St. Louis for the animal training on the mouse model for leishmaniasis. We thank the Canadian Institutes of Health Research (J.R.D.).

## REFERENCES

1. Das S, Das U, Varela-Ramirez A, Lema C, Aguilera RJ, Balzarini J, De Clercq E, Dimmock SG, Gorecki DK, Dimmock JR. 2011. Bis[3,5-bis(benzylidene)-4-oxo-1-piperidinyl]amides: a novel class of potent cytotoxins. *ChemMedChem* 6:1892–1899. <http://dx.doi.org/10.1002/cmdc.201100199>.

2. Alvar J, Velez ID, Bern C, Herrero M, Desieux P, Cano J, Jannin J, den Boer M, WHO Leishmaniasis Control Team. 2012. Leishmaniasis worldwide and global estimates of its incidence. *PLoS One* 7:e35671. <http://dx.doi.org/10.1371/journal.pone.0035671>.
3. Haldar AK, Sen P, Roy S. 2011. Use of antimony in the treatment of leishmaniasis: current status and future directions. *Mol Biol Int* 2011: 571242. <http://dx.doi.org/10.4601/2011/571242>.
4. Croft SL, Barrett MP, Urbina JA. 2005. Chemotherapy of trypanosomiasis and leishmaniasis. *Trends Parasitol* 21:508–512. <http://dx.doi.org/10.1016/j.pt.2005.08.026>.
5. Desjeux P. 2004. Leishmaniasis: current situation and new perspectives. *Comp Immunol Microbiol Infect Dis* 27:305–318. <http://dx.doi.org/10.1016/j.cimid.2004.03.004>.
6. Nzila A, Okombo J, Becker RP, Chilengi R, Lang T, Niehues T. 2010. Anticancer agents against malaria: time to revisit? *Trends Parasitol* 26: 125–129. <http://dx.doi.org/10.1016/j.pt.2009.12.002>.
7. Steverding D. 2010. The development of drugs for treatment of sleeping sickness: a historical review. *Parasit Vectors* 3:15. <http://dx.doi.org/10.1186/1756-3305-3-15>.
8. Wenzel NI, Chavain N, Wang Y, Friebolin W, Maes L, Pradines B, Lanzer M, Yardley V, Brun R, Herold-Mende C, Biot C, Toth K, Davioud-Charvet E. 2010. Antimalarial versus cytotoxic properties of dual drugs derived from 4-aminoquinolines and Mannich bases: interaction with DNA. *J Med Chem* 53:3214–3226. <http://dx.doi.org/10.1021/jm9018383>.
9. Dimmock JR, Sidhu KK, Chen M, Reid RS, Allen TM, Kao GY, Truitt GA. 1993. Evaluation of some Mannich bases of cycloalkanones and related compounds for cytotoxic activity. *Eur J Med Chem* 28:313–322.
10. Mutus B, Wagner JD, Talpas CJ, Dimmock JR, Phillips OA, Reid RS. 1989. 1-*p*-Chlorophenyl-4,4-dimethyl-5-diethylamino-1-penten-3-one hydrochloride, a sulfhydryl-specific compound which reacts irreversibly with protein thiols but reversibly with small molecular weight thiols. *Anal Biochem* 177:237–243.
11. Baluja G, Municio AM, Vega S. 1964. Reactivity of some  $\alpha,\beta$ -unsaturated ketones towards sulfhydryl compounds and their antifungal activity. *Chem Ind* 1964:2053–2054.
12. Benvenuto JA, Connor TH, Monteith DK, Laidlaw JL, Adams SC, Matney TS, Theiss JC. 1993. Degradation and inactivation of antitumor drugs. *J Pharm Sci* 82:988–991. <http://dx.doi.org/10.1002/jps.2600821003>.
13. Thalhoffer CJ, Graff JW, Love-Homan L, Hickerson SM, Craft N, Beverley SM, Wilson ME. 2010. *In vivo* imaging of transgenic *Leishmania* parasites in a live host. *J Vis Exp* (2010):1980. <http://dx.doi.org/10.3791/1980>.
14. Capul AA, Barron T, Dobson DE, Turco SJ, Beverley SM. 2007. Two functionally divergent UDP-Gal nucleotide sugar transporters participate in phosphoglycan synthesis in *Leishmania major*. *J Biol Chem* 282:14006–14017. <http://dx.doi.org/10.1074/jbc.M610869200>.
15. Capul AA, Hickerson S, Barron T, Turco SJ, Beverley SM. 2007. Comparisons of mutants lacking the Golgi UDP-galactose or GDP-mannose transporters establish that phosphoglycans are important for promastigote but not amastigote virulence in *Leishmania major*. *Infect Immun* 75:4629–4637. <http://dx.doi.org/10.1128/IAI.00735-07>.
16. Nohara LL, Lema C, Bader JO, Aguilera RJ, Almeida IC. 2010. High-content imaging for automated determination of host-cell infection rate by the intracellular parasite *Trypanosoma cruzi*. *Parasitol Int* 59:565–570. <http://dx.doi.org/10.1016/j.parint.2010.07.007>.
17. Martínez A, Carreon T, Iniguez E, Anzellotti A, Sanchez A, Tyan M, Sattler A, Herrera L, Maldonado RA, Sanchez-Delgado RA. 2012. Searching for new chemotherapies for tropical diseases: ruthenium-clotrimazole complexes display high *in vitro* activity against *Leishmania major* and *Trypanosoma cruzi* and low toxicity toward normal mammalian cells. *J Med Chem* 55:3867–3877. <http://dx.doi.org/10.1021/jm300070h>.
18. Lonardoni MVC, Russo M, Jancar S. 2000. Essential role of platelet-activating factor in control of *Leishmania (Leishmania) amazonensis* infection. *Infect Immun* 68:6355–6361. <http://dx.doi.org/10.1128/IAI.68.11.6355-6361.2000>.