

2017

## Expression of Ifnlr1 on intestinal epithelial cells is critical to the antiviral effects of IFN-lambda against norovirus and reovirus

Megan T. Baldrige  
*Washington University School of Medicine in St. Louis*

Sanghyun Lee  
*Washington University School of Medicine in St. Louis*

Judy J. Brown  
*Vanderbilt University Medical Center*

Nicole McAllister  
*University of Pittsburgh School of Medicine*

Kelly Urbanek  
*University of Pittsburgh School of Medicine*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Baldrige, Megan T.; Lee, Sanghyun; Brown, Judy J.; McAllister, Nicole; Urbanek, Kelly; Dermody, Terence S.; Nice, Timothy J.; and Virgin, Herbert W., "Expression of Ifnlr1 on intestinal epithelial cells is critical to the antiviral effects of IFN-lambda against norovirus and reovirus." *Journal of Virology*. 91, 7. e02079-16. (2017).

[https://digitalcommons.wustl.edu/open\\_access\\_pubs/5500](https://digitalcommons.wustl.edu/open_access_pubs/5500)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

---

**Authors**

Megan T. Baldrige, Sanghyun Lee, Judy J. Brown, Nicole McAllister, Kelly Urbanek, Terence S. Dermody, Timothy J. Nice, and Herbert W. Virgin



# Expression of *Ifnlr1* on Intestinal Epithelial Cells Is Critical to the Antiviral Effects of Interferon Lambda against Norovirus and Reovirus

Megan T. Baldrige,<sup>a\*</sup> Sanghyun Lee,<sup>a</sup> Judy J. Brown,<sup>b</sup> Nicole McAllister,<sup>c</sup> Kelly Urbaneck,<sup>d</sup> Terence S. Dermody,<sup>c,d</sup> Timothy J. Nice,<sup>e</sup> Herbert W. Virgin<sup>a</sup>

Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, Missouri, USA<sup>a</sup>; Department of Pathology, Microbiology, and Immunology, Vanderbilt University Medical Center, Nashville, Tennessee, USA<sup>b</sup>; Department of Microbiology and Molecular Genetics, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA<sup>c</sup>; Department of Pediatrics, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA<sup>d</sup>; Department of Molecular Microbiology and Immunology, Oregon Health and Science University, Portland, Oregon, USA<sup>e</sup>

**ABSTRACT** Lambda interferon (IFN- $\lambda$ ) has potent antiviral effects against multiple enteric viral pathogens, including norovirus and rotavirus, in both preventing and curing infection. Because the intestine includes a diverse array of cell types, however, the cell(s) upon which IFN- $\lambda$  acts to exert its antiviral effects is unclear. Here, we sought to identify IFN- $\lambda$ -responsive cells by generation of mice with lineage-specific deletion of the receptor for IFN- $\lambda$ , *Ifnlr1*. We found that expression of IFNLR1 on intestinal epithelial cells (IECs) in the small intestine and colon is required for enteric IFN- $\lambda$  antiviral activity. IEC *Ifnlr1* expression also determines the efficacy of IFN- $\lambda$  in resolving persistent murine norovirus (MNoV) infection and regulates fecal shedding and viral titers in tissue. Thus, the expression of *Ifnlr1* by IECs is necessary for the response to both endogenous and exogenous IFN- $\lambda$ . We further demonstrate that IEC *Ifnlr1* expression is required for the sterilizing innate immune effects of IFN- $\lambda$  by extending these findings in *Rag1*-deficient mice. Finally, we assessed whether our findings pertained to multiple viral pathogens by infecting mice specifically lacking IEC *Ifnlr1* expression with reovirus. These mice phenocopied *Ifnlr1*-null animals, exhibiting increased intestinal tissue titers and enhanced reovirus fecal shedding. Thus, IECs are the critical cell type responding to IFN- $\lambda$  to control multiple enteric viruses. This is the first genetic evidence that supports an essential role for IECs in IFN- $\lambda$ -mediated control of enteric viral infection, and these findings provide insight into the mechanism of IFN- $\lambda$ -mediated antiviral activity.

**IMPORTANCE** Human noroviruses (HNoVs) are the leading cause of epidemic gastroenteritis worldwide. Type III interferons (IFN- $\lambda$ ) control enteric viral infections in the gut and have been shown to cure mouse norovirus, a small-animal model for HNoVs. Using a genetic approach with conditional knockout mice, we identified IECs as the dominant IFN- $\lambda$ -responsive cells in control of enteric virus infection *in vivo*. Upon murine norovirus or reovirus infection, *Ifnlr1* depletion in IECs largely recapitulated the phenotype seen in *Ifnlr1*<sup>-/-</sup> mice of higher intestinal tissue viral titers and increased viral shedding in the stool. Moreover, IFN- $\lambda$ -mediated sterilizing immunity against murine norovirus requires the capacity of IECs to respond to IFN- $\lambda$ . These findings clarify the mechanism of action of this cytokine and emphasize the therapeutic potential of IFN- $\lambda$  for treating mucosal viral infections.

**KEYWORDS** innate immunity, interferons, mucosal immunity, norovirus, reovirus

Received 27 October 2016 Accepted 6 January 2017

Accepted manuscript posted online 11 January 2017

**Citation** Baldrige MT, Lee S, Brown JJ, McAllister N, Urbaneck K, Dermody TS, Nice TJ, Virgin HW. 2017. Expression of *Ifnlr1* on intestinal epithelial cells is critical to the antiviral effects of interferon lambda against norovirus and reovirus. *J Virol* 91:e02079-16. <https://doi.org/10.1128/JVI.02079-16>.

**Editor** Susana López, Instituto de Biotecnología/UNAM

**Copyright** © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Timothy J. Nice, [nice@ohsu.edu](mailto:nice@ohsu.edu), or Herbert W. Virgin, [virgin@wustl.edu](mailto:virgin@wustl.edu).

\* Present address: Megan T. Baldrige, Department of Medicine, Washington University School of Medicine, St. Louis, Missouri, USA.

M.T.B. and S.L. contributed equally to this work.

Norovirus and rotavirus are viral pathogens that infect at mucosal surfaces and induce gastroenteritis, characterized by vomiting, diarrhea, and malaise (1, 2). Viral gastroenteritis causes significant morbidity and mortality in children, the elderly, and immunocompromised persons, thus representing a substantial health care burden (3, 4). Treatments for these illnesses have been limited thus far to symptomatic care, including rehydration, because currently there is no specific antiviral therapy for these viral pathogens. Lambda interferon (IFN- $\lambda$ ; also called type III IFN) is an antiviral cytokine that regulates viral infection at mucosal surfaces and in the liver and brain (5–8). Administration of recombinant IFN- $\lambda$  can prevent and resolve viral infections in the gastrointestinal tract (8, 9) and at other sites in mice (10). These effects are observed for murine norovirus (MNoV) in mice lacking adaptive immunity, thus representing sterilizing innate immunity in the intestine (8). These studies indicate the potential for IFN- $\lambda$  as a therapeutic for viral infections, including those causing gastroenteritis, in humans, including immunocompromised hosts (11). Better understanding of the mechanisms by which this antiviral cytokine functions is essential to understanding basic mechanisms of intestinal control of viral infection and for potential therapeutic application in humans.

Binding of IFN- $\lambda$  to its receptor, a heterodimer of interleukin-10R2 (IL-10R2) and IFNLR1 (12, 13), induces an antiviral gene expression program similar to that induced by type I IFN, with substantial overlap in gene sets *in vitro* (10, 14, 15). However, type I and III IFNs exhibit unique antiviral properties *in vivo*. *Ifnlr1*<sup>-/-</sup> mice exhibit elevated intestinal tissue replication and enhanced fecal shedding of a persistent strain of MNoV (8, 16), a model virus which allows for more tractable *in vitro* and *in vivo* analyses than human norovirus (reviewed in references 17 and 18). Recombinant IFN- $\lambda$  treatment is sufficient to prevent and cure MNoV infection (8). In contrast, mice deficient for *Ifnar1* (the receptor for type I IFNs) show enhanced extraintestinal spread of virus, but levels of MNoV fecal shedding are comparable to those of wild-type mice (8, 16). Similarly, IFNLR1 restricts growth in the epithelium and fecal shedding of reovirus, while IFNAR1 instead regulates reovirus growth in the lamina propria (19). IFN- $\lambda$  exhibits an antiviral role exclusive of type I IFNs against a murine rotavirus strain (9) but cooperates with type I IFNs to limit intestinal replication of a heterologous simian strain in neonatal but not adult mice (20). These findings indicate the likely importance of tissue compartment-, development-, and cell type-specific effects of type I and III IFNs *in vivo*. These effects may be secondary to unique virulence factors that counter specific IFNs or to differential expression of the IFN receptors (21, 22).

IFNAR1 is thought to be expressed ubiquitously and at especially high levels on cells of hematopoietic origin (reviewed in references 23 and 24), whereas expression of detectable IFNLR1 appears to be limited to mucosal epithelial cells (25), human hepatocytes (6), and neutrophils (26). Although IFNLR1 expression on peripheral leukocytes has also been reported, it does not appear to be functional (27). Upon IFN- $\lambda$  treatment, IFN-stimulated genes accumulate in intestinal epithelial cells (IECs), indicating functional IFNLR1 expression (9, 19, 20). In contrast, in IECs of adult mice, IFNAR1 may be expressed at lower levels or alternately trafficked, such as only to the apical portion of the cell (9, 20). Differential receptor expression thus could account for complementary roles for different IFNs in protection against systemic infection (type I) and infection of mucosal (type III) sites. Importantly, however, it has been reported that cells that do not express detectably high levels of IFNLR1, such as the endothelial cells of the blood-brain barrier, may still respond to endogenous and exogenous IFN- $\lambda$  with protective antiviral effects (10). Thus, to successfully identify the cell types required for the antiviral response to IFN- $\lambda$ , analysis of receptor expression levels may be insufficient, and definitive resolution requires a genetic approach to selectively delete receptor expression in specific cell types.

To identify the cell types that respond to IFN- $\lambda$  *in vivo* in the intestine, we generated mice with a conditional mutant allele for *Ifnlr1* and crossed them to mice expressing Cre recombinase via the action of different cell type-specific promoters (Table 1). *Ifnlr1* was targeted in cell types expected to express high receptor levels (intestinal epithelial cells

**TABLE 1** Mouse lines, nomenclature, and cell types targeted by specific Cre lines


<i>Ifnlr1</i> and Cre mouse line(s)	Line name <sup>a</sup>	Cell type(s) targeted (reference)
<i>Ifnlr1</i> <sup>tm1Palu</sup> ; no Cre line	<i>Ifnlr1</i> <sup>-/-</sup>	All cells (28)
<i>Ifnlr1</i> <sup>tm1a(EUCOMM)Wtsi</sup> ; Villin-Cre	<i>Ifnlr1</i> <sup>f/f</sup> -Villincre	Intestinal epithelial cells (29)
<i>Ifnlr1</i> <sup>tm1a(EUCOMM)Wtsi</sup> ; MRP8-Cre	<i>Ifnlr1</i> <sup>f/f</sup> -MRP8cre	Neutrophils (30)
<i>Ifnlr1</i> <sup>tm1a(EUCOMM)Wtsi</sup> ; CD11c-Cre	<i>Ifnlr1</i> <sup>f/f</sup> -CD11ccre	Dendritic cells and alveolar macrophages (31)
<i>Ifnlr1</i> <sup>tm1a(EUCOMM)Wtsi</sup> ; LysM-Cre	<i>Ifnlr1</i> <sup>f/f</sup> -LysMcre	Macrophages, neutrophils, some dendritic cells (32, 33)
<i>Ifnlr1</i> <sup>tm1a(EUCOMM)Wtsi</sup> ; Deleter-Cre	<i>Ifnlr1</i> <sup>f-/-</sup>	All cells (34)

<sup>a</sup>A conditional allele of *Ifnlr1* (*Ifnlr1*<sup>f/f</sup>) was crossed to multiple different Cre lines for lineage-specific deletion of *Ifnlr1* in the specific cell types.

[25] and neutrophils [26]) and cells that are known to be permissive for MNoV replication in tissue culture (macrophages and dendritic cells [35]). Of all the cell types tested, only intestinal epithelial cells (IECs) required expression of *Ifnlr1* for the antiviral effects of IFN-λ against MNoV. To show the generality of our findings, we demonstrated the importance of IEC expression of this receptor for control of reovirus infection. This is the first study to genetically define IFN-λ-responsive cells *in vivo* in the context of two independent mucosal viral infections. This study also confirms that the cells required for responding to endogenous IFN-λ to attenuate MNoV infection are the same as those that respond to exogenous IFN-λ administration, including in the elicitation of sterilizing innate immunity.

**RESULTS**

***Ifnlr1* is expressed in the epithelial fraction along the length of the gastrointestinal tract.** Tissue from adult mice homozygous for a null mutation in *Ifnlr1* (28) or wild-type controls was collected from sites along the intestine, lung, mesenteric lymph node (MLN), or spleen (Fig. 1A). The small intestine was also dissociated into epithelial and lamina propria fractions as previously described (36), and RNA was isolated from




**FIG 1** *Ifnlr1* is expressed in the epithelial fraction along the length of the intestine. (A) RNA was isolated from sites along the intestine and the lung, indicated by red boxes, whole mesenteric lymph node (MLN) and spleen, and epithelial and lamina propria (LP) fractions of the small intestine from *Ifnlr1*-sufficient and -deficient mice. (B) *Ifnlr1* expression was quantified by quantitative real-time PCR of RNA from sites depicted in panel A. *n* = 4 to 6 samples per group, from two independent experiments, analyzed by Mann-Whitney test. \*\*, *P* < 0.01; ns, not significant. Prox, proximal; dist, distal.

these fractions and tissues. Expression of *Ifnlr1* was detected by quantitative real-time PCR of cDNA generated from these RNA samples. We found that *Ifnlr1* was expressed along the length of the intestine and in the lung, as well as in systemic tissues, including MLN and spleen (Fig. 1B). Intestinal *Ifnlr1* expression was substantially enriched (at least 30-fold;  $P = 0.0381$ ) in the epithelial fraction compared to the lamina propria fraction (Fig. 1B), consistent with previous reports (9, 25). As expected, no transcript was detected in any tissue in *Ifnlr1*<sup>-/-</sup> mice (Fig. 1B).

***Ifnlr1* expression in the small and large intestine is significantly diminished in *Ifnlr1*<sup>fl/fl</sup>-Villincre mice.** Embryonic stem (ES) cells targeted with a construct containing sequences homologous to *Ifnlr1*, an FLP recombinant target (FRT)-flanked *lacZ* and neomycin cassette, and loxP sites flanking exon 2 were provided by the Wellcome Trust Sanger Institute (Fig. 2A). Mice derived from these ES cells were crossed with mice expressing Flp recombinase for deletion of the FRT-flanked cassette (38), leaving a conditional allele of *Ifnlr1*, referred to as *Ifnlr1*<sup>f</sup> (Fig. 2A). Following removal of the floxed region in cells expressing Cre, the resulting transcript is predicted to produce a truncated protein product (Fig. 2B). For disruption in specific cell lineages, *Ifnlr1*<sup>fl/fl</sup> mice were crossed with various Cre mouse lines (Table 1). For each line, Cre(+) *Ifnlr1*<sup>fl/fl</sup> mice were compared to Cre(-) *Ifnlr1*<sup>fl/fl</sup> littermates to assess the effects of cell type-specific deletion upon *Ifnlr1* expression along the intestine and in extraintestinal tissues (Fig. 1A). *Ifnlr1*<sup>fl/fl</sup>-Villincre mice showed significantly diminished *Ifnlr1* expression in the small and large intestine (Fig. 2C). Fractionation of the small intestine into epithelial and lamina propria fractions revealed efficient deletion of *Ifnlr1* in the epithelium of these mice (Fig. 2C). In contrast, *Ifnlr1*<sup>fl/fl</sup>-MRP8cre, *Ifnlr1*<sup>fl/fl</sup>-LysMcre, and *Ifnlr1*<sup>fl/fl</sup>-CD11ccre mice showed no alterations in intestinal *Ifnlr1* expression at the level of the whole tissues tested (Fig. 2D, E, and F). *Ifnlr1*<sup>fl/fl</sup>-MRP8cre, *Ifnlr1*<sup>fl/fl</sup>-LysMcre, and *Ifnlr1*<sup>fl/fl</sup>-CD11ccre mice did exhibit substantial depletion of *Ifnlr1* in their respectively targeted cell types of neutrophils (~85%), macrophages (~91%), and dendritic cells (~85%), consistent with a previous report (39) (Fig. 2D, E, and F). Expression of *Ifnlr1* remained unchanged in lung, MLN, spleen, stomach, and duodenum in *Ifnlr1*<sup>fl/fl</sup>-Villincre mice, indicating expression of Cre specific to distal small intestine and colon (Fig. 2C), consistent with previous reports (29, 40).


**Expression of *Ifnlr1* in intestinal epithelium regulates MNoV shedding and response to recombinant IFN-λ.** *Ifnlr1*<sup>-/-</sup> mice and wild-type controls were inoculated with CR6, a persistent strain of MNoV that replicates well in the intestine, is shed into the feces at readily detectable levels, and is sensitive to treatment with IFN-λ (8, 41). As described previously (8), *Ifnlr1*<sup>-/-</sup> mice allow higher levels of fecal MNoV shedding than do wild-type mice at early time points (Fig. 3A and G) and are insensitive to IFN-λ treatment, although this treatment terminates MNoV replication in wild-type mice (Fig. 3A). These results were also observed in a novel *Ifnlr1*-deficient mouse model (*Ifnlr1*<sup>f-/-</sup>) (Table 1 and Fig. 3B). This assay was next applied to the four mouse strains with lineage-specific deletion of *Ifnlr1* (Table 1). *Ifnlr1*<sup>fl/fl</sup>-Villincre mice phenocopied *Ifnlr1*<sup>-/-</sup> and *Ifnlr1*<sup>f-/-</sup> mice, exhibiting both elevated fecal shedding of MNoV and resistance to IFN-λ treatment (Fig. 3C). In contrast, *Ifnlr1*<sup>fl/fl</sup>-MRP8cre, *Ifnlr1*<sup>fl/fl</sup>-LysMcre, and *Ifnlr1*<sup>fl/fl</sup>-CD11ccre mice exhibited viral loads and response to IFN-λ equivalent to those of *Ifnlr1*<sup>fl/fl</sup> controls (Fig. 3D, E, and F). At day 7 postinoculation, IFNLR1 regulated fecal shedding of MNoV, as seen by comparing wild-type and *Ifnlr1*<sup>-/-</sup> levels (Fig. 3G). *Ifnlr1*<sup>fl/fl</sup>-Villincre mice allowed fecal shedding equivalent to *Ifnlr1*<sup>-/-</sup> and *Ifnlr1*<sup>f-/-</sup> mice, suggesting that control of MNoV fecal shedding can be fully accounted for by IFNLR1 in Villin-expressing cells (Fig. 3G). Similarly, *Ifnlr1*<sup>fl/fl</sup>-Villincre mice exhibited no difference in comparison to *Ifnlr1*<sup>-/-</sup> and *Ifnlr1*<sup>f-/-</sup> mice along the full time course of infection (Fig. 3H).

**Expression of *Ifnlr1* in intestinal epithelium is essential for induction of IFN-λ-mediated sterilizing innate immunity to MNoV infection.** We previously reported that recombinant IFN-λ can cure persistently infected mice in the absence of adaptive immunity (8). To determine whether expression of *Ifnlr1* in IECs is required for IFN-λ-mediated sterilizing innate immunity to persistent MNoV infection, we established




**FIG 2** *Ifnlr1* expression is decreased in the small and large intestines of *Ifnlr1<sup>fl/fl</sup>-Villincre* mice. (A) Schematic depicting the *Ifnlr1* gene locus in *Ifnlr1<sup>tm1a(EUCOMM)Wtsi</sup>* mice. After crossing with mice expressing Flp recombinase (+Flp recombinase), the region between the two FRT sites was deleted, leaving conditional-ready *Ifnlr1<sup>fl/fl</sup>* mice. In the absence of Cre, all exons are present. With the addition of Cre recombinase, the floxed exon 2 is deleted. (B) In the absence of Cre [Cre(-)], the IFNL1 protein is expressed. In the presence of Cre [Cre(+)], the protein sequence is altered at amino acid 20 and a premature stop codon is introduced at amino acid 42. (C to F) *Ifnlr1* expression was assessed by quantitative real-time PCR of sites along the intestine and the lung, MLN and spleen, and epithelial and LP fractions from *Ifnlr1<sup>fl/fl</sup>-Villincre* (C), *Ifnlr1<sup>fl/fl</sup>-MRP8cre* (D), *Ifnlr1<sup>fl/fl</sup>-LysMcre* (E), and *Ifnlr1<sup>fl/fl</sup>-CD11ccre* (F) mice compared to their *Ifnlr1<sup>fl/fl</sup>* littermates. *Ifnlr1* expression was also assessed by quantitative real-time PCR of isolated bone marrow neutrophils from *Ifnlr1<sup>fl/fl</sup>-MRP8cre* (D), splenic macrophages from *Ifnlr1<sup>fl/fl</sup>-LysMcre* (E), and splenic dendritic cells from *Ifnlr1<sup>fl/fl</sup>-CD11ccre* (F) mice compared to their *Ifnlr1<sup>fl/fl</sup>* littermates. *n* = 4 to 7 samples per group, from two independent experiments, analyzed by Mann-Whitney test. \*, *P* < 0.05; \*\*, *P* < 0.01; ns, not significant.





**FIG 3** Expression of *Ifnlr1* on intestinal epithelial cells is required for the antiviral effects of endogenous and exogenous IFN-λ against MNoV. (A to E) Time course of MNoV genome copies shed into fecal pellets with time points at 7, 14, 21, 24, 28, and 35 days after CR6 infection. PBS or recombinant IFN-λ was injected intraperitoneally on day 21 into wild-type and *Ifnlr1*<sup>-/-</sup> (A), wild-type and *Ifnlr1*<sup>fl/fl</sup> (B), *Ifnlr1*<sup>fl/fl</sup>-Villin<sup>cre</sup> (C), *Ifnlr1*<sup>fl/fl</sup>-MRP8<sup>cre</sup> (D), *Ifnlr1*<sup>fl/fl</sup>-LysM<sup>cre</sup> (E), or *Ifnlr1*<sup>fl/fl</sup>-CD11<sup>cre</sup> (F) mice and their *Ifnlr1*<sup>fl/fl</sup> littermates. *n* = 6 to 12 mice per group, from two to three independent experiments, analyzed by two-way ANOVA followed by Tukey's multiple-comparison test; a *P* value of <0.001 by ANOVA column factor was found for panels A to F. (G) Individual data points depicting MNoV genome copies shed into fecal pellets on day 7 from panels A to F. *n* = 9 to 21 mice per group, from two to three independent experiments, analyzed by one-way ANOVA followed by Tukey's multiple-comparison test; a *P* value of <0.001 was determined by ANOVA. (H) Fecal shedding data from PBS-treated mice in panels A to C is shown superimposed to facilitate comparison between strains. *n* = 8 to 11 mice per group, from two to three independent experiments, analyzed by two-way ANOVA followed by Tukey's multiple-comparison test; a *P* value of <0.001 was determined by ANOVA column factor. \*, *P* < 0.05; \*\*, *P* < 0.01; \*\*\*, *P* < 0.001; ns, not significant.






**FIG 4** Expression of *Ifnl1* on intestinal epithelial cells is required for the antiviral effects of IFN-λ against MNoV in the absence of adaptive immunity. (A) Time course of MNoV genome copies shed into fecal pellets with time points at 7, 14, 21, 24, 28, and 35 days after CR6 infection. PBS or recombinant IFN-λ was injected intraperitoneally on day 21 and day 23 into *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* or *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup> mice. *n* = 6 to 14 mice per group, combined from three independent experiments, analyzed by two-way ANOVA followed by Tukey's multiple-comparison test; a *P* value of <0.001 was found by ANOVA column factor. (B) Individual data points depicting MNoV genome copies shed into fecal pellets on day 7 from *Rag1*<sup>-/-</sup>, *Rag1*<sup>-/-</sup> *Ifnl1*<sup>-/-</sup> double knockouts or mice depicted in panel A. *n* = 6 to 22 mice per group, combined from two to three independent experiments, analyzed by Mann-Whitney test. \*\*\*, *P* < 0.001; ns, not significant.

*Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* conditional double knockout mice. *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* mice were orally inoculated with CR6, and viral shedding in the stool was quantified by quantitative PCR (qPCR). *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* mice showed increased viral shedding throughout the infection time course (Fig. 4A). Injection of recombinant IFN-λ terminated MNoV replication in *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup> mice but did not affect MNoV loads in *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* mice (Fig. 4A). At 7 days postinoculation, *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* mice had significantly higher viral shedding than *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup> mice, and the level of viral shedding in *Rag1*<sup>-/-</sup> *Ifnl1*<sup>ff</sup>-*Villincre* mice was comparable to the level of viral shedding in *Rag1*<sup>-/-</sup> *Ifnl1*<sup>-/-</sup> mice (Fig. 4B). Therefore, IFN-λ responses in IECs limited persistent MNoV infection in the absence of adaptive immunity, and IFN-λ signaling in IECs was essential for clearance of persistently infected MNoV by IFN-λ-mediated sterilizing innate immunity.

**Control of reovirus in intestinal tissue by IFN-λ depends upon the expression of *Ifnl1* in epithelial cells.** To assess whether *Ifnl1* expression on IECs was required for control of other enteric pathogens, *Ifnl1*<sup>-/-</sup> and *Ifnl1*<sup>ff</sup>-*Villincre* mice were orally inoculated with 10<sup>8</sup> PFU of reovirus strain type 1 Lang (T1L). At 4 days postinfection, viral titers in small intestinal tissues, including duodenum, jejunum, and ileum, as well as viral shedding in stools, were significantly higher in *Ifnl1*<sup>-/-</sup> mice (Fig. 5A and B), consistent with a previous report using another strain of reovirus, type 3 Dearing (19). Control of reovirus was predominantly through the expression of IFNLR1 on IECs, as *Ifnl1*<sup>ff</sup>-*Villincre* mice displayed increased titers of reovirus in small intestinal tissues as well as enhanced fecal shedding (Fig. 5A and B). These results demonstrate that expression of IFNLR1 in epithelial cells is essential for the control of reovirus infection by IFN-λ in the gut and indicate that IFN-λ signaling in IECs is an antiviral mechanism common to multiple enteric viral pathogens.

**Interferon-stimulated gene expression in the intestine depends upon the expression of *Ifnl1* in epithelial cells.** Ileum and proximal colon tissues were isolated from wild-type (WT), *Ifnl1*<sup>-/-</sup>, *Ifnl1*<sup>ff</sup>, and *Ifnl1*<sup>ff</sup>-*Villincre* mice 1 day posttreatment with either PBS or IFN-λ. These tissues were then assessed for expression of canonical antiviral interferon-stimulated genes (ISGs) *Oas1a* (42), *Ift1* (43), and *Ifl44* (44) (Fig. 6A




**FIG 5** *Ifnlr1* expression on intestinal epithelial cells limits reovirus infection. (A and B) Titers of reovirus strain T1L were assessed at day 4 postinoculation in the different compartments of the small intestine (A) and stool (B) from wild-type, *Ifnlr1*<sup>-/-</sup>, *Ifnlr1*<sup>fl/fl</sup>-Villincre, and *Ifnlr1*<sup>fl/fl</sup> littermate control mice. The small intestine was resected from the pylorus to the cecum and sectioned into three equal parts, representing the duodenum, jejunum, and ileum. Titers are expressed as PFU per milliliter of tissue homogenate or gram of stool. *n* = 6 to 8 mice per group, combined from two independent experiments, analyzed by one-way ANOVA followed by Tukey's multiple-comparison test; a *P* value of <0.001 was determined by ANOVA column factor for all tissues and stool. \*, *P* < 0.05; \*\*, *P* < 0.01; \*\*\*, *P* < 0.001; ns, not significant.

to C). While intestinal tissues from WT and *Ifnlr1*<sup>fl/fl</sup> mice exhibited robust ISG induction in response to IFN- $\lambda$  treatment, tissues from *Ifnlr1*<sup>-/-</sup> and *Ifnlr1*<sup>fl/fl</sup>-Villincre mice failed to significantly upregulate these ISGs in response to IFN- $\lambda$ . These data correlate with the impaired antiviral response against MNoV in *Ifnlr1*<sup>-/-</sup> and *Ifnlr1*<sup>fl/fl</sup>-Villincre mice after IFN- $\lambda$  treatment (Fig. 3A and C), consistent with a potentially critical role for IFNLR1 expression on epithelial cells for induction of antiviral ISGs in response to IFN- $\lambda$  treatment.

## DISCUSSION

In this study, we found that IECs are the predominant cell type expressing *Ifnlr1* in the small intestine and colon and that this cell type plays a major role in IFN- $\lambda$ -mediated antiviral immunity in the intestine. Antiviral immunity elicited by IFN- $\lambda$  to enteric reovirus and norovirus infection depends upon IFNLR1 signaling in Villin-positive IECs. Using four mouse strains with lineage-specific deletion of *Ifnlr1* to study persistent infection and IFN- $\lambda$ -mediated clearance, we found that only *Ifnlr1*<sup>fl/fl</sup>-Villincre mice exhibited a complete phenocopy of *Ifnlr1*<sup>-/-</sup> mice. Targeting *Ifnlr1* in other cells, including dendritic cells, macrophages, and neutrophils, had no detectable effect on basal levels of viral shedding or IFN- $\lambda$ -mediated clearance of MNoV. The dominant IFN- $\lambda$ -dependent antiviral contribution by IECs also was confirmed with reovirus infection. In studies using reovirus T1L, we observed viral titers in the small intestine of *Ifnlr1*<sup>fl/fl</sup>-Villincre mice increased comparably to those detected in *Ifnlr1*<sup>-/-</sup> mice, although we cannot rule out a minor role for other IFN- $\lambda$ -responding cells in the ileum. Therefore, IECs are the functionally dominant IFN- $\lambda$ -responding cells for endogenous and exogenous IFN- $\lambda$  control of viruses in the intestine. There is a clear correlation between IFN- $\lambda$ -mediated induction of antiviral ISGs and IEC expression of IFNLR1, suggesting that induction of ISGs in IECs is the mechanism by which IFN- $\lambda$  exerts its antiviral effects.

Expression of *Ifnlr1* mRNA throughout the gut and in other extraintestinal tissues (MLN, lung, and spleen) was quantified by qPCR analysis. In lamina propria cells, there were fewer than 500 copies of *Ifnlr1* mRNA per 1  $\mu$ g total RNA. In contrast, IECs express



**FIG 6** *Ifnlr1* expression on intestinal epithelial cells is necessary for induction of interferon-stimulated genes. *Oas1a* (A), *Ifit1* (B), and *Ifi44* (C) expression was assessed by quantitative real-time PCR of RNA from distal ileum and proximal colon tissue from wild-type (WT), *Ifnlr1*<sup>-/-</sup>, *Ifnlr1*<sup>fl/fl</sup>, and *Ifnlr1*<sup>fl/fl</sup>-VillinCre mice at 1 day posttreatment with PBS or recombinant IFN- $\lambda$ . *n* = 5 to 9 mice per group, combined from two independent experiments, analyzed by one-way ANOVA followed by Tukey's multiple-comparison test; a *P* value of <0.001 was determined by ANOVA column factor for all tissues. \*\*\*, *P* < 0.001; ns, not significant.

more than 10,000 copies of *Ifnlr1* mRNA per 1  $\mu$ g total RNA throughout the small intestine. Thus, IECs are the dominant *Ifnlr1*-expressing cells and function as the major IFN- $\lambda$ -responding cells for antiviral immunity in the intestine. In MLN, spleen, and lung, we detected comparable expression of *Ifnlr1* mRNA. Villin-positive cells were not the major cell type responsible for *Ifnlr1* expression in these tissues, and neither were neutrophils, dendritic cells, or macrophages. Therefore, there may exist some other cell types that are important for IFN- $\lambda$  responses in these tissues. Lung epithelial cells, which do not express Villin, likely reflect a major source of *Ifnlr1* in that tissue (29). B or T cells, which have been reported to express *Ifnlr1* even though they lack a robust response to IFN- $\lambda$ , may account for *Ifnlr1* expression in the MLN and spleen (27). Another possible cellular source for this expression is endothelial cells, based on the report that blood-brain barrier endothelial cells respond to IFN- $\lambda$  to restrict West Nile virus neuroinvasion (10). Thus, it would be interesting to study the role of IFN- $\lambda$  in extraintestinal tissues in control of other pathogens and define the IFN- $\lambda$ -responsive cell types in these contexts.

In some tissues, such as lung and vagina, there is redundancy between type I and III IFN-mediated antiviral responses. IFN- $\lambda$  controls influenza virus, severe acute respiratory syndrome (SARS) coronavirus, respiratory syncytial virus infection in the lung (45–47), and herpes simplex virus 2 (HSV-2) infection in the genital tract (48), redundantly with type I IFNs. In the intestine, however, IFN- $\lambda$ -mediated antiviral immunity does not redundantly overlap type I IFNs (8, 9, 19). Adult IECs have polarized apical IFNAR1 expression only at low levels (9), and although IECs in neonates exhibit robust STAT1 activation after type I IFN treatment, in adult mice they are largely unresponsive to type I IFN treatment *in vivo* (9, 19, 20). Moreover, the expression level of *Ifnlr1* mRNA is highly enriched in IECs but minimally detectable in other compartments of intestinal tissue (Fig. 1 and 2). This study, bolstering previous findings of alternate cellular expression patterns for type I and III IFN receptors, helps explain why IFN- $\lambda$ -mediated immunity in the intestine is nonredundant with IFN- $\alpha/\beta$  in adult mice, even though they may stimulate transcription of highly overlapping sets of antiviral genes (7, 20). Our data support a role for IECs as sentinels for enteric virus infection via their response to compartment-specific IFN- $\lambda$  signaling (19, 20).

One of the important features of IFN- $\lambda$ -mediated immunity is its sterilizing activity against persistent MNoV infection in the absence of adaptive immunity (8). We observed that persistent MNoV infection of *Rag1*<sup>-/-</sup> *Ifnlr1*<sup>f/f</sup>-Villin<sup>cre</sup> mice was not resolved by IFN- $\lambda$  treatment and showed increased viral titers in the stool, similar to our observations with *Rag1*<sup>-/-</sup> *Ifnlr1*<sup>-/-</sup> mice. Thus, IFN- $\lambda$ -mediated sterilizing innate immunity requires IEC expression of the receptor. Since only macrophages, dendritic cells, and B cells are known to be susceptible to MNoV infection *in vitro* (35, 49), it is not clear how the IFN- $\lambda$  response in IECs ablates persistent MNoV infection in the absence of adaptive immunity. One possible explanation is that there is a secondary *trans*-acting molecule induced by IFN- $\lambda$  in IECs that clears MNoV in other cell types. A related study has demonstrated that rotavirus can be terminated by injecting IL-22 and IL-18 into *Rag1*<sup>-/-</sup> mice, but this IL-22- and IL-18-mediated viral clearance does not induce IFN- $\lambda$  or Stat1 activation (50). Thus, there may be multiple innate immunological mechanisms to resolve persistent viral infection in the absence of adaptive immunity. Identifying the effectors of IFN- $\lambda$ -mediated sterilizing immunity is an important area to pursue in IFN- $\lambda$  immunology.

This study reveals that *Ifnlr1* expression in IECs is required for control of enteric MNoV and reovirus infections. Using a genetic approach with conditional knockout mice, we identified IECs as the dominant cell type that responds to endogenous and exogenous IFN- $\lambda$  to control enteric viruses. Understanding the identity of IFN- $\lambda$ -responsive cell types provides further insight into mechanisms that control enteric viruses and will enhance future development of IFN- $\lambda$ -mediated antiviral therapeutics.

## MATERIALS AND METHODS

**Generation of MNoV stocks and determination of titers.** Stocks of MNoV strain CR6 were generated from a molecular clone as previously described (51). Briefly, a plasmid encoding the CR6 genome was transfected into 293T cells to generate infectious virus, which was subsequently passaged on BV2 cells. After two passages, BV2 cultures were frozen and thawed to liberate virions. Cultures then were cleared of cellular debris and virus was concentrated by ultracentrifugation through a 30% sucrose cushion. Titers of virus stocks were determined by plaque assay on BV2 cells (51).

**Generation of reovirus stocks and determination of titers.** Spinner-adapted murine L929 (L) cells were grown in either suspension or monolayer cultures in Joklik's modified Eagle's minimal essential medium (SMEM; Lonza) supplemented to contain 5% fetal bovine serum (Gibco), 2 mM L-glutamine, 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin (Gibco), and 25 ng/ml amphotericin B (Sigma). BHK-T7 cells were grown in Dulbecco's modified Eagle's minimal essential medium (DMEM; Gibco) supplemented to contain 5% fetal bovine serum, 2 mM L-glutamine, 1 mg/ml Geneticin (Gibco), and nonessential amino acids (Sigma).

Recombinant reoviruses were generated using plasmid-based reverse genetics (52). Recombinant strain type 1 Lang (T1L) is a stock generated by plasmid-based rescue from cloned T1L cDNAs (53). After 3 to 5 days of incubation, cells were frozen and thawed three times, and virus was isolated by plaque purification using monolayers of L cells (54). Purified reovirus virions were generated from second- or third-passage L-cell lysate stocks (55). Viral particles were extracted from infected cell lysates using Vertrel XF (Dupont), layered onto 1.2- to 1.4-g/cm<sup>3</sup> CsCl gradients, and centrifuged at 62,000  $\times$  g for 16 h. Bands corresponding to virions (1.36 g/cm<sup>3</sup>) were collected and dialyzed in virion storage buffer (150 mM NaCl, 15 mM MgCl<sub>2</sub>, and 10 mM Tris-HCl [pH 7.4]) (56). The concentration of reovirus virions in purified preparations was determined from the following equivalence: one optical density (OD) unit at 260 nm equals 2.1  $\times$  10<sup>12</sup> virions (56). Viral titer was determined by plaque assay using L cells (54).

For analysis of reoviral titers in organs, mice were euthanized at various intervals postinoculation, and organs were harvested into 1 ml of PBS and homogenized by freeze-thaw and bead beating. For analysis of viral titer in stool, samples were harvested at various intervals, weighed, stored in 1 ml of PBS, and homogenized by freeze-thaw and bead beating. Viral titers in organs and stool homogenates were determined by plaque assay using L cells (54). Titers are expressed as PFU per milliliter of tissue homogenate or per gram of stool.

**Mice, infections, and IFN- $\lambda$  treatment.** Wild-type (WT) C57BL/6J mice (stock number 000664) were purchased from Jackson Laboratories (Bar Harbor, ME) and housed at the Washington University School of Medicine under specific-pathogen-free conditions (57) according to university guidelines. *Ifnlr1*<sup>-/-</sup> (B6.Cg-*Ifnlr1*<sup>tm1Palu</sup>) mice were obtained from Bristol-Myers Squibb (Seattle, WA) and backcrossed using speed congenics onto a C57BL/6J background (28).

To generate mice conditionally deficient for *Ifnlr1*, *Ifnlr1*<sup>tm1a(EUCOMM)Wtsi</sup> ES cells on a C57BL/6N background were provided by the Wellcome Trust Sanger Institute. A conditional-ready (floxed) allele in which exon 2 is flanked by loxP sites, designated *Ifnlr1*<sup>flf</sup>, was created (Fig. 2A) (38). *Ifnlr1*<sup>flf</sup> mice were crossed to Villin-Cre (intestinal epithelial cells [29]), LysM-Cre (macrophages and neutrophils, as well as some dendritic cells [32, 33]), CD11c-Cre (dendritic cells and alveolar macrophages [31]), and MRP8-Cre (neutrophils [30]) lines for selective disruption of *Ifnlr1* in different cell types *in vivo*. *Ifnlr1*<sup>flf</sup> mice were also crossed to a Deleter-Cre line (34) to generate an alternate *Ifnlr1*<sup>-/-</sup> line, here designated *Ifnlr1*<sup>f-/-</sup>. *Ifnlr1*<sup>f-/-</sup> mice were backcrossed using speed congenics onto a C57BL/6J background. Mouse lines and naming conventions are summarized in Table 1.

For MNoV infections, mice were inoculated with a dose of 10<sup>6</sup> PFU of strain CR6 at 6 to 8 weeks of age by the oral route in a volume of 25  $\mu$ l. For reovirus infections, mice were orally gavaged with a dose of 10<sup>8</sup> PFU of strain T1L virus at 6 to 8 weeks in a volume of 100  $\mu$ l.

Recombinant IFN- $\lambda$  was provided by Bristol-Myers Squibb (Seattle, WA) as a monomeric conjugate comprised of 20-kDa linear polyethylene glycol (PEG) attached to the amino terminus of murine IFN- $\lambda$ , as previously reported (8). For treatment of mice, 25  $\mu$ g of IFN- $\lambda$  diluted in PBS was injected intraperitoneally.

Stool and tissues were harvested into 2-ml tubes (Sarstedt, Germany) with 1-mm-diameter zirconia/silica beads (Biospec, Bartlesville, OK). Tissues were flash frozen in a bath of ethanol and dry ice and either processed on the same day or stored at -80°C.

**Isolation of epithelial and lamina propria fractions of small intestine.** Epithelial and lamina propria fractions were prepared as previously described (36). In brief, after mice were euthanized, small intestines were collected. Intestinal tissues were washed with cold PBS twice and then chopped and transferred to new tubes. The tissues were incubated with stripping buffer (10% bovine calf serum, 15 mM HEPES, 5 mM EDTA, 5 mM dithiothreitol [DTT] in 1  $\times$  Hanks' balanced salt solution [HBSS]) for 20 min at 37°C. The dissociated cells were collected as the epithelial fraction, consisting predominantly of IECs. The remaining tissue was used as the lamina propria fraction.

**Isolation of neutrophils, macrophages, and dendritic cells.** Neutrophils were isolated from *Ifnlr1*<sup>flf-MRP8cre</sup> mice and *Ifnlr1*<sup>flf</sup> littermates by collecting bone marrow from femurs and tibiae. Red blood cells were lysed using red blood cell lysis buffer (Sigma, St. Louis, MO), and neutrophils were isolated using the mouse neutrophil isolation kit (Miltenyi Biotec, Germany). Isolated neutrophils were confirmed to be 95 to 98% double positive for CD11b-allophycocyanin (APC) and Ly6G-fluorescein isothiocyanate (FITC) (BioLegend, San Diego, CA) (data not shown). Macrophages were isolated from *Ifnlr1*<sup>flf-LysMcre</sup> mice and *Ifnlr1*<sup>flf</sup> littermates by collecting and homogenizing spleens, lysing red blood cells (RBCs), and enriching for macrophages using mouse anti-F4/80 UltraPure MicroBeads (Miltenyi Biotec). Isolated macrophages were confirmed to be 70 to 85% positive for F4/80-AF488

(Thermo Fisher Scientific) as well as CD11b-APC and CD45.2-phycoerythrin (PE) (BioLegend) (data not shown). Dendritic cells were isolated from *Ifnlr1<sup>fl/fl</sup>-CD11ccre* mice and *Ifnlr1<sup>fl/fl</sup>* littermates by collecting and homogenizing spleens, lysing RBCs, and enriching for dendritic cells using the mouse pan-dendritic cell isolation kit (Miltenyi Biotec). Isolated dendritic cells were confirmed to be 70 to 85% CD11c-AF488 (BioLegend) single positive or CD11c-AF488 and B220-PE (BD Bioscience) double positive (data not shown).

**Quantitative reverse transcription-PCR.** RNA from stool was isolated using a ZR-96 viral RNA kit (Zymo Research, Irvine, CA). RNA from tissues or cells was isolated using TRI Reagent (Invitrogen) and a direct-zol-96 RNA kit (Zymo Research, Irvine, CA) according to the manufacturer's protocol. Five microliters of RNA from stool or 1  $\mu$ g of RNA from tissue was used as a template for cDNA synthesis with the ImPromII reverse transcriptase system (Promega, Madison, WI). DNA contamination was removed using the DNFree kit (Life Technologies).

MNoV TaqMan assays were performed, using a standard curve for determination of absolute viral genome copies, as described previously (58). Quantitative PCR for housekeeping gene *Rps29* was performed with forward primer 5'-GCAAATACGGGCTGAACATG-3', reverse primer 5'-GTCCAACCTAATG AAGCCTATGTC-3', and probe 5'-/5HEX/CCTTCGCGT/ZEN/ACTGCCGAAGC/3IABkFQ/-3' (where 3IABkFQ is 3' Iowa Black fluorescence quencher; Integrated DNA Technologies), each at a concentration of 0.2  $\mu$ M, using AmpliTaq gold DNA polymerase (Applied Biosystems). Quantitative PCRs for *Ifnlr1* (Mm.PT.58.10781457), *Oas1a* (Mm.PT.58.30459792), *Ifi44* (Mm.PT.58.12162024), and *Ifit1* (Mm.PT.58.32674307) were similarly performed using PrimeTime qPCR assays (Integrated DNA Technologies). Standard curves for quantitative PCR assays were generated to facilitate absolute quantification of transcript copy numbers. For *Rps29*, the PCR product using the above-described primers was cloned into the p-ENTR/D-TOPO vector (Thermo Fisher Scientific), and for *Ifnlr1* a full-length *Ifnlr1* cDNA clone (5036481; Open Biosystems) was used. Plasmids were Sanger sequenced to confirm the identity of the inserts. For *Oas1a*, *Ifit1*, and *Ifi44*, absolute transcripts were quantitated based on target sequence-containing gBlocks (Integrated DNA Technologies). Cycling parameters for *Rps29*, *Ifnlr1*, *Oas1a*, *Ifit1*, and *Ifi44* were identical to those for MNoV TaqMan. Absolute values of *Ifnlr1*, *Oas1a*, *Ifit1*, and *Ifi44* per microgram of RNA were normalized to the within-tissue average of housekeeping gene *Rps29*. No significant changes in absolute copy number of *Rps29* were detected between comparison groups (data not shown).

**Statistical analysis.** Data were analyzed with Prism 7 software (GraphPad Software, San Diego, CA). In all graphs, three asterisks indicate a *P* value of <0.001, two asterisks indicate a *P* value of <0.01, one asterisk indicates a *P* value of <0.05, and ns indicates not significant (*P* > 0.05) as determined by Mann-Whitney test, one-way analysis of variance (ANOVA), or two-way ANOVA with Tukey's multiple-comparison test, as specified in the relevant figure legends.

## ACKNOWLEDGMENTS

We thank D. Kreamalmeyer for animal care and breeding, S. Peterson for technical assistance, members of the Virgin laboratory for manuscript review and discussion, S. Doyle and Bristol-Myers Squibb for providing *Ifnlr1<sup>tm1Palu</sup>* mice, and Bill Skarnes and Wendy Bushell at the Wellcome Trust Sanger Institute for providing *Ifnlr1<sup>tm1a(EUCOMM)Wtsi</sup>* ES cells. Experimental support was provided by the Speed Con- genics Facility of the Rheumatic Diseases Core Center.

H.W.V. was supported by National Institutes of Health (NIH) grant U19 AI109725 and the Crohn's and Colitis Foundation grant 326556. M.T.B. was supported by NIH training grant 5T32CA009547 and the W. M. Keck Fellowship from Washington University. T.J.N. was supported by NIH training grant 5T32A100716334 and postdoctoral fellowships from the Cancer Research Institute and American Cancer Society. S.L. was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Education (NRF-2016R1A6A3A03012352). J.B. was supported by NIH training grant 5T32HL007751 and predoctoral fellowship F31DK108562. T.S.D. was supported by NIH grant R01 AI038296. Research reported in this publication was supported by NIH award number P30AR048335. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## REFERENCES

- Karst SM. 2010. Pathogenesis of noroviruses, emerging RNA viruses. *Viruses* 2:748–781. <https://doi.org/10.3390/v2030748>.
- Glass RI, Parashar UD, Estes MK. 2009. Norovirus gastroenteritis. *N Engl J Med* 361:1776–1785. <https://doi.org/10.1056/NEJMra0804575>.
- Koo HL, Ajami N, Atmar RL, DuPont HL. 2010. Noroviruses: the leading cause of gastroenteritis worldwide. *Discov Med* 10:61–70.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 17:7–15. <https://doi.org/10.3201/eid1701.P11101>.
- Lazear HM, Nice TJ, Diamond MS. 2015. Interferon-lambda: immune functions at barrier surfaces and beyond. *Immunity* 43:15–28. <https://doi.org/10.1016/j.immuni.2015.07.001>.
- Hermant P, Michiels T. 2014. Interferon-lambda in the context of viral infections: production, response and therapeutic implications. *J Innate Immun* 6:563–574. <https://doi.org/10.1159/000360084>.



7. Egli A, Santer DM, O'Shea D, Tyrrell DL, Houghton M. 2014. The impact of the interferon-lambda family on the innate and adaptive immune response to viral infections. *Emerg Microbes Infect* 3:e51. <https://doi.org/10.1038/emi.2014.51>.
8. Nice TJ, Baldrige MT, McCune BT, Norman JM, Lazear HM, Artyomov M, Diamond MS, Virgin HW. 2015. Interferon-lambda cures persistent murine norovirus infection in the absence of adaptive immunity. *Science* 347:269–273. <https://doi.org/10.1126/science.1258100>.
9. Pott J, Mahlakoiv T, Mordstein M, Duerr CU, Michiels T, Stockinger S, Staeheli P, Hornef MW. 2011. IFN-lambda determines the intestinal epithelial antiviral host defense. *Proc Natl Acad Sci U S A* 108:7944–7949. <https://doi.org/10.1073/pnas.1100552108>.
10. Lazear HM, Daniels BP, Pinto AK, Huang AC, Vick SC, Doyle SE, Gale M, Jr, Klein RS, Diamond MS. 2015. Interferon-lambda restricts West Nile virus neuroinvasion by tightening the blood-brain barrier. *Sci Transl Med* 7:284ra259.
11. Muir AJ, Arora S, Everson G, Flisiak R, George J, Ghalib R, Gordon SC, Gray T, Greenbloom S, Hassanein T, Hillson J, Horga MA, Jacobson IM, Jeffers L, Kowdley KV, Lawitz E, Lueth S, Rodriguez-Torres M, Rustgi V, Shemanski L, Shiffman ML, Srinivasan S, Vargas HE, Vierling JM, Xu D, Lopez-Talavera JC, Zeuzem S, EMERGE Study Group. 2014. A randomized phase 2b study of peginterferon lambda-1a for the treatment of chronic HCV infection. *J Hepatol* 61:1238–1246. <https://doi.org/10.1016/j.jhep.2014.07.022>.
12. Kutenko SV, Gallagher G, Baurin VV, Lewis-Antes A, Shen M, Shah NK, Langer JA, Sheikh F, Dickensheets H, Donnelly RP. 2003. IFN-lambdas mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat Immunol* 4:69–77. <https://doi.org/10.1038/ni875>.
13. Sheppard P, Kindsvogel W, Xu W, Henderson K, Schlutsmeyer S, Whitmore TE, Kuestner R, Garrigues U, Birks C, Roraback J, Ostrand C, Dong D, Shin J, Presnell S, Fox B, Haldeman B, Cooper E, Taft D, Gilbert T, Grant FJ, Tackett M, Krivan W, McKnight G, Clegg C, Foster D, Klucher KM. 2003. IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol* 4:63–68. <https://doi.org/10.1038/ni873>.
14. Bolen CR, Ding S, Robek MD, Kleinstein SH. 2014. Dynamic expression profiling of type I and type III interferon-stimulated hepatocytes reveals a stable hierarchy of gene expression. *Hepatology* 59:1262–1272. <https://doi.org/10.1002/hep.26657>.
15. Kohli A, Zhang X, Yang J, Russell RS, Donnelly RP, Sheikh F, Sherman A, Young H, Imamichi T, Lempicki RA, Masur H, Kottlilil S. 2012. Distinct and overlapping genomic profiles and antiviral effects of Interferon-lambda and -alpha on HCV-infected and noninfected hepatoma cells. *J Viral Hepat* 19:843–853. <https://doi.org/10.1111/j.1365-2893.2012.01610.x>.
16. Baldrige MT, Nice TJ, McCune BT, Yokoyama CC, Kambal A, Wheadon M, Diamond MS, Ivanova Y, Artyomov M, Virgin HW. 2015. Commensal microbes and interferon-lambda determine persistence of enteric murine norovirus infection. *Science* 347:266–269. <https://doi.org/10.1126/science.1258025>.
17. Karst SM, Wobus CE, Goodfellow IG, Green KY, Virgin HW. 2014. Advances in norovirus biology. *Cell Host Microbe* 15:668–680. <https://doi.org/10.1016/j.chom.2014.05.015>.
18. Wobus CE, Thackray LB, Virgin HW. 2006. Murine norovirus: a model system to study norovirus biology and pathogenesis. *J Virol* 80:5104–5112. <https://doi.org/10.1128/JVI.02346-05>.
19. Mahlakoiv T, Hernandez P, Gronke K, Diefenbach A, Staeheli P. 2015. Leukocyte-derived IFN-alpha/beta and epithelial IFN-lambda constitute a compartmentalized mucosal defense system that restricts enteric virus infections. *PLoS Pathog* 11:e1004782. <https://doi.org/10.1371/journal.ppat.1004782>.
20. Lin JD, Feng N, Sen A, Balan M, Tseng HC, McElrath C, Smirnov SV, Peng J, Yasukawa LL, Durbin RK, Durbin JE, Greenberg HB, Kutenko SV. 2016. Distinct roles of type I and type III interferons in intestinal immunity to homologous and heterologous rotavirus infections. *PLoS Pathog* 12:e1005600. <https://doi.org/10.1371/journal.ppat.1005600>.
21. Visel A, Thaller C, Eichele G. 2004. GenePaint.org: an atlas of gene expression patterns in the mouse embryo. *Nucleic Acids Res* 32:D552–D556. <https://doi.org/10.1093/nar/gkh029>.
22. Diez-Roux G, Banfi S, Sultan M, Geffers L, Anand S, Rozado D, Magen A, Canidio E, Pagani M, Peluso I, Lin-Marq N, Koch M, Bilio M, Cantiello I, Verde R, De Masi C, Bianchi SA, Cicchini J, Perroud E, Mehmeti S, Dagand E, Schrinner S, Nurnberger A, Schmidt K, Metz K, Zwimgmann C, Brieske N, Springer C, Hernandez AM, Herzog S, Grabbe F, Sieverding C, Fischer B, Schrader K, Brockmeyer M, Dettmer S, Helbig C, Alunni V, Battaini MA, Mura C, Henrichsen CN, Garcia-Lopez R, Echevarria D, Puellas E, Garcia-Calero E, Kruse S, Uhr M, Kauck C, Feng G, Milyaev N, Ong CK, Kumar L, Lam M, Semple CA, Gyenesei A, Mundlos S, Radelof U, Lehrach H, Sarmientos P, Raymond A, Davidson DR, Dolle P, Antonarakis SE, Yaspo ML, Martinez S, Baldock RA, Eichele G, Ballabio A. 2011. A high-resolution anatomical atlas of the transcriptome in the mouse embryo. *PLoS Biol* 9:e1000582. <https://doi.org/10.1371/journal.pbio.1000582>.
23. Randall RE, Goodbourn S. 2008. Interferons and viruses: an interplay between induction, signalling, antiviral responses and virus countermeasures. *J Gen Virol* 89:1–47. <https://doi.org/10.1099/vir.0.83391-0>.
24. de Weerd NA, Samarajiva SA, Hertzog PJ. 2007. Type I interferon receptors: biochemistry and biological functions. *J Biol Chem* 282:20053–20057. <https://doi.org/10.1074/jbc.R700006200>.
25. Sommereyns C, Paul S, Staeheli P, Michiels T. 2008. IFN-lambda (IFN-lambda) is expressed in a tissue-dependent fashion and primarily acts on epithelial cells in vivo. *PLoS Pathog* 4:e1000017. <https://doi.org/10.1371/journal.ppat.1000017>.
26. Blazek K, Eames HL, Weiss M, Byrne AJ, Perocheau D, Pease JE, Doyle S, McCann F, Williams RO, Udalova IA. 2015. IFN-lambda resolves inflammation via suppression of neutrophil infiltration and IL-1beta production. *J Exp Med* 212:845–853. <https://doi.org/10.1084/jem.20140995>.
27. Witte K, Gruetz G, Volk HD, Looman AC, Asadullah K, Sterry W, Sabat R, Wolk K. 2009. Despite IFN-lambda receptor expression, blood immune cells, but not keratinocytes or melanocytes, have an impaired response to type III interferons: implications for therapeutic applications of these cytokines. *Genes Immun* 10:702–714. <https://doi.org/10.1038/gene.2009.72>.
28. Ank N, Iversen MB, Bartholdy C, Staeheli P, Hartmann R, Jensen UB, Dagnaes-Hansen F, Thomsen AR, Chen Z, Haugen H, Klucher K, Paludan SR. 2008. An important role for type III interferon (IFN-lambda/IL-28) in TLR-induced antiviral activity. *J Immunol* 180:2474–2485. <https://doi.org/10.1049/jimmunol.180.4.2474>.
29. Madison BB, Dunbar L, Qiao XT, Braunstein K, Braunstein E, Gumucio DL. 2002. Cis elements of the villin gene control expression in restricted domains of the vertical (crypt) and horizontal (duodenum, cecum) axes of the intestine. *J Biol Chem* 277:33275–33283. <https://doi.org/10.1074/jbc.M204935200>.
30. Passegue E, Wagner EF, Weissman IL. 2004. JunB deficiency leads to a myeloproliferative disorder arising from hematopoietic stem cells. *Cell* 119:431–443. <https://doi.org/10.1016/j.cell.2004.10.010>.
31. Stranges PB, Watson J, Cooper CJ, Choisy-Rossi CM, Stonebraker AC, Beighton RA, Hartig H, Sundberg JP, Servick S, Kaufmann G, Fink PJ, Chervovsky AV. 2007. Elimination of antigen-presenting cells and autoreactive T cells by Fas contributes to prevention of autoimmunity. *Immunity* 26:629–641. <https://doi.org/10.1016/j.immuni.2007.03.016>.
32. Clausen BE, Burkhardt C, Reith W, Renkawitz R, Forster I. 1999. Conditional gene targeting in macrophages and granulocytes using LysMcre mice. *Transgenic Res* 8:265–277. <https://doi.org/10.1023/A:1008942828960>.
33. Jakubczik K, Bogunovic M, Bonito AJ, Kuan EL, Merad M, Randolph GJ. 2002. Lymph-migrating, tissue-derived dendritic cells are minor constituents within steady-state lymph nodes. *J Exp Med* 205:2839–2850. <https://doi.org/10.1084/jem.20081430>.
34. Schwenk F, Baron U, Rajewsky K. 1995. A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Res* 23:5080–5081. <https://doi.org/10.1093/nar/23.24.5080>.
35. Wobus CE, Karst SM, Thackray LB, Chang KO, Sosnovtsev SV, Belliot G, Krug A, Mackenzie JM, Green KY, Virgin HW. 2004. Replication of Norovirus in cell culture reveals a tropism for dendritic cells and macrophages. *PLoS Biol* 2:e432. <https://doi.org/10.1371/journal.pbio.0020432>.
36. Lefrancois L, Lycke N. 2001. Isolation of mouse small intestinal intraepithelial lymphocytes, Peyer's patch, and lamina propria cells. *Curr Protoc Immunol* Chapter 3:Unit 3.19.
37. Reference deleted.
38. Kanki H, Suzuki H, Itohara S. 2006. High-efficiency CAG-FLPe deleter mice in C57BL/6J background. *Exp Anim* 55:137–141. <https://doi.org/10.1538/expanim.55.137>.
39. Abram CL, Roberge GL, Hu Y, Lowell CA. 2014. Comparative analysis of the efficiency and specificity of myeloid-Cre deleting strains using ROSA-EYFP reporter mice. *J Immunol Methods* 408:89–100. <https://doi.org/10.1016/j.jim.2014.05.009>.
40. Zhong Z, Baker JJ, Zylstra-Diegel CR, Williams BO. 2012. Lrp5 and Lrp6 play compensatory roles in mouse intestinal development. *J Cell Biochem* 113:31–38. <https://doi.org/10.1002/jcb.23324>.



41. Nice TJ, Strong DW, McCune BT, Pohl CS, Virgin HW. 2013. A single-amino-acid change in murine norovirus NS1/2 is sufficient for colonic tropism and persistence. *J Virol* 87:327–334. <https://doi.org/10.1128/JVI.01864-12>.
42. Elkhateeb E, Tag-El-Din-Hassan HT, Sasaki N, Torigoe D, Morimatsu M, Agui T. 2016. The role of mouse 2',5'-oligoadenylate synthetase 1 paralogs. *Infect Genet Evol* 45:393–401. <https://doi.org/10.1016/j.meegid.2016.09.018>.
43. Reynaud JM, Kim DY, Atasheva S, Rasalouslykaya A, White JP, Diamond MS, Weaver SC, Frolova EI, Frolov I. 2015. IFIT1 differentially interferes with translation and replication of alphavirus genomes and promotes induction of type I interferon. *PLoS Pathog* 11:e1004863. <https://doi.org/10.1371/journal.ppat.1004863>.
44. Carlton-Smith C, Elliott RM. 2012. Viperin, MTAP44, and protein kinase R contribute to the interferon-induced inhibition of Bunyamwera Orthobunyavirus replication. *J Virol* 86:11548–11557. <https://doi.org/10.1128/JVI.01773-12>.
45. Crotta S, Davidson S, Mahlakoiv T, Desmet CJ, Buckwalter MR, Albert ML, Staeheli P, Wack A. 2013. Type I and type III interferons drive redundant amplification loops to induce a transcriptional signature in influenza-infected airway epithelia. *PLoS Pathog* 9:e1003773. <https://doi.org/10.1371/journal.ppat.1003773>.
46. Mordstein M, Kochs G, Dumoutier L, Renaud JC, Paludan SR, Klucher K, Staeheli P. 2008. Interferon-lambda contributes to innate immunity of mice against influenza A virus but not against hepatotropic viruses. *PLoS Pathog* 4:e1000151. <https://doi.org/10.1371/journal.ppat.1000151>.
47. Mordstein M, Neugebauer E, Ditt V, Jessen B, Rieger T, Falcone V, Sorgeloos F, Ehl S, Mayer D, Kochs G, Schwemmler M, Gunther S, Drosten C, Michiels T, Staeheli P. 2010. Lambda interferon renders epithelial cells of the respiratory and gastrointestinal tracts resistant to viral infections. *J Virol* 84:5670–5677. <https://doi.org/10.1128/JVI.00272-10>.
48. Ank N, West H, Bartholdy C, Eriksson K, Thomsen AR, Paludan SR. 2006. Lambda interferon (IFN-lambda), a type III IFN, is induced by viruses and IFNs and displays potent antiviral activity against select virus infections in vivo. *J Virol* 80:4501–4509. <https://doi.org/10.1128/JVI.80.9.4501-4509.2006>.
49. Jones MK, Watanabe M, Zhu S, Graves CL, Keyes LR, Grau KR, Gonzalez-Hernandez MB, Iovine NM, Wobus CE, Vinje J, Tibbetts SA, Walle SM, Karst SM. 2014. Enteric bacteria promote human and mouse norovirus infection of B cells. *Science* 346:755–759. <https://doi.org/10.1126/science.1257147>.
50. Zhang B, Chassaing B, Shi Z, Uchiyama R, Zhang Z, Denning TL, Crawford SE, Puijssers AJ, Iskarpatyoti JA, Estes MK, Dermody TS, Ouyang W, Williams IR, Vijay-Kumar M, Gewirtz AT. 2014. Viral infection. Prevention and cure of rotavirus infection via TLR5/NLRC4-mediated production of IL-22 and IL-18. *Science* 346:861–865.
51. Strong DW, Thackray LB, Smith TJ, Virgin HW. 2012. Protruding domain of capsid protein is necessary and sufficient to determine murine norovirus replication and pathogenesis in vivo. *J Virol* 86:2950–2958. <https://doi.org/10.1128/JVI.07038-11>.
52. Kobayashi T, Antar AA, Boehme KW, Danthi P, Eby EA, Guglielmi KM, Holm GH, Johnson EM, Maginnis MS, Naik S, Skelton WB, Wetzel JD, Wilson GJ, Chappell JD, Dermody TS. 2007. A plasmid-based reverse genetics system for animal double-stranded RNA viruses. *Cell Host Microbe* 1:147–157. <https://doi.org/10.1016/j.chom.2007.03.003>.
53. Kobayashi T, Ooms LS, Ikizler M, Chappell JD, Dermody TS. 2010. An improved reverse genetics system for mammalian orthoreoviruses. *Virology* 398:194–200. <https://doi.org/10.1016/j.virol.2009.11.037>.
54. Virgin HWT, Bassel-Duby R, Fields BN, Tyler KL. 1988. Antibody protects against lethal infection with the neurally spreading reovirus type 3 (Dearing). *J Virol* 62:4594–4604.
55. Furlong DB, Nibert ML, Fields BN. 1988. Sigma 1 protein of mammalian reoviruses extends from the surfaces of viral particles. *J Virol* 62:246–256.
56. Smith RE, Zweerink HJ, Joklik WK. 1969. Polypeptide components of virions, top component and cores of reovirus type 3. *Virology* 39:791–810. [https://doi.org/10.1016/0042-6822\(69\)90017-8](https://doi.org/10.1016/0042-6822(69)90017-8).
57. Cadwell K, Patel KK, Maloney NS, Liu TC, Ng AC, Storer CE, Head RD, Xavier R, Stappenbeck TS, Virgin HW. 2010. Virus-plus-susceptibility gene interaction determines Crohn's disease gene Atg16L1 phenotypes in intestine. *Cell* 141:1135–1145. <https://doi.org/10.1016/j.cell.2010.05.009>.
58. Baert L, Wobus CE, Van Coillie E, Thackray LB, Debevere J, Uyttendaele M. 2008. Detection of murine norovirus 1 by using plaque assay, transfection assay, and real-time reverse transcription-PCR before and after heat exposure. *Appl Environ Microbiol* 74:543–546. <https://doi.org/10.1128/AEM.01039-07>.