

2017

The retina visual cycle is driven by cis retinol oxidation in the outer segments of cones

Shinya Sato
Washington University School of Medicine in St. Louis

Rikard Frederiksen
Boston University

M. Carter Cornwall
Boston University

Vladimir J. Kefalov
Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Sato, Shinya; Frederiksen, Rikard; Cornwall, M. Carter; and Kefalov, Vladimir J., "The retina visual cycle is driven by cis retinol oxidation in the outer segments of cones." *Visual Neuroscience*. 34, e004. (2017).
https://digitalcommons.wustl.edu/open_access_pubs/5726

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

BRIEF COMMUNICATION

The retina visual cycle is driven by *cis* retinol oxidation in the outer segments of cones

SHINYA SATO,¹ RIKARD FREDERIKSEN,² M. CARTER CORNWALL,² AND VLADIMIR J. KEFALOV¹

¹Department of Ophthalmology and Visual Sciences, Washington University School of Medicine, Saint Louis, Missouri 63110

²Department of Physiology and Biophysics, Boston University School of Medicine, Boston, MA 02118

(RECEIVED December 8, 2016; ACCEPTED January 9, 2017)

Abstract

Vertebrate rod and cone photoreceptors require continuous supply of chromophore for regenerating their visual pigments after photoactivation. Cones, which mediate our daytime vision, demand a particularly rapid supply of 11-*cis* retinal chromophore in order to maintain their function in bright light. An important contribution to this process is thought to be the chromophore precursor 11-*cis* retinol, which is supplied to cones from Müller cells in the retina and subsequently oxidized to 11-*cis* retinal as part of the retina visual cycle. However, the molecular identity of the *cis* retinol oxidase in cones remains unclear. Here, as a first step in characterizing this enzymatic reaction, we sought to determine the subcellular localization of this activity in salamander red cones. We found that the onset of dark adaptation of isolated salamander red cones was substantially faster when exposing directly their outer *vs.* their inner segment to 9-*cis* retinol, an analogue of 11-*cis* retinol. In contrast, this difference was not observed when treating the outer *vs.* inner segment with 9-*cis* retinal, a chromophore analogue which can directly support pigment regeneration. These results suggest, surprisingly, that the *cis*-retinol oxidation occurs in the outer segments of cone photoreceptors. Confirming this notion, pigment regeneration with exogenously added 9-*cis* retinol was directly observed in the truncated outer segments of cones, but not in rods. We conclude that the enzymatic machinery required for the oxidation of recycled *cis* retinol as part of the retina visual cycle is present in the outer segments of cones.

Keywords: Cone photoreceptors, Pigment regeneration, Dark adaptation, Retinol dehydrogenase, Visual cycle

Introduction

The vertebrate visual pigment, the light receptor molecule of rod and cone photoreceptor cells, is composed of a protein moiety, opsin, and chromophore, typically 11-*cis* retinal. Upon absorbing light, 11-*cis* retinal is photoisomerized to its all-*trans* form to activate the pigment before eventually dissociating from opsin [reviewed in (Ebrey & Koutalos, 2001; Imamoto & Shichida, 2014)]. The continuous function of photoreceptor cells requires regeneration of their visual pigments with newly synthesized 11-*cis* retinal. In vertebrate eyes, there are two metabolic pathways for 11-*cis* retinal synthesis: the retinal pigment epithelium (RPE) visual cycle and the retina visual cycle (Wang & Kefalov, 2011; Saari, 2012; Tang et al., 2013; Wright et al., 2015). The RPE visual cycle is the canonical pathway in which 11-*cis* retinal is synthesized in the RPE cells and supplied to both rods and cones. The retina visual cycle, on the other hand, is a recently discovered pathway in which 11-*cis* retinol is synthesized in Müller cells, supplied to cones, and then oxidized there to 11-*cis* retinal. This *cis* retinol oxidation

activity is cone-specific (Jones et al., 1989; Miyazono et al., 2008; Ala-Laurila et al., 2009; Sato & Kefalov, 2016). As a result, the retina visual cycle can supply chromophore exclusively to cones, enhancing in that way their fast dark adaptation and their function in bright light (Mata et al., 2002; Kolesnikov et al., 2011; Wang et al., 2014). However, little is known about the molecular mechanisms of the retina visual cycle. Although recent studies have identified several putative components of the pathway for 11-*cis* retinoid synthesis in Müller cells (Kaylor et al., 2013; Kaylor et al., 2014; Xue et al., 2015), the enzyme(s) responsible for the *cis* retinol oxidation in cones is still unknown.

Two hypotheses about this cone-specific reaction have been recently proposed. One suggestion is that the oxidation of *cis* retinol is carried out in the outer segments of cones by retinol dehydrogenase 8 (RDH8). This idea is based on the observation that deletion of RDH8 blocks the *cis* retinol-dependent sensitivity recovery in mouse M-cones (Kolesnikov et al., 2015). However, RDH8 is expressed in the outer segments of both rods and cones (Maeda et al., 2005; Miyazono et al., 2008) and is thought to be primarily a reductase of all-*trans* retinal rather than an oxidase of *cis* retinol (Rattner et al., 2000). The alternative hypothesis is that the oxidation of *cis* retinol takes place in cone inner segments and is carried out by RDH13-like (RDH13L) and its homologs (Sato et al., 2015).

Address correspondence to: Vladimir J. Kefalov, Ph.D, Department of Ophthalmology and Visual Sciences, Washington University in Saint Louis, 660 S. Euclid Avenue, Saint Louis, MO 63110. E-mail: kefalov@wustl.edu

RDH13L is a cone-specific enzyme identified from purified carp cone inner segment membranes. It oxidizes *cis* retinol with concomitant reduction of NADP⁺ or hydrophobic aldehydes. However, RDH13L is found only in fish and is, therefore, not a component of the mammalian retina visual cycle. Although its functional homolog, RDH14 (Sato et al., 2015), is suggested to be expressed in mouse cones (Kanan et al., 2008), its role in the cone-specific oxidation of *cis* retinol has not been examined.

Here, as a way of addressing this issue, we sought to identify the cellular compartment of salamander red cones where *cis* retinol oxidation takes place. The salamander retina has a high percentage of cones (~40%) (Sherry et al., 1998) and its photoreceptors are >5 times larger in width than those of the mouse retina [see for instance (Koutalos & Cornwall, 2010)]. These features allowed us targeted application of *cis* retinal or *cis* retinol selectively to the inner or outer segment of individual cones in an effort to determine the sub-cellular localization of the *cis* retinol oxidation activity in cones.

Materials and methods

Single-cell suction recordings

All experiments were performed in accordance with protocols approved by the Washington University Animal Studies Committee and the Institutional Animal Care and Use Committee of Boston University School of Medicine, following the Guide for the Care and Use of Laboratory Animals and the Animal Welfare Act. Larval tiger salamanders (*Ambystoma tigrinum*, Charles D. Sullivan Co. Inc., Nashville, TN) were maintained on a 12-h day–night cycle and were dark-adapted overnight before use. Single-cell recordings from salamander cones were carried out as previously described (Kefalov et al., 2001). For the cone dark-adaptation experiments, the inner or outer segment of a red cone was first drawn in the suction pipette and after measuring its dark current and sensitivity, the cell was exposed to brief bright 505 nm light estimated to bleach >99% of its visual pigment. Bleaching light duration was determined from the relation $F = 1 - \exp(-IPt)$, where F is the fraction of pigment to be bleached (>0.99), t is the duration of light exposure (>18 s), I is the light intensity (1.2×10^8 photons· $\mu\text{m}^{-2}\cdot\text{sec}^{-1}$), and P is the photosensitivity of salamander red cones at 505 nm (2.3×10^{-9} μm^2), estimated from the action spectrum of salamander red cones (Makino & Dodd, 1996; Sato & Kefalov, 2016) and the reported photosensitivity value at the wavelength of peak absorbance (6.0×10^{-9} μm^2) (Jones et al., 1993). Following the bleach, the cell was incubated in darkness for 20 min to allow the photoactivated visual pigments to decay to free opsin. The bleach-adapted cell was then treated with 800 μl of 125 μM retinoid solution which was added to the recording chamber. The retinoid solution was prepared as follows: 300 μg of dry retinoid (9-*cis* retinol: Santa Cruz Biotechnology, Inc., Santa Cruz, CA; 9-*cis* retinal: Sigma-Aldrich, Inc., Saint Louis, MO) was dissolved into 11 μl ethanol and kept on ice until used. Just before adding to the recording chamber, 1 μl of the ethanolic retinoid solution was mixed with 100 μl of Ringer's solution first and then additional 700 μl of Ringer's solution was added gradually to obtain 125 μM retinoid solution. The retinoid solution thus prepared was added into the recording chamber containing ~2 ml of perfusion solution (i.e., final concentration: ~40 μM) using a pipette with a wide-bore tip. To optimize the delivery of retinoids to the cone held in the suction electrode, the solution in the chamber was stirred by flushing some of it in and out of the pipette 3 times. As the solution was applied to the bath, only the

segment of the cell exposed to it and not drawn in the suction electrode was treated with the retinoid (Jin et al., 1994). To monitor the recovery of cone sensitivity following the application of the retinoid, dim flash responses (with amplitudes below 30% of the maximum recorded from each cone) were recorded every 1 min for 9-*cis* retinal-treated cones or every 2 min for 9-*cis* retinol-treated cones, up to 20 min after the addition of retinoid solution. Flash sensitivities were determined as the ratio of dim flash response amplitude and flash intensity. Electrophysiology data were analyzed using pClamp 9 (Molecular Devices, LLC, Sunnyvale, CA) and Origin 7.5 (OriginLab, Northampton, MA).

Microspectrophotometry measurements

The microspectrophotometry measurements were performed on a previously described instrument (Frederiksen et al., 2012). At the beginning of an MSP experiment, a dark-adapted retina was chopped and triturated. Solution containing isolated intact cells and outer segments was placed on a quartz cover-slip window located in the bottom of a 2 mm-deep Plexiglass recording chamber. The tissue was allowed to settle and adhere to the quartz window for up to 10 min to minimize movement of the cells, after which the tissue in the chamber was superfused at a rate of 4 ml/min with Ringer's solution identical to that used in the electrophysiological experiments. At this time, an isolated intact photoreceptor cell or outer segments were identified from their projected image on the screen of the infrared video system. A baseline absorbance spectrum was measured in an area adjacent to the selected outer segment. Following this measurement, the outer segment was oriented in the beam path with the polarization of the incident measuring beam parallel to the plane of the disks (rods) or invaginations (cones), and absorbance spectra in this area were measured in steps of 2 nm step width. Generally, ten complete sample scans and ten baseline scans were averaged to increase the signal-to-noise ratio of a measured spectrum. The absorbance spectrum was calculated using Beers' Law according to the relation; $\text{OD} = \log_{10}(I_0/I_t)$, where OD is optical density or absorbance, I_0 is the light transmitted through a cell-free space adjacent to the outer segment, and I_t is the light transmitted through the outer segment.

Dark-adapted spectra of the visual pigment were measured in this way from rod and cone outer segments. Once the dark-adapted spectral absorbance was established, the pigment was bleached (>99%) using a calibrated (505 nm; 1.3×10^7 photons· $\mu\text{m}^{-2}\cdot\text{sec}^{-1}$) LED light source. Bleaching photoproducts were monitored periodically and were allowed to decay to undetectable levels. This period was 10 min for cone outer segments, and about 60 min for rods. Following this, we measured the bleached absorbance spectrum. Next, we treated the preparation with 1 ml of 40 μM 9-*cis* retinol for 10 min. Spectral absorbance was measured and the pigment regeneration was monitored until no further change in spectral absorbance could be detected. Then the regenerated pigment (cones) was bleached once more to confirm that it was photosensitive. If no pigment was regenerated (rods) the preparation was treated with 1 ml of 40 μM 11-*cis* retinal for 10 min and allowed to regenerate visual pigment.

Statistics

A two-tailed unpaired Student's *t*-test was used to test for the significance of difference in the mean values of two sample groups. *P*-values of <0.05 were considered to be statistically significant.




Fig. 1. Kinetics of red cone sensitivity recovery when exposing the inner segment to exogenous 9-*cis* chromophore. (A) Image of a red cone in the outer segment-in configuration under microscope with infrared illumination. (B) Dim flash responses from bleached red cones at indicated times after treatment with 9-*cis* retinal (black traces) and 9-*cis* retinol (red traces). Response amplitudes are normalized to the flash intensity (i.e. response

Results

Location of the cis retinol oxidation reaction in salamander red cones

In the living retina, fresh 11-*cis* retinol is provided by the Müller cells to cones, and ultimately used for pigment regeneration in the cone outer segments. However, as Müller cell processes do not make direct contact with the cone outer segments (Sarantis & Mobbs, 1992; Garlipp & Gonzalez-Fernandez, 2013), their 11-*cis* retinol is most likely delivered to the cone inner segments (Insinna et al., 2012). It is currently unclear whether the subsequent oxidation of 11-*cis* retinol to 11-*cis* retinal takes place in the inner or outer segments of cones. To determine the location of *cis* retinol recovery in cones, we compared the time course of sensitivity recovery of fully bleached (>99% of visual pigment) isolated salamander red cones in a suction electrode when their outer vs. inner segment was exposed to 9-*cis* retinol, a commercially available more stable analogue of 11-*cis* retinol. As the suction electrode fits tightly around the photoreceptor cell, chromophore movement from the bath into the electrode was restricted so that only the part of the cell outside the electrode was exposed to it (Jin et al., 1994). Thus, we reasoned that exposing directly the part of the cell where oxidation takes place to 9-*cis* retinol would result in faster pigment regeneration compared to the reverse cell configuration. Isolated salamander red cones were drawn into the suction pipette from the outer segment to the mid-point of the inner segment (outer segment-in configuration; Fig. 1A) or from the cell body to the base of the outer segment (outer segment-out configuration; Fig. 2A). Cone pigments were fully bleached (>99%) by 505 nm light and then the cells were kept in darkness for 20 min until their sensitivity reached a steady state. At this point, 9-*cis* retinal or 9-*cis* retinol was added to the bath of the recording chamber and the change in cone sensitivity was monitored for the next 10 min for 9-*cis* retinal or 20 min for 9-*cis* retinol. Although the relation between the fraction of bleached pigment and sensitivity is non-linear (Jones et al., 1993; Kefalov et al., 2005), physiological measurements of cone sensitivity provide a reliable method for monitoring the onset and kinetics of cone pigment regeneration. Thus, we used the kinetics of cone dark adaptation to estimate the rate of pigment regeneration in the outer segment, and the initial delay in this process to estimate the time required for the production/delivery of 9-*cis* retinal to the outer segment.

We began by treating the inner segments of bleached red cones with 9-*cis* retinal in order to determine the kinetics of delivery of this retinoid to the outer segment and the subsequent rate of cone pigment regeneration in a configuration similar to that *in vivo*, with the retinoid delivered from the Müller cells. As 9-*cis* retinal can be used directly for pigment regeneration, the dark adaptation of cones in this case was rate-limited by its delivery to the outer segment. For outer segment-in cones (Fig. 1A), sensitivity recovery began shortly upon exposure to 9-*cis* retinal (Fig. 1B, black traces). The recovery onset was delayed by ~1 min (Fig. 1C, black filled squares), possibly reflecting the time required for 9-*cis* retinal taken up by the inner segment from the bath to reach the outer segment of the cell, sheltered inside the suction electrode. The subsequent dark adaptation of the cones proceeded rapidly and was largely completed within 5–6 min.

per photon) to allow direct comparison of sensitivity. (C) Sensitivity recovery time course of bleached red cones in the outer segment-in (OS-in) configuration incubated with 9-*cis* retinal (filled black squares; $n = 10$) or 9-*cis* retinol (filled red circles; $n = 8$). Data are shown as mean \pm SEM.




Fig. 2. Kinetics of red cone sensitivity recovery when exposing the outer segment to exogenous 9-*cis* chromophore. (A) Image of a red cone in the outer segment-out configuration under microscope with infrared illumination. (B) Sensitivity recovery time course of bleached red cones in the outer segment-out (OS-out) configuration incubated with 9-*cis* retinal (open black squares; $n = 7$) or 9-*cis* retinol (open red circles; $n = 8$). Data are shown as mean \pm SEM.

We next treat the inner segments of bleached cones with 9-*cis* retinol, a functional analogue to the 11-*cis* retinol produced by Müller cells in the retina (Sato & Kefalov, 2016). In comparison with 9-*cis* retinal, use of 9-*cis* retinol requires the additional step of oxidation before it could be used for pigment regeneration. Thus, the overall time course of cone dark adaptation in this case was determined by two processes: production of 9-*cis* retinal and its delivery to the outer segment. As expected, the administration of 9-*cis* retinol to the inner segments resulted in gradual recovery of cone sensitivity, indicating that cones could oxidize 9-*cis* retinol and use it for pigment regeneration. However, this recovery was delayed by ~ 6 min (Fig. 1B, red traces; Fig. 1C, filled red circles) from the time of retinoid application. This delay in the onset of cone recovery with 9-*cis* retinol was substantially larger than the ~ 1 min delay observed for 9-*cis* retinal in the same configuration.

The slight transient decrease in sensitivity upon the application of 9-*cis* retinol was likely due to toxicity of its high concentration in the bath when the perfusion was stopped during retinoid application. Indeed, this decrease was not observed when the 9-*cis* retinol solution was diluted 10-fold and the perfusion was not halted during its application, thus rapidly removing excess retinoid from the recording chamber (data not shown). The kinetics of the 9-*cis* retinol-driven recovery were also >3 -fold slower than the recovery driven by 9-*cis* retinal so that cone dark adaptation was still incomplete 20 min after treatment. Thus, when the retinoid was applied to the cone inner segment, the recovery driven by 9-*cis* retinol was delayed by several min and proceeded >3 -fold slower than the recovery driven by 9-*cis* retinal. This could reflect the different traffic rates of 9-*cis* retinal and 9-*cis* retinol through the cone inner segment, or could be caused by the extra oxidation step of 9-*cis* retinol to 9-*cis* retinal. However, based only on the results above, we could not determine which of these two factors causes the delay or the slower recovery of cones when their inner segments were treated with 9-*cis* retinol vs. 9-*cis* retinal.

To address this question and distinguish between these two possibilities, we treated directly the outer segments of bleached cones with 9-*cis* retinal and 9-*cis* retinol by using the outer segment-out configuration (Fig. 2A). When applied directly to the outer segment, 9-*cis* retinal could be immediately taken up by the cone and began driving pigment regeneration and dark adaptation within seconds (Fig. 2B; black open squares). Direct comparison of cones treated with 9-*cis* retinal in the outer segment-in (Fig. 3A, open squares) and outer segment-out (Fig. 3A, filled squares) configurations demonstrated that their kinetics of dark adaptation were comparable. The only difference was the ~ 1 min delay in cone recovery when the chromophore was applied to the inner segment of the cones, compared to its direct application to the outer segment. This delay when 9-*cis* retinal is applied to the cone inner segment is comparable to that previously observed with 11-*cis* retinal (Jin et al., 1994) and likely reflects the time required for the retinoid to reach the outer segment. Notably, the ensuing 9-*cis* retinal-driven pigment regeneration in our experiments was independent of the site of retinoid application and completed within 5–6 min.

Having established the time course of dark adaptation driven by exogenous *cis* retinal in the two cell configurations, we next compared the kinetics of cone dark adaptation with *cis* retinol. Pigment regeneration in this case requires the additional step of oxidation to *cis* retinal before pigment regeneration. To investigate the subcellular localization and the kinetics of *cis* retinol oxidation in cones, we compared the kinetics of 9-*cis* retinol-driven cone dark adaptation in the outer-segment-in configuration with that of bleached cones in the outer segment-out configuration. Notably, similar to the treatment with 9-*cis* retinal in this configuration, and unlike the treatment with 9-*cis* retinol in the reverse configuration, the recovery of cone sensitivity in this case began immediately after the application of the retinoid (Fig. 2B, red open circles). Thus, cones could rapidly and efficiently utilize 9-*cis* retinol for robust pigment regeneration when the retinoid was applied directly to their outer segments. The kinetics of the 9-*cis* retinol-driven recovery of sensitivity were slower than those of the 9-*cis* retinal-driven recovery in this configuration (Fig. 2B), but comparable to those of the 9-*cis* retinol-driven recovery in the outer segment-in configuration (Fig. 3B). In both recording configurations, the kinetics of recovery with 9-*cis* retinol was ~ 3 -fold slower than that with 9-*cis* retinal (Figs. 1C and 2B). Together, these results suggest that the delay in cone recovery when 9-*cis* retinol is applied to the inner segment is most likely due to the slow transport of this retinoid to the outer segment. The immediate




Fig. 3. Comparison of the kinetics of bleached red cone recovery in outer segment-in vs. outer segment-out configurations (data replotted from Fig. 1C and 2B). (A) Sensitivity recovery time course of bleached red cones incubated with 9-*cis* retinal in the OS-out (filled black squares) or OS-in (open black squares) configuration. (B) Sensitivity recovery time course of bleached red cones incubated with 9-*cis* retinal in the OS-out (filled red circles) or OS-in (open red circles) configuration.

onset of cone recovery when 9-*cis* retinal is applied directly to the outer segment is consistent with the notion that its oxidation to *9-cis* retinal occurs in the outer segment itself rather than in the inner segment. The ~3-fold slower dark adaptation with 9-*cis* retinal over 9-*cis* retinal likely reflects the relatively slow oxidation of *cis* retinal to *cis* retinal in the outer segments of cones.

cis Retinol oxidation in the truncated cone outer segment

Our physiological data is consistent with the presence of *cis* retinal oxidation activity in cone outer segments. However, the spatial isolation between the outer and inner segments by our suction electrode was likely not ideal. Thus, the possibility still existed that in the outer segment-out configuration, 9-*cis* retinal from the bath

could reach the tip of the inner segment and get oxidized there instead of in the outer segment. A second complication of the interpretation of our recordings arose from the slight decrease of cone sensitivity in the outer segment-in configuration upon the application of 9-*cis* retinal (Fig. 1C). This transient sensitivity suppression, which was not observed in the outer segment-out configuration, could also have masked any early onset of cone recovery mediated by oxidation in the inner segment. To rule out these possibilities, we tested directly the ability of truncated salamander cone outer segments to oxidize 9-*cis* retinol and use it for pigment regeneration. As suction recordings from truncated cone outer segments are quite challenging, we used microspectrophotometry (MSP) to measure directly pigment regeneration in 9-*cis* retinol-treated cone outer segments broken off from the rest of the cell. Truncated outer segments from dark-adapted red cones were identified by their shape and their characteristic absorption spectrum (Fig. 4A) with a peak at 610 ± 4 nm (mean ± s.d., n = 4), reflecting the mix of 11-*cis* A₁ and 11-*cis* A₂ chromophore in the retina of larval tiger salamanders (Makino et al., 1999). The outer segments were then bleached by bright light, which, as expected, resulted in the ablation of the pigment peak in the absorption spectrum (Fig. 4B). Critically, treatment of such bleached truncated cone outer segments with 9-*cis* retinal resulted in robust 9-*cis* pigment regeneration as demonstrated by the reappearance of a blue-shifted pigment spectrum (Fig. 4C) with a peak at 544 ± 9 nm (mean ± s.d., n = 4). This peak wavelength is comparable to the reported value for the 9-*cis* pigment in salamander red cones (542 nm) (Makino et al., 1999). The absorption peak disappeared after an additional bright light illumination, confirming that it originated from bleachable pigment and not from nonspecific retinoid products. This result clearly demonstrates that 9-*cis* retinal can be oxidized in the truncated cone outer segment and used for 9-*cis* pigment formation. As a control, the same experiments were done with intact salamander red rods (Fig. 4E) following a bleach (Fig. 4F). In this case, pigment regeneration was not observed after 9-*cis* retinal treatment (Fig. 4G). However, subsequent treatment with 11-*cis* retinal produced robust pigment regeneration (Fig. 4H). The inability of salamander rods to utilize 9-*cis* retinal for pigment regeneration is consistent with our recent finding that 9-*cis* retinol oxidation activity is cone-specific (Sato & Kefalov, 2016). Importantly, the lack of regeneration in rods under these conditions also ruled out contamination with 9-*cis* retinal in our 9-*cis* retinal preparation. Together, these MSP results substantiate our findings from single-cell recordings above and clearly demonstrate that the oxidation of 9-*cis* retinol takes place in cone outer segments and can drive robust cone pigment regeneration.

Discussion

Oxidation and traffic of 9-cis retinol in cones

This study builds on two elegant previous studies that showed that *cis* retinol can drive cone pigment regeneration (Jones et al., 1989) and that the movement of retinoid from the inner to outer segment of salamander cones can be studied by restricting access with a suction electrode (Jin et al., 1994). Here, we used a similar approach to determine the cone compartment where oxidation of *cis* retinol takes place. We found that when the cone outer segment was treated with 9-*cis* retinal, dark adaptation began essentially immediately (Fig. 2B). When the cone inner segment was treated with 9-*cis* retinal, the recovery onset was delayed by ~1 min (Fig. 1C) reflecting the time required for the movement of 9-*cis* retinal to the outer segment before pigment regeneration can begin. The kinetics




Fig. 4. Microspectrophotometric measurement of pigment regeneration in a truncated cone outer segment and in an intact rod. (A) Absorbance spectrum of a dark-adapted cone outer segment. Fitted to the data is a visual pigment template with peak wavelength at 605 nm and an optical density of 0.039. (B) Absorbance spectrum of the same cone outer segment as in (A) recorded after >99% of the visual pigment had been bleached. (C) Absorbance spectrum of the same bleached cone outer segment as in (B) recorded after treatment with 9-*cis* retinol. Fitted to the data is a visual pigment template with peak wavelength at 549 nm and an optical density of 0.040. (D) Absorbance spectrum of the same regenerated cone outer segment as in (C) recorded after a second >99% bleach of its visual pigment. (E) Absorbance spectrum of the outer segment of an intact dark-adapted red rod. Fitted to the data is a visual pigment template with peak wavelength at 518 nm and an optical density of 0.14. (F) Absorbance spectrum of the same rod as in (E) recorded after >99% of the visual pigment had been bleached. (G) Absorbance spectrum of the same bleached rod as in (F) recorded after treatment with 9-*cis* retinol. (H) Absorbance spectrum of the same bleached rod as in (G) recorded after treatment with 11-*cis* retinal. Fitted to the data is a visual pigment template with peak wavelength at 500 nm and an optical density of 0.12.

of the subsequent dark adaptation was identical in the two configurations (Fig. 3A). Notably, after its onset, the kinetics of cone recovery with 9-*cis* retinol in the two configurations were also identical (Fig. 3B) but ~3-fold slower than the recovery kinetics with 9-*cis* retinal. This result suggests that the oxidation of *cis* retinol proceeds relatively slowly and might be rate-limiting for the regeneration of pigment *via* the retina visual cycle. On the other hand, after exposing the inner segment of bleached cones to 9-*cis* retinol, sensitivity recovery onset was delayed by ~6 min. The absence of such delay in the outer segment-out configuration (Fig. 2B) allowed us to rule out slow 9-*cis* retinol oxidation as its source. Instead, this result suggested that this delay likely reflects the time required for 9-*cis* retinol to travel from the inner to the outer segments of cones, similar to the case of 9-*cis* retinal. Thus, the most likely explanation for our electrophysiological results is that 9-*cis* retinol is oxidized in the cone outer segment.

Notably, the delay in recovery onset observed in outer segment-in cones with 9-*cis* retinol (Fig. 1C) was similar to the delay observed in intact salamander retina [Fig. 5A in (Wang et al., 2009)]. Thus, even though isomeric differences between endogenous 11-*cis* retinoids and their exogenous 9-*cis* analogs used in this study could exist, they do not appear to affect dramatically cone dark adaptation. Together, our results suggest that in the intact eye, *cis* retinol is transported from the microvilli of Müller cells to cone outer segments *via* the inner segments and is then oxidized to the *cis* retinal needed for pigment regeneration. The substantially longer time for 9-*cis* retinol delivery from the inner to the outer

segment compared to 9-*cis* retinal (Fig. 1C) suggests the presence of an efficient mechanism for transporting *cis* retinal from cone inner segment to the outer segment. Alternatively, the movement of *cis* retinol through the cone inner segment could somehow be obstructed. Photoreceptors are thought not to have retinoid binding proteins that regulate the transport of hydrophobic retinoids within the cytosol. Indeed, cellular retinal binding protein (Collery et al., 2008) and cellular retinol binding protein (Bok et al., 1984) have not been detected from photoreceptors. Thus, chromophore-binding proteins are not likely to be involved in the regulation of retinoid transport through the cone inner segment. Finally, it is also worth pointing out that due to their different chemical nature, retinol is less hydrophobic than retinal, and would therefore be expected to partition and diffuse differently through the cellular membranes and cytosol of cone photoreceptors.

cis Retinol oxidase in cone outer segment

Our MSP data (Fig. 4) clearly demonstrate the presence of *cis* retinol oxidase in the outer segments of cones. This result is consistent with the hypothesis that RDH8 is the enzyme responsible for this reaction. RDH8 is expressed in cone outer segments (Maeda et al., 2005; Miyazono et al., 2008) and shows weak but detectable activity to *cis* retinoid (Rattner et al., 2000). In addition, deletion of RDH8 blocks the sensitivity recovery with exogenous 9-*cis* retinol in mouse M-cones (Kolesnikov et al., 2015). However, RDH8 is expressed not only in cones but also in rods (Maeda et al., 2005;

Miyazono et al., 2008), a distribution inconsistent with the cone-specific oxidation of *cis* retinol. One possibility is that RDH8 is not itself the *cis* retinol dehydrogenase, but rather it supports this reaction by providing NADP⁺ through the reduction of all-*trans* retinal (Kaylor et al., 2014). Further studies will be needed to determine the exact involvement of RDH8 in the retina visual cycle.

In contrast to our finding from salamander, a recent biochemical study reported *cis*-retinol oxidation activity in the carp cone inner segments but not in the outer segments (Sato et al., 2013; Sato et al., 2015). RDH13L, the enzyme responsible for this inner segment activity, is localized in the inner segments of carp cones but it is not found in the amphibian genome. Notably, a functional homologue of RDH13L, RDH14, is expressed in rod and cone outer segments in bovine retina (Haeseleer et al., 2002). However, the functional role of RDH14 in the retina visual cycle, and specifically in the oxidation of 11-*cis* retinol, is still unknown. The conflicting results obtained from salamander cones (this study) and from carp cones (Sato et al., 2013; Sato et al., 2015) may simply reflect interspecies difference. Alternatively, co-factors or other reaction modifiers could be required for the efficient oxidation of *cis* retinol in cones. In that case, their availability or absence in different experimental conditions could determine the subcellular location of this reaction. Similarly, the presence or absence of cellular compartmentalization in different experimental conditions could also affect the efficiency of *cis* retinol oxidation. Nevertheless, the robust pigment regeneration and dark adaptation observed in our experiments with cone outer segment-delivered 9-*cis* retinol clearly demonstrate the presence of *cis* retinol dehydrogenase activity in the outer segments of cones but not rods.

Finally, it is also possible that another enzyme, different from RDH8 and RDH14, is responsible for the oxidation of *cis* retinol in the outer segments of cones. Notably, RPE65, the isomerohydrolase of the canonical RPE visual cycle, was recently shown to be expressed in the outer segments of cones in some mouse strains (Tang et al., 2011b) as well as in humans (Tang et al., 2011a). It has been suggested that this protein might be involved in the cone visual cycle by acting as a retinol-binding protein (Tang et al., 2013). An interesting possibility is that RPE65 might interact with the cone-specific RDH and, in that way, facilitate the oxidation of *cis* retinol in cones. However, the functional role of RPE65 in cones is still unclear and its possible involvement in the retina visual cycle has not been explored.

Co-factor driving the *cis* retinol oxidation in cones

Previous studies with amphibian rods (Kawamura & Murakami, 1989) and fish rods and cones (Takemoto et al., 2009) both suggest that the inside of the truncated outer segment remains accessible to bath-applied compounds. Thus, the truncated outer segments in our experiments also likely remained open, allowing wash out of small molecules from the cytoplasm. NADP⁺ is thought to be a cofactor in the cone-specific *cis* retinol oxidation (Mata et al., 2002). However, we observed 9-*cis* retinol-driven pigment regeneration in the truncated cone outer segment (Fig. 4) even though NADP⁺ was expected to be washed out rapidly. Thus, the cofactor for *cis* retinol oxidation in the truncated cone outer segment is not likely to be NADP⁺. A possible alternative co-factor is all-*trans* retinal, which is insoluble and shown to work as a co-factor in the *cis* retinol oxidation reaction by carp RDH13L and mouse RDH14 (Sato et al., 2015). All-*trans* retinal is released at a high concentration (up to 3 mM) in

cone outer segments following the bleaching of their visual pigments (Harosi, 1975). In the intact cones, all-*trans* retinal is then rapidly converted to all-*trans* retinol by NADPH-driven reduction (Ala-Laurila et al., 2006). However, our attempt to detect acceleration of *cis* retinol oxidation by all-*trans* retinal was not successful: addition of all-*trans* retinal to the 9-*cis* retinol treatment of either outer segment-in (Fig. 1A) or outer segment-out (Fig. 2A) bleached cones did not affect the kinetics of cone dark adaptation (data not shown). Thus, the cofactor for *cis*-retinol oxidation is still unclear and further experiments will be needed to clarify the molecular mechanism of this key reaction of the retina visual cycle. Our finding that the oxidation of *cis* retinol takes place in the outer segments of cones should inform future studies and help in the eventual identification of the enzyme responsible for this reaction.

Author contribution

SS and VJK designed the study and wrote the paper. SS and VJK designed, performed, and analyzed the experiments shown in Figs. 1, 2, and 3. RF and MCC designed, performed, and analyzed the experiments shown in Fig. 4. All authors reviewed the results and approved the final version of the manuscript.

Acknowledgments

This work was supported by National Institutes of Health (NIH) grants EY019312 and EY025696 (V.J.K.), EY001157 (M.C.C.), EY002687 to the Department of Ophthalmology and Visual Sciences at Washington University, and by Research to Prevent Blindness. We are grateful to the National Eye Institute and Rosalie Crouch from the Medical University of South Carolina for the 11-*cis* retinal.

References

- ALA-LAURILA, P., CORNWALL, M.C., CROUCH, R.K. & KONO, M. (2009). The action of 11-*cis*-retinol on cone opsins and intact cone photoreceptors. *Journal of Biological Chemistry* **284**, 16492–16500.
- ALA-LAURILA, P., KOLESNIKOV, A.V., CROUCH, R.K., TSINA, E., SHUKOLYUKOV, S.A., GOVARDOVSKII, V.I., KOUTALOS, Y., WIGGERT, B., ESTEVEZ, M.E. & CORNWALL, M.C. (2006). Visual cycle: Dependence of retinol production and removal on photoproduct decay and cell morphology. *Journal of General Physiology* **128**, 153–169.
- BOK, D., ONG, D.E. & CHYTL, F. (1984). Immunocytochemical localization of cellular retinol binding protein in the rat retina. *Investigative Ophthalmology & Visual Science* **25**, 877–883.
- COLLERY, R., MCLOUGHLIN, S., VENDRELL, V., FINNEGAN, J., CRABB, J.W., SAARI, J.C. & KENNEDY, B.N. (2008). Duplication and divergence of zebrafish CRALBP genes uncovers novel role for RPE- and Muller-CRALBP in cone vision. *Investigative Ophthalmology & Visual Science* **49**, 3812–3820.
- EBREY, T. & KOUTALOS, Y. (2001). Vertebrate photoreceptors. *Progress in Retinal and Eye Research* **20**, 49–94.
- FREDERIKSEN, R., BOYER, N.P., NICKLE, B., CHAKRABARTI, K.S., KOUTALOS, Y., CROUCH, R.K., OPRIAN, D. & CORNWALL, M.C. (2012). Low aqueous solubility of 11-*cis*-retinal limits the rate of pigment formation and dark adaptation in salamander rods. *Journal of General Physiology* **139**, 493–505.
- GARLIPP, M.A. & GONZALEZ-FERNANDEZ, F. (2013). Cone outer segment and Muller microvilli pericellular matrices provide binding domains for interphotoreceptor retinoid-binding protein (IRBP). *Exp. Eye Res.* **113**, 192–202.
- HAESELEER, F., JANG, G.F., IMANISHI, Y., DRIESSEN, C.A., MATSUMURA, M., NELSON, P.S. & PALCZEWSKI, K. (2002). Dual-substrate specificity short chain retinol dehydrogenases from the vertebrate retina. *Journal of Biological Chemistry* **277**, 45537–45546.
- HAROSI, F.I. (1975). Absorption spectra and linear dichroism of some amphibian photoreceptors. *Journal of General Physiology* **66**, 357–382.

- IMAMOTO, Y. & SHICHIDA, Y. (2014). Cone visual pigments. *Biochimica et Biophysica Acta* **1837**, 664–673.
- INSINNA, C., DANIELE, L.L., DAVIS, J.A., LARSEN, D.D., KUEMME, C., WANG, J., NIKONOV, S.S., KNOX, B.E. & PUGH, E.N., Jr. (2012). An S-opsin knock-in mouse (F81Y) reveals a role for the native ligand 11-*cis*-retinal in cone opsin biosynthesis. *Journal of Neuroscience* **32**, 8094–8104.
- JIN, J., JONES, G.J. & CORNWALL, M.C. (1994). Movement of retinal along cone and rod photoreceptors. *Visual Neuroscience* **11**, 389–399.
- JONES, G.J., CROUCH, R.K., WIGGERT, B., CORNWALL, M.C. & CHADER, G.J. (1989). Retinoid requirements for recovery of sensitivity after visual-pigment bleaching in isolated photoreceptors. *Proceedings of the National Academy of Sciences of the United States of America* **86**, 9606–9610.
- JONES, G.J., FEIN, A., MACNICHOL, E.F., Jr. & CORNWALL, M.C. (1993). Visual pigment bleaching in isolated salamander retinal cones. Microspectrophotometry and light adaptation. *Journal of General Physiology* **102**, 483–502.
- KANAN, Y., KASUS-JACOBI, A., MOISEYEV, G., SAWYER, K., MA, J.X. & AL-UBAIDI, M.R. (2008). Retinoid processing in cone and Muller cell lines. *Experimental Eye Research* **86**, 344–354.
- KAWAMURA, S. & MURAKAMI, M. (1989). Regulation of cGMP levels by guanylate cyclase in truncated frog rod outer segments. *Journal of General Physiology* **94**, 649–668.
- KAYLOR, J.J., COOK, J.D., MAKSHANOFF, J., BISCHOFF, N., YONG, J. & TRAVIS, G.H. (2014). Identification of the 11-*cis*-specific retinyl-ester synthase in retinal Muller cells as multifunctional *O*-acyltransferase (MFAT). *Proceedings of the National Academy of Sciences of the United States of America* **111**, 7302–7307.
- KAYLOR, J.J., YUAN, Q., COOK, J., SARFARE, S., MAKSHANOFF, J., MIU, A., KIM, A., KIM, P., HABIB, S., ROYBAL, C.N., XU, T., NUSINOWITZ, S. & TRAVIS, G.H. (2013). Identification of DES1 as a vitamin A isomerase in Muller glial cells of the retina. *Nature Chemical Biology* **9**, 30–36.
- KEFALOV, V.J., CROUCH, R.K. & CORNWALL, M.C. (2001). Role of noncovalent binding of 11-*cis*-retinal to opsin in dark adaptation of rod and cone photoreceptors. *Neuron* **29**, 749–755.
- KEFALOV, V.J., ESTEVEZ, M.E., KONO, M., GOLETZ, P.W., CROUCH, R.K., CORNWALL, M.C. & YAU, K.W. (2005). Breaking the covalent bond—A pigment property that contributes to desensitization in cones. *Neuron* **46**, 879–890.
- KOLESNIKOV, A.V., MAEDA, A., TANG, P.H., IMANISHI, Y., PALCZEWSKI, K. & KEFALOV, V.J. (2015). Retinol dehydrogenase 8 and ATP-binding cassette transporter 4 modulate dark adaptation of M-cones in mammalian retina. *Journal of Physiology* **593**, 4923–4941.
- KOLESNIKOV, A.V., TANG, P.H., PARKER, R.O., CROUCH, R.K. & KEFALOV, V.J. (2011). The mammalian cone visual cycle promotes rapid M/L-cone pigment regeneration independently of the interphotoreceptor retinoid-binding protein. *Journal of Neuroscience* **31**, 7900–7909.
- KOUTALOS, Y. & CORNWALL, M.C. (2010). Microfluorometric measurement of the formation of all-*trans*-retinol in the outer segments of single isolated vertebrate photoreceptors. *Methods in Molecular Biology* **652**, 129–147.
- MAEDA, A., MAEDA, T., IMANISHI, Y., KUKSA, V., ALEKSEEV, A., BRONSON, J.D., ZHANG, H., ZHU, L., SUN, W., SAPERSTEIN, D.A., RIEKE, F., BAEHR, W. & PALCZEWSKI, K. (2005). Role of photoreceptor-specific retinol dehydrogenase in the retinoid cycle in vivo. *Journal of Biological Chemistry* **280**, 18822–18832.
- MAKINO, C.L. & DODD, R.L. (1996). Multiple visual pigments in a photoreceptor of the salamander retina. *Journal of General Physiology* **108**, 27–34.
- MAKINO, C.L., GROESBEEK, M., LUGTENBURG, J. & BAYLOR, D.A. (1999). Spectral tuning in salamander visual pigments studied with dihydroretinal chromophores. *Biophysical Journal* **77**, 1024–1035.
- MATA, N.L., RADU, R.A., CLEMMONS, R.C. & TRAVIS, G.H. (2002). Isomerization and oxidation of vitamin A in cone-dominant retinas: A novel pathway for visual-pigment regeneration in daylight. *Neuron* **36**, 69–80.
- MIYAZONO, S., SHIMAUCHI-MATSUKAWA, Y., TACHIBANAKI, S. & KAWAMURA, S. (2008). Highly efficient retinal metabolism in cones. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 16051–16056.
- RATTNER, A., SMALLWOOD, P.M. & NATHANS, J. (2000). Identification and characterization of all-*trans*-retinol dehydrogenase from photoreceptor outer segments, the visual cycle enzyme that reduces all-*trans*-retinal to all-*trans*-retinol. *Journal of Biological Chemistry* **275**, 11034–11043.
- SAARI, J.C. (2012). Vitamin A metabolism in rod and cone visual cycles. *Annual Review of Nutrition* **32**, 125–145.
- SARANTIS, M. & MOBBS, P. (1992). The spatial relationship between Muller cell processes and the photoreceptor output synapse. *Brain Research* **584**, 299–304.
- SATO, S., FUKAGAWA, T., TACHIBANAKI, S., YAMANO, Y., WADA, A. & KAWAMURA, S. (2013). Substrate specificity and subcellular localization of the aldehyde-alcohol redox-coupling reaction in carp cones. *Journal of Biological Chemistry* **288**, 36589–36597.
- SATO, S. & KEFALOV, V.J. (2016). *cis* Retinol oxidation regulates photoreceptor access to the retina visual cycle and cone pigment regeneration. *Journal of Physiology* **594**, 6753–6765.
- SATO, S., MIYAZONO, S., TACHIBANAKI, S. & KAWAMURA, S. (2015). RDH13L, an enzyme responsible for the aldehyde-alcohol redox coupling reaction (AL-OL coupling reaction) to supply 11-*cis* retinal in the carp cone retinoid cycle. *Journal of Biological Chemistry* **290**, 2983–2992.
- SHERRY, D.M., BUI, D.D. & DEGRIP, W.J. (1998). Identification and distribution of photoreceptor subtypes in the neonetic tiger salamander retina. *Visual Neuroscience* **15**, 1175–1187.
- TAKEMOTO, N., TACHIBANAKI, S. & KAWAMURA, S. (2009). High cGMP synthetic activity in carp cones. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 11788–11793.
- TANG, P.H., BUHUSI, M.C., MA, J.X. & CROUCH, R.K. (2011a). RPE65 is present in human green/red cones and promotes photopigment regeneration in an in vitro cone cell model. *Journal of Neuroscience* **31**, 18618–18626.
- TANG, P.H., KONO, M., KOUTALOS, Y., ABLONCZY, Z. & CROUCH, R.K. (2013). New insights into retinoid metabolism and cycling within the retina. *Progress in Retinal and Eye Research* **32**, 48–63.
- TANG, P.H., WHELESS, L. & CROUCH, R.K. (2011b). Regeneration of photopigment is enhanced in mouse cone photoreceptors expressing RPE65 protein. *Journal of Neuroscience* **31**, 10403–10411.
- WANG, J.S., ESTEVEZ, M.E., CORNWALL, M.C. & KEFALOV, V.J. (2009). Intra-retinal visual cycle required for rapid and complete cone dark adaptation. *Nature Neuroscience* **12**, 295–302.
- WANG, J.S. & KEFALOV, V.J. (2011). The cone-specific visual cycle. *Progress in Retinal and Eye Research* **30**, 115–128.
- WANG, J.S., NYMARK, S., FREDERIKSEN, R., ESTEVEZ, M.E., SHEN, S.Q., CORBO, J.C., CORNWALL, M.C. & KEFALOV, V.J. (2014). Chromophore supply rate-limits mammalian photoreceptor dark adaptation. *Journal of Neuroscience* **34**, 11212–11221.
- WRIGHT, C.B., REDMOND, T.M. & NICKERSON, J.M. (2015). A history of the classical visual cycle. *Progress in Molecular Biology and Translational Science* **134**, 433–448.
- XUE, Y., SHEN, S.Q., JUI, J., RUPP, A.C., BYRNE, L.C., HATTAR, S., FLANNERY, J.G., CORBO, J.C. & KEFALOV, V.J. (2015). CRALBP supports the mammalian retinal visual cycle and cone vision. *Journal of Clinical Investigation* **125**, 727–738.