

2017

Validation of ICD-9-CM diagnosis codes for surgical site infection and noninfectious wound complications after mastectomy

Margaret A. Olsen
Washington University School of Medicine in St. Louis

Kelly E. Ball
Washington University School of Medicine in St. Louis

Katelin B. Nickel
Washington University School of Medicine in St. Louis

Anna E. Wallace
HealthCore, Inc.

Victoria J. Fraser
Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Olsen, Margaret A.; Ball, Kelly E.; Nickel, Katelin B.; Wallace, Anna E.; and Fraser, Victoria J., "Validation of ICD-9-CM diagnosis codes for surgical site infection and noninfectious wound complications after mastectomy." *Infection Control and Hospital Epidemiology*. 38, 3. 334-339. (2017).
https://digitalcommons.wustl.edu/open_access_pubs/5745

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

ORIGINAL ARTICLE

Validation of ICD-9-CM Diagnosis Codes for Surgical Site Infection and Noninfectious Wound Complications After Mastectomy

Margaret A. Olsen, PhD, MPH;^{1,2} Kelly E. Ball, BSN, MPH, RN;¹ Katelin B. Nickel, MPH;¹ Anna E. Wallace, MPH;³ Victoria J. Fraser, MD¹

BACKGROUND. Few studies have validated ICD-9-CM diagnosis codes for surgical site infection (SSI), and none have validated coding for noninfectious wound complications after mastectomy.

OBJECTIVES. To determine the accuracy of *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) diagnosis codes in health insurer claims data to identify SSI and noninfectious wound complications, including hematoma, seroma, fat and tissue necrosis, and dehiscence, after mastectomy.

METHODS. We reviewed medical records for 275 randomly selected women who were coded in the claims data for mastectomy with or without immediate breast reconstruction and had an ICD-9-CM diagnosis code for a wound complication within 180 days after surgery. We calculated the positive predictive value (PPV) to evaluate the accuracy of diagnosis codes in identifying specific wound complications and the PPV to determine the accuracy of coding for the breast surgical procedure.

RESULTS. The PPV for SSI was 57.5%, or 68.9% if cellulitis-alone was considered an SSI, while the PPV for cellulitis was 82.2%. The PPVs of individual noninfectious wound complications ranged from 47.8% for fat necrosis to 94.9% for seroma and 96.6% for hematoma. The PPVs for mastectomy, implant, and autologous flap reconstruction were uniformly high (97.5%–99.2%).

CONCLUSIONS. Our results suggest that claims data can be used to compare rates of infectious and noninfectious wound complications after mastectomy across facilities, even though PPVs vary by specific type of postoperative complication. The accuracy of coding was highest for cellulitis, hematoma, and seroma, and a composite group of noninfectious complications (fat necrosis, tissue necrosis, or dehiscence).

Infect Control Hosp Epidemiol 2017;38:334–339

Health claims data are useful for retrospective surveillance of postoperative complications and for tracking complications across institutions and the spectrum of care. The accuracy of identification depends on the validity of the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) diagnosis codes chosen for surveillance. Results have been mixed regarding the accuracy of ICD-9-CM diagnosis codes for surgical site infection (SSI) after various procedures. Few studies have validated ICD-9-CM diagnosis codes for SSI after breast procedures,^{1–3} and to our knowledge, no group has validated the coding of noninfectious wound complications (eg, hematoma, fat necrosis) after mastectomy.

We previously reported on the incidence of SSI⁴ and noninfectious wound complications⁵ after mastectomy using data from private insurer claims. We used this population to determine the positive predictive value (PPV) of ICD-9-CM diagnosis codes to identify infectious and noninfectious wound complications and ICD-9-CM procedure and CPT-4

codes to identify breast surgical procedures in claims data compared to medical record review.

METHODS

From the HealthCore Integrated Research Database, a private insurer claims database, we established a retrospective cohort of women 18–64 years of age that had an ICD-9-CM procedure code or a CPT-4 code for mastectomy between January 1, 2004, and August 31, 2009, as described previously (see Supplemental Table 1).⁴ Newly coded SSIs, cellulitis, hematoma, seroma, dehiscence, fat necrosis, and tissue necrosis were identified by ICD-9-CM diagnosis codes in inpatient and outpatient facility and provider claims from 2 to 180 days after mastectomy (day 0 for hematoma), as described previously.^{4,5} The complication algorithm included diagnosis codes specific to breast complications (eg, 611.3 for fat necrosis), and general

Affiliations: 1. Division of Infectious Diseases, Department of Medicine, Washington University School of Medicine, St. Louis, Missouri; 2. Division of Public Health Sciences, Department of Surgery, Washington University School of Medicine, St. Louis, Missouri; 3. HealthCore, Inc., Wilmington, Delaware.

Received July 26, 2016; accepted October 17, 2016; electronically published December 15, 2016

© 2016 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2017/3803-0012. DOI: 10.1017/ice.2016.271

postoperative complications (eg, 998.59 for SSI) (Supplemental Tables 2 and 3).

We randomly selected a subset of women with infectious or noninfectious complications for review of medical records. We prioritized medical records from hospitals for patients with an inpatient claim coded for SSI and/or noninfectious wound complication. If there were no hospitalizations coded with a complication, we prioritized records from outpatient facilities, and finally, we considered medical records from providers. The medical records were obtained by a third-party vendor and were redacted of identifying information before review.

We received records that contained clinical information spanning the time period of the complication of interest for 132 women coded for SSI and/or cellulitis and 188 women coded in the claims data for ≥ 1 noninfectious wound complication. Signs and symptoms of postoperative complications documented by clinicians were abstracted from the medical records by a reviewer blinded to the codes in the claims data (K.E.B.). The Centers for Disease Control and Prevention/National Healthcare Safety Network (NHSN) definition was used to define SSIs in the medical record; the pre-2010 (including cellulitis as an SSI) and 2010 definitions (excluding cellulitis-only) were considered separately.^{6,7}

PPV was calculated as the number of cases confirmed by medical record review divided by the number identified by the claims algorithm. The 95% confidence intervals were calculated using a finite population correction factor. Data management and analyses were performed using SAS v9.3 (SAS Institute, Cary, NC) and SPSS version 21.0 (IBM

SPSS Statistics, Armonk, NY). This study was approved with a waiver of informed consent by the Washington University Human Research Protection Office and Quorum Review.

RESULTS

We obtained medical records of 275 women coded in the claims data for mastectomy; 260 records (94.5%) included information regarding whether mastectomy was performed. The records showed that 2 women had undergone breast-conserving surgery rather than mastectomy, yielding a PPV for mastectomy codes of 99.2%. The PPV for laterality of the mastectomy was 94.6%; 244 of 258 were coded correctly in the claims data as unilateral or bilateral according to medical record review.

In total, 259 records contained sufficient information to determine whether breast reconstruction was performed and the type of reconstruction. Of 122 women coded in the claims for breast expander or implant reconstruction, 119 were confirmed by medical record review (PPV 97.5%). The records of 48 women included codes for autologous flap reconstruction, and 47 of these cases were confirmed by medical record review (PPV, 97.9%).

The PPVs for individual complications are shown in Table 1. The PPV was 57.5% for SSI using the current NHSN definition of SSI (excluding cellulitis-only), but the PPV increased to 68.9% when the pre-2010 NHSN definition including cellulitis-only as SSI was used. The PPVs of individual noninfectious wound complications ranged from 47.8% for fat necrosis to 96.6% for hematoma.

TABLE 1. Positive Predictive Value of Coding in Administrative Claims Data Compared to Medical Record Review to Identify Surgical Site Infection (SSI) and Noninfectious Wound Complications^a After Mastectomy

Coded Complication	No. of Women Coded for Complication ^b	No. of Women With Confirmed Complication by Medical Record Review	Positive Predictive Value, % (95% Confidence Interval)
SSI and/or cellulitis ^c	132	91	68.9 (61.4–76.4)
SSI (not including cellulitis)	120	69	57.5 (49.1–65.9)
SSI (not including cellulitis or 998.51)	107	67	62.6 (53.9–71.3)
Cellulitis	45	37	82.2 (71.6–92.9)
Hematoma	58	56	96.6 (92.2–100.9)
Seroma	78	74	94.9 (90.2–99.6)
Dehiscence	41	26	63.4 (49.2–77.6)
Fat necrosis	23	11	47.8 (28.4–67.2)
Tissue necrosis	23	15	65.2 (46.5–84.0)
Dehiscence, fat necrosis, or tissue necrosis	76	66	86.8 (79.6–94.1)

^aWomen could have multiple different complications per the claims data. Of the 188 women coded for noninfectious wound complications, 157 were coded in the claims data for 1 noninfectious wound complication, 27 were coded for 2 different noninfectious wound complications, and 4 were coded for 3 different noninfectious wound complications (223 individual noninfectious complications). In total, 45 women were coded in the claims data for both SSI and a noninfectious wound complication.

^bNumber of women coded in the claims data for complication for which the medical record was received with clinical information spanning the time period of the complication.

^cPre-2010 NHSN definition included cellulitis as SSI. The PPV for this definition, excluding 998.51, was 75.0 (95% CI, 67.6–82.4).

TABLE 2. Summary of Complications That Were Not Confirmed and Complications Documented in the Medical Records

False-Positive Complication Based on ICD-9-CM Algorithm (Total False Positive)	No. Not Confirmed (n = 100) ^a	Complication Documented in the Medical Record at the Time of False-Positive Coding in the Claims Data
SSI (n = 51)	14	Cellulitis
	13	Seroma
	5	Hematoma
	12	Noninfectious wound complication (dehiscence/fat or tissue necrosis)
	3	Other infection (port, suture abscess)
	4	Nonspecific complication without specific indication of breast infection (chronic inflammation/mastitis, pain/swelling/fever, neutropenia)
	3	No wound complication (breast revision procedure, drain removal, drainage with no complication)
Cellulitis (n = 8)	3	SSI
	2	Other infection (cellulitis of leg/arm)
	2	Seroma
	1	Nonspecific complication without specific indication of breast infection (pain/swelling)
Hematoma (n = 2)	1	Allergic reaction
	2	SSI
	1	Seroma
Seroma (n = 4)	1	Noninfectious wound complication (dehiscence/fat or tissue necrosis)
	1	SSI
Dehiscence (n = 15)	3	Hematoma
	4	SSI
	2	Cellulitis
	1	Seroma
Fat necrosis (n = 12)	1	Hematoma
	3	Noninfectious wound complication (fat or tissue necrosis)
	2	No wound complication (breast reconstruction/revision, no complication)
	4	SSI
	3	Cellulitis
Tissue necrosis (n = 8)	2	Seroma
	1	Hematoma
	11	Noninfectious wound complication (dehiscence/tissue necrosis)
	1	SSI
	2	Hematoma
	3	Noninfectious wound complication (dehiscence/fat necrosis)
	1	Non-specific complication (poorly healing wound)
	2	No wound complication (drain removal, no complication)

NOTE. SSI, surgical site infection.

^aAn individual record could have multiple complications recorded.

The complications documented in the medical record at the time of false-positive coding for a wound complication are shown in Table 2. Of 100 discrepancies between the claims data and medical records among 90 women, 4 (4.0%) had the correctly coded wound complication at another anatomic site (eg, port infection, cellulitis), and 81 (81.0%) had a different breast wound complication recorded in the medical record. The most common complication noted in the medical records for women with false-positive coding for SSI was cellulitis, followed by seroma and noninfectious wound complications. Of the 13 false-positive SSIs with medical record documentation of seroma, 11 cases had been coded with the ICD-9-CM diagnosis code 998.51 (infected postoperative seroma). The most common error in the coding of noninfectious wound

complications involved incorrect use of a code for a different wound complication. Of these discrepancies, 24 involved codes in the claims data for fat or tissue necrosis or dehiscence with medical record documentation of another of these non-infectious complications (Table 2). In the case of fat necrosis, 11 of the 12 complications miscoded as fat necrosis had medical record documentation of tissue necrosis.

DISCUSSION

The PPVs of our ICD-9-CM diagnosis code algorithms to identify infectious and noninfectious complications after mastectomy were variable but generally showed moderate to good results. The PPVs of coding for SSI ranged from 58% to

69% depending on whether cellulitis-alone was considered to meet the SSI definition. Cellulitis-alone was excluded from the NHSN definition of SSI in 2010,^{6,7} and this change has had a considerable impact on reported infection rates because cellulitis is the most common postoperative breast infection.⁸ To address this impact, we calculated the PPV for SSI using the definition of SSI at the time of the study (2004–2009), as well as the current NHSN definition excluding cellulitis-only. Of the 51 miscoded SSIs (using the strict definition of SSI), 14 cases were recorded as cellulitis in the medical record, and 3 of the 8 cases miscoded with cellulitis were recorded in the medical record as an SSI without clinician documentation of cellulitis.

Most nonconfirmed individual complications coded in the claims were documented in the records as other noninfectious breast wound complications. Some of the miscodings could have been due to misinterpretation via term searches. For example, a common error involving false coding for SSI was due to the use of the diagnosis code for infected seroma (998.51), in women with a documented seroma (without evidence for infection). Similarly, difficulty appeared to exist in discriminating fat necrosis versus tissue necrosis.

These results suggest that care should be used with respect to the inclusion of certain codes in algorithms to identify specific complications (ie, avoid 998.51 for identifying SSIs). Because the majority of errors in coding noninfectious wound complications involved misinterpretation of the specific complication, it may be better to focus on a composite group consisting of fat necrosis, tissue necrosis, or dehiscence, which had a higher PPV. In contrast, the PPVs for seroma and hematoma were very high; therefore, these can be reliably identified as individual complications.

In the literature, validation of claims data for SSI surveillance has yielded mixed results. The sensitivities of algorithms using a comprehensive list of codes to identify SSI in several studies were high at 72%–99%,^{9–11} while the PPVs were lower, ranging from 15% to 51%.^{3,10–16} In contrast, in studies that used a small set of SSI-specific diagnosis codes,^{1,2,17–22} sensitivities varied greatly depending on the surgery and the data source, from 20% for 998.59 from a single academic medical center after general and vascular surgery¹⁸ to 100% after joint arthroplasty (using 998.5X and 996.66) or vascular surgery (using 998.5X and 996.62) using Medicare claims.¹⁷ The PPVs in these studies using SSI-specific codes were generally higher, with 4 of 6 studies reporting PPVs >50% and up to 88%.^{1,2,18–21}

Only 3 studies have assessed the accuracy of coding for SSI codes after breast surgery. We previously reported high sensitivity (88%), specificity (99%), and PPV (88%) for SSI using ICD-9-CM diagnosis codes 611.0, 682.2, 682.3, 996.69, and 998.5X from primarily inpatient billing data after breast surgery at an academic medical center.² In a multicenter 2-phase study, Yokoe et al¹ reported the sensitivity of SSI diagnosis codes 998.5X from inpatient billing data as 50% in phase 1 and 70% in phase 2, with PPVs of 58% and 79%,

respectively. Miner et al³ used claims data from a large health-care system to evaluate a complex algorithm for SSI using a variety of ICD-9-CM diagnosis codes and procedure codes for wound care and culture. The PPV of the SSI algorithm using medical claims from inpatient, outpatient, or emergency department encounters within 60 days after breast surgery was only 18%.³

The choice of codes used to identify SSIs and corresponding sensitivity and PPV depends on the intended goal of surveillance. Our intent was to develop an algorithm to compare complication rates among facilities and over time; thus, our focus was on maximizing the PPV to identify SSI and non-infectious wound complications. Interestingly, Calderwood et al^{12,16} reported low PPVs of a comprehensive list of diagnosis and procedure codes used to identify SSIs in Medicare claims after hip arthroplasty and vascular surgery, but they found similar PPVs between hospitals in the best- and worst-performing deciles in terms of risk-adjusted infection rates. These findings highlight the potential benefit of administrative data for comparing SSI rates between hospitals, even when PPVs are low.

The PPVs for the procedure codes to identify mastectomy and breast implant and autologous flap reconstruction were all >97%, consistent with previous reports.^{2,23,24} As described previously,⁴ before randomly selecting procedures for medical record review, we performed extensive filtering to exclude procedures that were unlikely mastectomy (eg, mastectomy coded only by an assistant, brachytherapy catheters at time of procedure). Thus, the true PPVs of the procedure codes for mastectomy and reconstruction procedures are likely lower than we report. Provided that a comprehensive algorithm is used to identify the appropriate patient population, the high PPVs of procedure codes suggest that accurate denominators can be calculated from claims or billing data to compare SSI rates across different surgical procedures or institutions.

We are aware of some limitations in our study. First, we only reviewed the record from 1 source (facility or provider) per patient, so it is possible that documentation of a complication may have been available in an alternative record. We could only determine the PPV of the ICD-9-CM diagnosis code algorithms and not sensitivity, specificity, or negative predictive values because we did not obtain medical records for women who were not coded for complications. A larger study would be helpful to assess whether our findings for individual noninfectious complications can be replicated.

Using health claims data, we captured complications coded at the time of an inpatient or outpatient hospital visit at the same or different institution than the index mastectomy, as well as complications treated in outpatient clinics. We found moderate PPVs for ICD-9-CM diagnosis codes for fat necrosis, tissue necrosis, and dehiscence, and we found high PPVs for coding of cellulitis, hematoma, and seroma. The PPV was modest for SSI without cellulitis, but it improved when cellulitis-only was considered an infection and when the code for infected seroma was excluded from the algorithm.

Given the relatively high PPVs for noninfectious complications, including noninfectious wound complications along with SSI, may provide a more robust measure of quality of care by which to compare complication rates between facilities. New surveillance algorithms to identify infectious and noninfectious wound complications must be developed using ICD-10 codes in future, but this study can be used as the basis to develop ICD-10 algorithms. Finally, the high PPVs for identifying breast procedures are encouraging because they indicate high accuracy of the denominators used to calculate wound complication rates from claims data. Although imperfect, claims data can be used to screen for possible complications, with subsequent confirmation by medical record review, to improve the efficiency of routine SSI surveillance.

ACKNOWLEDGMENTS

We thank Cherie Hill for database and computer management support.

Financial support: Funding for this work was provided by the National Institutes of Health (NIH) (grant no. 5R01CA149614 to M.A.O.). Additional support was provided by the Centers for Disease Control and Prevention (CDC) Epicenters Program (grant no. U54CK000162 to V.J.F.).

Potential conflicts of interest: M.A.O. reports consultant work with Pfizer, Merck, and Sanofi Pasteur and grant funding through Pfizer, Sanofi Pasteur, and Cubist Pharmaceuticals for work outside the submitted manuscript. A.E.W. is an employee of HealthCore, a wholly owned subsidiary of Anthem, Inc., a health insurance company. She has received Anthem stock options and participates in an Anthem employee stock purchase plan. V.J.F. reports that her spouse is Senior Vice President and Chief Medical Officer at Express Scripts. K.E.B. and K.B.N. report no conflicts of interest.

Address correspondence to Margaret A. Olsen, PhD, MPH, Professor of Medicine and Surgery, Washington University School of Medicine, Campus Box 8051, 660 S. Euclid Ave., St. Louis, MO 63110 (molsen@wustl.edu).

SUPPLEMENTARY MATERIAL

To view supplementary material for this article, please visit <https://doi.org/10.1017/ice.2016.271>

REFERENCES

1. Yokoe DS, Noskin GA, Cunningham SM, et al. Enhanced identification of postoperative infections among inpatients. *Emerg Inf Dis* 2004;10:1924–1930.
2. Olsen MA, Fraser VJ. Use of diagnosis codes and/or wound culture results for surveillance of surgical site infection after mastectomy and breast reconstruction. *Infect Control Hosp Epidemiol* 2010;31:544–547.
3. Miner AL, Sands KE, Yokoe DS, et al. Enhanced identification of postoperative infections among outpatients. *Emerg Inf Dis* 2004;10:1931–1937.
4. Olsen MA, Nickel KB, Fox IK, et al. Incidence of surgical site infection following mastectomy with and without immediate reconstruction using private insurer claims data. *Infect Control Hosp Epidemiol* 2015;36:907–914.
5. Nickel KB, Fox IK, Margenthaler JA, Wallace AE, Fraser VJ, Olsen MA. Effect of noninfectious wound complications after mastectomy on subsequent surgical procedures and early implant loss. *J Am Coll Surg* 2016;222:844–852.
6. National Healthcare Safety Network (NHSN) Procedure-Associated (PA) module: surgical site infection (SSI) event. Centers for Disease Control and Prevention website. <http://www.cdc.gov/nhsn/PDFs/pscManual/9pscSSICurrent.pdf>. Updated 2016. Accessed November 14, 2013.
7. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1999;20:250–278.
8. Degnim AC, Throckmorton AD, Boostrom SY, et al. Surgical site infection after breast surgery: impact of 2010 CDC reporting guidelines. *Ann Surg Oncol* 2012;19:4099–4103.
9. Sands KE, Yokoe DS, Hooper DC, et al. Detection of postoperative surgical-site infections: comparison of health plan-based surveillance with hospital-based programs. *Infect Control Hosp Epidemiol* 2003;24:741–743.
10. Rhee C, Huang SS, Berrios-Torres SI, et al. Surgical site infection surveillance following ambulatory surgery. *Infect Control Hosp Epidemiol* 2015;36:225–228.
11. Calderwood MS, Kleinman K, Murphy MV, Platt R, Huang SS. Improving public reporting and data validation for complex surgical site infections after coronary artery bypass graft surgery and hip arthroplasty. *Open Forum Infect Dis* 2014;1:ofu106.
12. Calderwood MS, Kleinman K, Bratzler DW, et al. Use of Medicare claims to identify US hospitals with a high rate of surgical site infection after hip arthroplasty. *Infect Control Hosp Epidemiol* 2013;34:31–39.
13. Stevenson KB, Khan Y, Dickman J, et al. Administrative coding data, compared with CDC/NHSN criteria, are poor indicators of health care-associated infections. *Am J Infect Control* 2008;36:155–164.
14. West J, Khan Y, Murray DM, Stevenson KB. Assessing specific secondary ICD-9-CM codes as potential predictors of surgical site infections. *Am J Infect Control* 2010;38:701–705.
15. Julian KG, Brumbach AM, Chicora MK, et al. First year of mandatory reporting of healthcare-associated infections, Pennsylvania: an infection control-chart abstractor collaboration. *Infect Control Hosp Epidemiol* 2006;27:926–930.
16. Calderwood MS, Kleinman K, Bratzler DW, et al. Medicare claims can be used to identify US hospitals with higher rates of surgical site infection following vascular surgery. *Med Care* 2014;52:918–925.
17. Calderwood MS, Ma A, Khan YM, et al. Use of Medicare diagnosis and procedure codes to improve detection of surgical site infections following hip arthroplasty, knee arthroplasty, and vascular surgery. *Infect Control Hosp Epidemiol* 2012;33:40–49.
18. Hollenbeak CS, Boltz MM, Nikkel LE, Schaefer E, Ortenzi G, Dillon PW. Electronic measures of surgical site infection: implications for estimating risks and costs. *Infect Control Hosp Epidemiol* 2011;32:784–790.
19. Lawson EH, Louie R, Zingmond DS, et al. A comparison of clinical registry versus administrative claims data for reporting of 30-day surgical complications. *Ann Surg* 2012;256:973–981.
20. Bolon MK, Hooper D, Stevenson KB, et al. Improved surveillance for surgical site infections after orthopedic implantation procedures: extending applications for automated data. *Clin Infect Dis* 2009;48:1223–1229.

21. Yokoe DS, Khan Y, Olsen MA, et al. Enhanced surgical site infection surveillance following hysterectomy, vascular, and colorectal surgery. *Infect Control Hosp Epidemiol* 2012;33:768–773.
22. Knepper BC, Young H, Jenkins TC, Price CS. Time-saving impact of an algorithm to identify potential surgical site infections. *Infect Control Hosp Epidemiol* 2013;34:1094–1098.
23. Fisher ES, Whaley FS, Krushat WM, et al. The accuracy of Medicare's hospital claims data: progress has been made, but problems remain. *Am J Public Health* 1992;82:243–248.
24. Talia J, Agarwal S, Momoh AO, Wilkins EG, Kozlow JH. The validity of hospital discharge data for autologous breast reconstruction research. *Plast Reconstr Surg* 2015;135:368–374.