

2017

## Transmesenteric robot-assisted pyeloplasty for ureteropelvic junction obstruction in horseshoe kidney

Aaron M. Potretzke  
*Washington University School of Medicine in St. Louis*

Anand Mohapatra  
*Washington University School of Medicine in St. Louis*

Jeffrey A. Larson  
*Washington University School of Medicine in St. Louis*

Brian M. Benway  
*Cedars-Sinai Medical Center*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

**Please let us know how this document benefits you.**

---

### Recommended Citation

Potretzke, Aaron M.; Mohapatra, Anand; Larson, Jeffrey A.; and Benway, Brian M., "Transmesenteric robot-assisted pyeloplasty for ureteropelvic junction obstruction in horseshoe kidney." *International Brazilian Journal of Urology*. 42, 3. 626-627. (2017).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/5966](https://digitalcommons.wustl.edu/open_access_pubs/5966)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).



# Transmesenteric robot-assisted pyeloplasty for ureteropelvic junction obstruction in horseshoe kidney

Aaron M. Potretzke <sup>1</sup>, Anand Mohapatra <sup>1</sup>, Jeffrey A. Larson <sup>1</sup>, Brian M. Benway <sup>2</sup>

<sup>1</sup> Washington University School of Medicine, Division of Urologic Surgery, St. Louis, MO, USA; <sup>2</sup> Urology Academic Practice, Cedars-Sinai Medical Center, Los Angeles, CA, USA

## INTRODUCTION

Transmesenteric laparoscopic pyeloplasty has previously been reported in two pediatric patients. To our knowledge, there has yet to be a report of a robotic-assisted transmesenteric pyeloplasty. We sought to present a video display and step-by-step demonstration of this approach.

## CASE

The patient is a 28-year old female with an obstructed ureteropelvic junction (UPJ) of the left moiety of a horseshoe kidney. The Da Vinci S robotic platform was used. After transperitoneal access was obtained, a window in the mesentery was identified, and the renal pelvis was exposed. A dismembered, spatulated pyeloplasty was performed with transposition of the UPJ to a more dependent portion of the renal pelvis. A ureteral stent and closed surgical drain were placed.

## RESULTS

The case was completed without complications. Operative time was 207 minutes. Foley

catheter and drain were removed on postoperative days 1 and 2, respectively. The patient was discharged home on postoperative day 2, and the ureteral stent was removed after 8 weeks. Her diuretic T1/2 improved from 39 to 16 minutes. The differential function of the left moiety improved from 31% to 42%.

## CONCLUSIONS

Robot-assisted transmesenteric pyeloplasty is safe and feasible for the management of UPJ obstruction in select patients with a horseshoe kidney. Future study of its use in a larger number of patients is necessary to define its role in this unique population.

## CONFLICT OF INTEREST

None declared.

## ARTICLE INFO

Available at: [www.int brazjurol.com.br/video-section/potretzke\\_626\\_627/](http://www.int brazjurol.com.br/video-section/potretzke_626_627/)

Int Braz J Urol. 2016; 42 (Video #5): 626-7

---

Submitted for publication:  
June 06, 2015

---

Accepted after revision:  
July 17, 2015

---

**Correspondence address:**

Aaron M. Potretzke, MD  
Division of Urology, Washington  
University School of Medicine  
4960 Children's Place  
St. Louis, MO, 63110, USA  
Fax: +1 314 367-5016  
E-mail: potretzkea@wudosis.wustl.edu

**EDITORIAL COMMENT**

This is a well-done video that adds to the literature in the fledgling field of minimally-invasive reconstructive urology. This video should encourage urologists to counsel their patients that this procedure can be performed safely using minimally-invasive techniques with good outcomes. The difficulty with laparoscopic and/or robotic ap-

proaches to upper ureteric strictures is when direct visualization shows that the fibrosis extends beyond what was thought preoperatively. In this situation, the surgeon is faced with the choice of trying to reposition the patient for minimally-invasive psoas hitch/Boari flap, opening, or trying to obtain more length of ureter or bridge a gap that may cause undue tension. While the outcome here is excellent, proper patient selection is key.

*Lucas Wiegand, MD*  
*Assistant Professor*  
*University of South Florida Morsani College of Medicine*  
*Tampa, FL, USA*  
*E-mail: lwiegand@health.usf.edu*