

2017

Assessment of healthcare worker protocol deviations and self-contamination during personal protective equipment donning and doffing

Jennie H. Kwon
Washington University School of Medicine in St. Louis

Carey-Ann D. Burnham
Washington University School of Medicine in St. Louis

Kimberly A. Reske
Washington University School of Medicine in St. Louis

Stephen Y. Liang
Washington University School of Medicine in St. Louis

Tiffany Hink
Washington University School of Medicine in St. Louis

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Kwon, Jennie H.; Burnham, Carey-Ann D.; Reske, Kimberly A.; Liang, Stephen Y.; Hink, Tiffany; Wallace, Meghan A.; Shupe, Angela; Seiler, Sondra; Cass, Candice; Fraser, Victoria J.; and Dubberke, Erik R., "Assessment of healthcare worker protocol deviations and self-contamination during personal protective equipment donning and doffing." *Infection Control & Hospital Epidemiology*. 38, 9. 1077-1083. (2017). https://digitalcommons.wustl.edu/open_access_pubs/6103

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Jennie H. Kwon, Carey-Ann D. Burnham, Kimberly A. Reske, Stephen Y. Liang, Tiffany Hink, Meghan A. Wallace, Angela Shupe, Sondra Seiler, Candice Cass, Victoria J. Fraser, and Erik R. Dubberke

ORIGINAL ARTICLE

Assessment of Healthcare Worker Protocol Deviations and Self-Contamination During Personal Protective Equipment Donning and Doffing

Jennie H. Kwon, DO, MSCI;¹ Carey-Ann D. Burnham, PhD;² Kimberly A. Reske, MPH;¹ Stephen Y. Liang, MD, MPHS;¹ Tiffany Hink, BS;¹ Meghan A. Wallace, BS;² Angela Shupe, BS;² Sondra Seiler, BA;¹ Candice Cass, AA;¹ Victoria J. Fraser, MD;¹ Erik R. Dubberke, MD, MSPH¹ for the CDC Prevention Epicenters

OBJECTIVE. To evaluate healthcare worker (HCW) risk of self-contamination when donning and doffing personal protective equipment (PPE) using fluorescence and MS2 bacteriophage.

DESIGN. Prospective pilot study.

SETTING. Tertiary-care hospital.

PARTICIPANTS. A total of 36 HCWs were included in this study: 18 donned/doffed contact precaution (CP) PPE and 18 donned/doffed Ebola virus disease (EVD) PPE.

INTERVENTIONS. HCWs donned PPE according to standard protocols. Fluorescent liquid and MS2 bacteriophage were applied to HCWs. HCWs then doffed their PPE. After doffing, HCWs were scanned for fluorescence and swabbed for MS2. MS2 detection was performed using reverse transcriptase PCR. The donning and doffing processes were videotaped, and protocol deviations were recorded.

RESULTS. Overall, 27% of EVD PPE HCWs and 50% of CP PPE HCWs made ≥ 1 protocol deviation while donning, and 100% of EVD PPE HCWs and 67% of CP PPE HCWs made ≥ 1 protocol deviation while doffing ($P = .02$). The median number of doffing protocol deviations among EVD PPE HCWs was 4, versus 1 among CP PPE HCWs. Also, 15 EVD PPE protocol deviations were committed by doffing assistants and/or trained observers. Fluorescence was detected on 8 EVD PPE HCWs (44%) and 5 CP PPE HCWs (28%), most commonly on hands. MS2 was recovered from 2 EVD PPE HCWs (11%) and 3 CP PPE HCWs (17%).

CONCLUSIONS. Protocol deviations were common during both EVD and CP PPE doffing, and some deviations during EVD PPE doffing were committed by the HCW doffing assistant and/or the trained observer. Self-contamination was common. PPE donning/doffing are complex and deserve additional study.

Infect Control Hosp Epidemiol 2017;38:1077–1083

Personal protective equipment (PPE) is used in healthcare settings to protect healthcare workers (HCWs) from exposure to pathogens and to prevent the spread of pathogens to other patients. Proper use of PPE is crucial when HCWs care for patients with highly pathogenic organisms, such as the Ebola virus. To date, studies on PPE effectiveness are uncommon, small, and potentially out of date, and many evaluate PPE types no longer in use.¹ The 2014–2016 Ebola virus disease (EVD) outbreak revealed the need for better empirical data regarding best practices to safely don and doff PPE.^{1–5}

Although EVD is a high-visibility, high-impact disease, HCWs are much more likely to encounter pathogens on a daily basis, including methicillin-resistant *Staphylococcus aureus* (MRSA),

vancomycin-resistant *Enterococcus*, *Clostridium difficile*, and carbapenem-resistant *Enterobacteriaceae*. The primary forms of PPE used to protect HCWs and other patients against these important hospital-associated pathogens are associated with contact precautions (CP), which includes gown and gloves. Few data exist regarding whether HCWs follow guidelines for donning and doffing CP PPE and their risk of self-contamination.^{6,7}

One of the primary challenges when designing studies to evaluate PPE or donning/doffing procedures is determining how to model pathogen transmission. The most commonly used surrogate marker for the presence of pathogens is fluorescence, and it can be delivered in a variety of forms: powder,

Affiliations: 1. Division of Infectious Diseases, Washington University School of Medicine, St Louis, Missouri; 2. Department of Pathology and Immunology, Washington University School of Medicine, St Louis, Missouri.

Received January 31, 2017; accepted May 14, 2017; electronically published June 13, 2017

© 2017 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2017/3809-0011. DOI: 10.1017/ice.2017.121

liquid, lotion, etc.^{7–14} Fluorescent markers are inexpensive, easy to use, and because the read out is visual, they can provide immediate feedback to HCWs; however, fluorescence may not be an appropriate surrogate for contamination with infectious viral particles.¹⁰ An alternate marker for viral infection is the MS2 bacteriophage, a single-strand RNA bacteriophage that is a bio-safety level 1 agent and nonpathogenic to humans. MS2 has been used previously in several studies of PPE transmission and/or disinfection,^{10,11,15,16} in a long-term-care facility,¹⁷ a hotel,¹⁸ and an office.¹⁹ Commercial preparations of MS2 are expensive and require significant laboratory expertise to use, but they may provide a more accurate surrogate marker for how pathogens spread in the environment than fluorescence.^{3,10}

The purpose of this study was to evaluate HCW risk of self-contamination when donning and doffing EVD PPE and CP PPE using MS2 bacteriophage and a fluorescent marker as surrogates for pathogen transmission. The frequencies and types of protocol deviations that occurred were documented, and associations between HCW self-contamination after doffing and particular doffing protocol deviations or HCW characteristics were determined.

METHODS

This prospective pilot study was performed at Barnes-Jewish Hospital (BJH), a 1,250-bed, tertiary-care hospital in St Louis, Missouri. The study was approved by the Washington University Human Research Protection Office, and all participants provided written, informed consent. During the study period, EVD PPE consisted of inner and outer gloves (Esteem XP, Cardinal Health, Dublin OH), boot covers (Convertors FullGuard High Top Shoe Covers, Cardinal Health), impervious gown with Velcro on the back of the neck (Convertors SmartSleeve, Cardinal Health), a powered air-purifying respirator (PAPR) and hood with face shield (Versaflo, 3M, Maplewood, MN), and an outer apron (Tyvek apron, Uline, Pleasant Prairie, WI). CP PPE consisted of gloves and a gown (Cardinal Health).

HCW Characteristics

Two sets of HCWs were enrolled in this study. EVD PPE HCWs were enrolled during EVD PPE practice sessions and included respiratory therapists, nurses, infection control preventionists, and critical care physicians. CP PPE HCWs were recruited from BJH hospital wards during their normal shifts and included nurses, patient care technicians, and physicians. HCWs were interviewed regarding demographics and years of service, previous PPE training; HCW height and weight were measured and body mass index (BMI) was calculated.

PPE Donning and Doffing and Contamination Procedures

During EVD PPE training sessions, the donning and doffing processes were aided by a donning/doffing assistant and a

trained observer who instructed HCWs step-by-step as per CDC guidelines.²⁰ HCWs using CP PPE were not given donning or doffing instructions; they were encouraged to proceed according to their usual practices.

After consent, participants were scanned for baseline fluorescence using an UV-A light. Any areas of fluorescence detected were cleaned and noted. Next, HCWs were instructed to don the PPE. Upon completion, HCWs were instructed to close their eyes, and the MS2 bacteriophage and fluorescent marker were applied to HCW palms, abdomens, and ankles (for EVD PPE HCWs) or palms and abdomens (for CP PPE HCWs). Dummy applications of molecular grade water were applied to HCW shoulders. After donning, EVD PPE HCWs practiced various EVD patient care activities before doffing. CP PPE HCWs proceeded directly to doffing. The order and technique used to don and doff the PPE were videotaped and recorded. Immediately after doffing, the participant was scanned for fluorescence. Any areas of fluorescence detected were photographed and sampled utilizing a flocked swab in universal transport medium (Quidel, San Diego, CA). HCW hands (1 swab for both hands), coat sleeves or wrist, and peri-orbital/nasal/oral areas were swabbed regardless of fluorescence.

Donning and doffing videos were reviewed and protocol deviations were recorded. A second reviewer randomly reviewed selected videos to ensure accuracy. Protocol deviations were grouped into categories based on site and the donning/doffing procedural step during which they occurred (ie, glove removal and hand hygiene; PAPR and hood removal). Proper CP PPE and EVD PPE removal sequence were based on recommendations from the CDC^{6,20} and on written protocols used by the BJH infection prevention team.

A commercially available preparation of MS2 (Zeptomatrix, Buffalo, NY), supplied as a stock solution of 1.0×10^9 PFU/mL, was utilized as a surrogate for viral transmission. This substance was diluted to a 1:10 solution in viral transport medium for a working solution of 1.0×10^8 PFU/mL. GloGerm Mist liquid was selected as the fluorescent marker (GloGerm, Moab, UT). A mixture of 100 μ L GloGerm Mist liquid with 0.5 mL working solution MS2 was applied to each contamination site. This combination was tested, and there was no negative effect on MS2 recovery and detection. The mixture of GloGerm liquid and MS2 was drawn into a 3-mL syringe with a needleless, Luer-lock tip (Becton Dickinson, Franklin Lakes, NJ). The syringe was attached to a pediatric intranasal laryngeal mask airway mucosal atomization device (LMA MAD Nasal, Teleflex, Westmeath, Ireland). Syringes were not reused.

MS2 RNA was extracted utilizing a QIAamp viral RNA mini kit (Qiagen, Valencia, CA). MS2 detection was performed using reverse-transcriptase polymerase chain reaction (PCR) using previously described primers²¹ and the Cepheid Smart Cycler with QuantiTect Probe RT-PCR Kit (Qiagen, Valencia, CA). A positive control with MS2 RNA, and a negative control of PCR water was included in each run. The cycle threshold for all positive results was recorded.

Statistical Analyses

The primary outcomes of interest were the presence and frequency of MS2 and/or fluorescent contamination on the HCW after removal of PPE. The secondary endpoints were the correlations of the presence of contamination with the number of lapses in PPE doffing techniques, years of experience, type of PPE, and BMI. Univariate analyses were performed, and $P \leq .05$ was considered significant. We used χ^2 or univariate logistic regression for categorical variables, and we used the Mann-Whitney test for continuous variables. Analyses were performed with SPSS version 24 (IBM, Armonk, NY).

RESULTS

In total, 36 HCWs were enrolled in the study: 18 with EVD PPE and 18 with CP PPE. Most HCWs were nurses: 78% of EVD PPE HCWs and 61% of CP PPE HCWs (Table 1). EVD PPE HCWs were significantly older than CP PPE HCWs (median age, 38 vs 28.5 years; $P = .02$), and there was a trend toward greater years of service among the EVD PPE HCWs (median years of service, 8.5 vs 5.25; $P = .10$).

Donning and Doffing Protocol Deviations

Donning videos were available for review for 15 EVD PPE HCWs (Table 2). Donning videos for the remaining 3 HCWs were unavailable because the HCW was donning simultaneously while another HCW was being recorded. Overall, 27% of EVD PPE HCWs made at least 1 donning protocol deviation, compared with 50% of CP PPE HCWs ($P = .28$). Protocol deviations occurred most often in the gloves and hand-hygiene steps (20% of EVD PPE HCWs and 33% of CP PPE HCWs).

All EVD PPE HCWs had at least 1 doffing protocol deviation, versus 67% of CP PPE HCWs ($P = .02$) (Table 2).

The median number of doffing protocol deviations was greater among EVD PPE HCWs (median, 4 vs 1 among CP PPE HCWs). Moreover, 15 protocol deviations during EVD PPE doffing were committed by the doffing assistant or trained observer: 6 during gown or apron removal, 2 involving hand hygiene, 2 during hood removal, 2 during boot-cover removal, 1 during PAPR removal, and 2 miscellaneous deviations. Among EVD PPE HCWs, the unique doffing step with the greatest number of protocol deviations was boot-cover removal: 78% of HCWs made at least 1 protocol deviation doffing boot covers. The doffing step category with the greatest number of HCWs that committed at least 1 protocol deviation (in both PPE types) was gown/apron removal (83% of EVD PPE HCWs; 50% of CP PPE HCWs), followed by glove removal/hand hygiene (67% of EVD PPE HCWs; 39% of CP PPE HCWs).

MS2 and Fluorescence

Overall, fluorescence was detected on 8 EVD PPE HCWs (44%) and 5 CP PPE HCWs (28%) ($P = .49$), and 21 unique HCW sites fluoresced. The most common site of fluorescence was HCW hands: 6 among EVD PPE HCWs and 5 among CP PPE HCWs (Table 3). Of the 125 samples tested for MS2, 5 were positive (4%). MS2 was recovered from 2 EVD PPE HCWs (11%) and 3 CP PPE HCWs (17%). The 2 EVD PPE HCW sites from which MS2 was recovered were from an alcohol foam pump in the doffing area and an HCW's hands (Table 3). The 3 CP PPE HCW sites from which MS2 was recovered were from the face of 1 HCW and from the sleeves/wrist of 2 HCWs. Among the 5 sites positive for MS2, 2 (40%) also fluoresced. The association between fluorescence and doffing protocol deviations is given in Table 4. There were no significant differences in detection of any fluorescence by protocol deviation type, although there was a trend toward

TABLE 1. Healthcare Worker Demographics

Characteristic	Ebola PPE HCW (n = 18), No. (%) ^a	Contact Precautions PPE HCW (n = 18), No. (%) ^a	P
Age, median (range)	38 (27–55)	28.5 (24–61)	.02
Female	15 (83)	15 (83)	1.00
Years of service, median (range)	8.5 (2.5–30)	5.25 (<1–30)	.10
Previous PPE training	17 (94)	13 (72)	.18
HCW type			
RN, PA, or NP	14 (78)	11 (61)	Ref
MD	2 (11)	2 (11)	.82
Other	2 (11)	5 (28)	.21
Left handed	3 (17)	1 (6)	.60
Body mass index			
Normal	7 (39)	6 (33)	Ref
Overweight	8 (44)	5 (28)	.69
Obese	3 (17)	7 (39)	.26

NOTE. PPE, personal protective equipment; HCW, healthcare worker; RN, registered nurse, PA, physician's assistant; NP, nurse practitioner; MD, medical doctor.

^aUnless otherwise specified.

TABLE 2. Donning and Doffing Protocol Deviations by Personal Protective Equipment (PPE) Type

Characteristic	Ebola PPE HCW, No (%) ^{a,b}	Contact precautions PPE HCW (n = 18), No. (%) ^a
Donning		
Any ^c	4 (27)	9 (50)
Gloves/Hand hygiene	3 (20)	6 (33)
Gown/Apron	3 (20)	4 (22)
PAPR/Hood	2 (13)	N/A
Other	1 (7)	0 (0)
No. of protocol deviations, median (range)	0 (0–4)	0.5 (0–2)
Doffing		
Any ^d	18 (100)	12 (67)
Boot-cover removal	14 (78)	N/A
Gloves/Hand hygiene	12 (67)	7 (39)
Gown/Apron	15 (83)	9 (50)
PAPR/Hood	7 (39)	N/A
Shoe disinfection	8 (44)	N/A
Other	2 (11)	0 (0)
No. of protocol deviations, median (range)	4 (2–8)	1 (0–2)

NOTE. HCW, healthcare worker; N/A, not applicable; PAPR, powered air-purifying respirator.

^aUnless otherwise specified.

^bn = 18 and n = 15 for donning; donning videos were available for 15 of 18 EVD PPE HCWs.

^cP = .28.

^dP = .02.

TABLE 3. Sites of Fluorescence and/or MS2 (N = 24)^a by PPE Type

Site of Fluorescence and/or MS2	EVD PPE No. of Detections	Contact Precautions PPE No. of Detections
Hands	7 ^b	5
Alcohol foam pump	2 ^c	0
Chest	0	2
Forearm	1	0
Knee	1	1
Sleeves/Wrist	1	2 ^d
Thigh	0	1
Face	0	1 ^e

NOTE. PPE, personal protective equipment; EVD, Ebola virus disease; HCW, healthcare worker.

^a6 HCWs had >1 site of fluorescence (none had >1 site of MS2).

^b6 were fluorescent; 1 was MS2 positive.

^c1 fluorescent only; 1 fluorescent and MS2 positive.

^dMS2 positive only.

^eFluorescent and MS2 positive.

significance with boot-cover removal (100% of EVD PPE HCWs with fluorescence detected had a boot cover protocol deviation, versus 60% of EVD PPE HCWs without fluorescence; P = .09).

TABLE 4. Fluorescence and Doffing Protocol Deviations

Doffing Protocol Deviation	Any Fluorescence Detected ^a	No Fluorescence ^b
Any	12 (92)	18 (78)
Boot-cover removal ^c	8 (100)	6 (60)
Gloves/Hand hygiene	8 (62)	11 (48)
Gown/Apron	10 (77)	14 (61)
PAPR/Hood ^a	3 (38)	4 (40)
Shoe disinfection ^a	5 (63)	3 (30)
Other	0 (0)	2 (9)

^an = 13; n = 8 among EVD PPE HCWs.

^bn = 23; n = 10 among EVD PPE HCWs.

^cAmong EVD PPE HCWs only (n = 18).

HCW Characteristics and Donning/Doffing Protocol Deviations

Among EVD PPE HCWs, there was no significant difference in the median number of donning or doffing protocol deviations by years of service (data not shown). There were no significant differences in fluorescence and/or MS2 detection between BMI categories (ie, normal, overweight, or obese; data not shown). There also were no significant differences in the frequencies of types of donning or doffing protocol deviations by BMI (data not shown).

DISCUSSION

Proper use of PPE is essential to protecting patients and HCWs from infectious diseases. However, our results indicate that protocol deviations were common in both donning and doffing. Notably, we found that 100% of EVD PPE HCWs committed at least 1 protocol deviation during doffing, and 27% while donning. This finding is not surprising, given the complexity of EVD PPE, and it is consistent with previous studies.^{8,14,22} In a study involving 120 students, Casalino et al²² found that EVD PPE doffing errors occurred even after a 3-phase training program. While protocol deviations while doffing are a major focus for HCW self-contamination, donning deviations, such as an improperly tied gown (a deviation we observed) may increase the future risk of self-contamination while doffing. Furthermore, we demonstrated that not all protocol deviations were committed by the donning and doffing HCW. For example, several doffing assistants touched the inside of HCW gowns when undoing the neck Velcro, and trained observers occasionally failed to instruct HCWs to perform hand hygiene. While previous studies have evaluated HCW protocol deviations while doffing PPE, few have evaluated the role of other HCWs in the doffing process. This is an important area for future investigation.

Boot-cover removal was particularly problematic. HCWs received varied instructions on the specifics of the boot-cover removal process. HCWs struggled to balance their legs in the air or rest their legs on their scrubs without contaminating themselves, and left-handed HCWs struggled to use right-handed scissors. All 3 left-handed EVD PPE HCWs made ≥1

protocol deviation during boot-cover removal. Many HCWs touched their scrubs with their shoes or gown, both potentially contaminated, during boot-cover removal and shoe disinfection. Herlihey et al²³ also reported difficulties with shoe-cover removal. HCWs caring for EVD patients may be exposed to large amounts of environmental contamination;³ thus, this component of EVD PPE removal may benefit from process improvement. Notably, the recommendations for boot-cover removal have changed since this study was performed. At BJH, the revised process doffs gowns before boot covers, eliminates the use of scissors, and allows HCWs to keep their feet on the ground. We hypothesize that these changes will decrease protocol deviations, but more studies are needed to confirm this.

Hand hygiene and glove removal protocol deviations were common during doffing of both EVD and CP PPE (67% and 39% of HCWs made ≥ 1 error, respectively). During EVD PPE doffing, common protocol deviations included touching outer gloves with inner gloved hands and touching the outside of gloves with bare hands. Herlihey et al²³ reported similar challenges doffing multiple pairs of gloves. Casanova et al¹⁵ compared HCW self-contamination after doffing PPE with single gloves versus double gloves, using MS2 as a marker, and found that although double gloves reduced viral transfer, MS2 was still recovered from the hands of 23% of HCWs after doffing. These results may not be directly comparable to our study because it is unclear whether those HCWs performed hand hygiene after doffing. Regardless, hand hygiene and glove removal are high-risk opportunities for HCW self-contamination. For both the EVD and CP groups, we found fluorescence on HCW hands more often than any other site. HCWs may benefit from targeted training in the correct method for glove removal during EVD PPE doffing, and training should reinforce the fact that gloves are not a substitute for proper hand hygiene.

Measures to reduce HCW self-contamination rates include training and maintenance of training. Several previous studies have suggested that, regardless of PPE type, increased training and access to published donning/doffing guidelines improves HCWs' ability to don and doff PPE without protocol deviation.^{7,12,22,24,25} All EVD PPE HCWs in our study previously had received formal EVD PPE training. By contrast, although 72% of CP PPE HCWs reported having previous training in PPE donning/doffing, this training was often informal, "on the job" training from other HCWs. Similarly, Turnberg et al²⁶ reported that 15%–40% of HCWs had not received PPE training during the previous 12 months, and John et al²⁷ found that "on the job" training was the most common method of PPE training for HCWs. Despite the comparative simplicity of the CP PPE donning/doffing process, only half of the HCWs were able to don PPE without protocol deviation, and only approximately one-third were able to doff PPE without protocol deviation. Common CP PPE doffing protocol deviations included touching the front of the gown with bare hands or allowing the contaminated gown to brush against scrubs while disposing. Possibly, HCWs may be unaware that specific guidelines exist for donning/doffing CP precautions.⁶ Beam et al⁸ demonstrated that simple exposure to a

poster showing the correct donning/doffing sequence may not be enough to improve HCW practices. Tomas et al⁷ found that a training session on CP PPE doffing techniques led to a significant decrease in HCW self-contamination. Formal, targeted interventions or education programs may be needed to improve CP PPE donning/doffing practices.

We were unable to demonstrate clear superiority of either surrogate marker. Fluorescence was detected more frequently than MS2. MS2 was not detected from most sites with fluorescence, and MS2 was detected from 3 sites without fluorescence. Commercial preparations of MS2 are expensive; thus, fluorescent markers, which are inexpensive, may be preferable. Conversely, Casanova et al¹⁰ found considerable MS2 transfer to HCW hands and scrubs in the absence of fluorescence; thus, fluorescence may not accurately mimic transmission of viral particles. MS2 is not visible to the naked eye, and it is possible in our study that additional areas of MS2 contamination were not detected because, outside of HCW hands, face, and arms, we sampled only those areas that fluoresced. Additional data are needed on the relative benefits and limitations of these surrogate markers.

This study has several limitations. It was a relatively small pilot study and as such was underpowered. The small number of HCWs may not be reflective of HCW populations at large. The methods need replication in larger studies, and our methods and results may be useful in designing these. The end of the 2014–2015 EVD outbreak may remove the impetus for healthcare facilities to continue EVD PPE training programs, potentially making future studies of HCWs using EVD PPE more challenging. Some EVD PPE doffing recommendations have been revised since this study was performed. CP PPE, however, are routinely used in healthcare facilities, and larger studies may be possible. We used PCR for MS2 detection; therefore, MS2 detection may not be reflective of viable MS2.

In conclusion, PPE are critical for protecting both HCWs and patients from pathogens, regardless of whether the pathogen in question is high impact like EVD or commonly encountered like *C. difficile*. Previously published data on donning and doffing EVD PPE are limited, both by the number of studies available and the types of data and analyses.^{1,4} There are even fewer data on donning and doffing CP PPE. Our study highlights some potential areas for future research, including an improved boot-cover removal process, improved HCW education in the correct processes for glove removal, and an overall need for better training in the use of CP PPE. Both fluorescent markers and MS2 can be used safely as surrogates for pathogen transmission, although the relative strengths of each need further evaluation. Overall, improved processes for donning/doffing PPE and improved methods for evaluating these processes will help to protect both HCWs and patients from exposure to pathogens.

ACKNOWLEDGMENTS

The authors would like to thank the Barnes-Jewish Hospital Infection Prevention and Interventional Epidemiology Team for their support and assistance with this study.

Financial support: This study was supported by the Centers for Disease Control and Prevention (CDC) Prevention Epi-Center (grant no. 3U54CK000162-05S1). J.H.K. was supported by the Washington University Institute of Clinical and Translational Sciences and a grant (grant no. UL1TR000448, subaward KL2TR000450) from the National Center for Advancing Translational Sciences of the National Institutes of Health (NIH). The content is solely the responsibility of the authors and does not necessarily represent the official view of the NIH. S.Y.L. was supported by the KM1 Comparative Effectiveness Research Career Development Award (grant no. KM1CA156708-01), the Clinical and Translational Science Award program (grant no. UL1RR024992) of the National Center for Advancing Translational Sciences, and the Barnes-Jewish Patient Safety and Quality Career Development Program, which is funded by the Foundation for Barnes-Jewish Hospital.

Potential conflicts of interest: S.Y.L. serves as a subinvestigator for institutional research studies supported by Cepheid. V.J.F. reports that her spouse is Senior Vice President and Chief Medical Officer for Express Scripts; she has current funding from the NIH, the CDC, the Doris Duke Charitable Foundation, and the Foundation for Barnes-Jewish Hospital. She has past funding from the NIH, the CDC, and the AHRQ. All other authors report no conflicts of interest relevant to this article.

Address correspondence to Jennie H. Kwon, DO, MSCI, 4523 Clayton Ave, Campus Box 8051 St Louis, MO 63110 (j.kwon@wustl.edu).

REFERENCES

- Verbeek JH, Ijaz S, Mischke C, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev* 2016;4:Cd011621.
- Fischer WA 2d, Uyeki TM, Tauxe RV. Ebola virus disease: what clinicians in the United States need to know. *Am J Infect Control* 2015;43:788–793.
- Fischer WA 2d, Weber DJ, Wohl DA. Personal protective equipment: protecting health care providers in an Ebola outbreak. *Clin Ther* 2015;37:2402–2410.
- Sprecher AG, Caluwaerts A, Draper M, et al. Personal protective equipment for filovirus epidemics: a call for better evidence. *J Infect Dis* 2015;212:S98–S100.
- Weber DJ, Fischer WA, Wohl DA, Rutala WA. Protecting healthcare personnel from acquiring Ebola virus disease. *Infect Control Hosp Epidemiol* 2015;36:1229–1232.
- Sequence for putting on personal protective equipment (PPE). Centers for Disease Control and Prevention website. <http://www.cdc.gov/hai/pdfs/ppe/ppeposter8511.pdf>. Published 2014. Accessed September 8, 2016.
- Tomas ME, Kundrapu S, Thota P, et al. Contamination of health care personnel during removal of personal protective equipment. *JAMA Intern Med* 2015;175:1904–1910.
- Beam EL, Gibbs SG, Boulter KC, Beckerdite ME, Smith PW. A method for evaluating health care workers' personal protective equipment technique. *Am J Infect Control* 2011;39:415–420.
- Bell T, Smoot J, Patterson J, Smalligan R, Jordan R. Ebola virus disease: the use of fluorescents as markers of contamination for personal protective equipment. *IDCases* 2015;2:27–30.
- Casanova L, Alfano-Sobsey E, Rutala WA, Weber DJ, Sobsey M. Virus transfer from personal protective equipment to healthcare employees' skin and clothing. *Emerging Infect Dis* 2008;14:1291–1293.
- Casanova L, Rutala WA, Weber DJ, Sobsey MD. Methods for the recovery of a model virus from healthcare personal protective equipment. *J Appl Microbiol* 2009;106:1244–1251.
- Guo YP, Li Y, Wong PL. Environment and body contamination: a comparison of two different removal methods in three types of personal protective clothing. *Am J Infect Control* 2014;42:e39–e45.
- Wong TK, Chung JW, Li Y, et al. Effective personal protective clothing for health care workers attending patients with severe acute respiratory syndrome. *Am J Infect Control* 2004;32:90–96.
- Zamora JE, Murdoch J, Simchison B, Day AG. Contamination: a comparison of 2 personal protective systems. *CMAJ* 2006;175:249–254.
- Casanova LM, Rutala WA, Weber DJ, Sobsey MD. Effect of single- versus double-gloving on virus transfer to health care workers' skin and clothing during removal of personal protective equipment. *Am J Infect Control* 2012;40:369–374.
- Tomas ME, Cadnum JL, Jencson A, Donskey CJ. The Ebola disinfection booth: evaluation of an enclosed ultraviolet light booth for disinfection of contaminated personal protective equipment prior to removal. *Infect Control Hosp Epidemiol* 2015;36:1226–1228.
- Sassi HP, Sifuentes LY, Koenig DW, et al. Control of the spread of viruses in a long-term care facility using hygiene protocols. *Am J Infect Control* 2015;43:702–706.
- Sifuentes LY, Koenig DW, Phillips RL, Reynolds KA, Gerba CP. Use of hygiene protocols to control the spread of viruses in a hotel. *Food Environ Virol* 2014;6:175–181.
- Beamer PI, Plotkin KR, Gerba CP, Sifuentes LY, Koenig DW, Reynolds KA. Modeling of human viruses on hands and risk of infection in an office workplace using micro-activity data. *J Occupat Environ Hygiene* 2015;12:266–275.
- Guidance on personal protective equipment (PPE) to be used by healthcare workers during management of patients with confirmed Ebola or persons under investigation (PUIs) for Ebola who are clinically unstable or have bleeding, vomiting, or diarrhea in U.S. hospitals, including procedures for donning and doffing PPE. Centers for Disease Control and Prevention website. <http://www.cdc.gov/vhf/ebola/healthcare-us/ppe/guidance.html>. Published 2015. Accessed September 8, 2016.
- Ninove L, Nougairède A, Gazin C, et al. RNA and DNA bacteriophages as molecular diagnosis controls in clinical virology: a comprehensive study of more than 45,000 routine PCR tests. *PLoS One* 2011;6:e16142.
- Casalino E, Astocondor E, Sanchez JC, Diaz-Santana DE, Del Aguila C, Carrillo JP. Personal protective equipment for the Ebola virus disease: a comparison of 2 training programs. *Am J Infect Control* 2015;43:1281–1287.
- Herlihey TA, Gelmi S, Flewwelling CJ, et al. Personal protective equipment for infectious disease preparedness: a human factors evaluation. *Infect Control Hosp Epidemiol* 2016;37:1022–1028.
- Hon CY, Gamage B, Bryce EA, et al. Personal protective equipment in health care: can online infection control courses transfer knowledge and improve proper selection and use? *Am J Infect Control* 2008;36:e33–e37.
- Weber DJ, Rutala WA, Fischer WA, Kanamori H, Sickbert-Bennett EE. Emerging infectious diseases: focus on infection control issues for novel coronaviruses (Severe Acute Respiratory Syndrome-CoV and Middle East Respiratory Syndrome-CoV),

- hemorrhagic fever viruses (Lassa and Ebola), and highly pathogenic avian influenza viruses, A(H5N1) and A(H7N9). *Am J Infect Control* 2016;44(Suppl 5):e91–e100.
26. Turnberg W, Daniell W, Seixas N, et al. Appraisal of recommended respiratory infection control practices in primary care and emergency department settings. *Am J Infect Control* 2008;36:268–275.
27. John A, Tomas ME, Cadnum JL, et al. Are health care personnel trained in correct use of personal protective equipment? *Am J Infect Control* 2016;44:840–842.