

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2018

Attention-deficit/hyperactivity disorder symptoms are associated with overeating with and without loss of control in youth with overweight/obesity

Amy Heard Egbert
Loyola University Chicago

Denise E. Wilfley
Washington University School of Medicine in St. Louis

Kamryn T. Eddy
Massachusetts General Hospital and Harvard Medical School

Kerri N. Boutelle
University of California - San Diego

Nancy Zucker
Duke University Medical Center

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Egbert, Amy Heard; Wilfley, Denise E.; Eddy, Kamryn T.; Boutelle, Kerri N.; Zucker, Nancy; Peterson, Carol B.; Doyle, Angela Celio; Grange, Daniel Le; and Goldschmidt, Andrea B., "Attention-deficit/hyperactivity disorder symptoms are associated with overeating with and without loss of control in youth with overweight/obesity." *Childhood Obesity*. 14, 1. 50-57. (2018).
https://digitalcommons.wustl.edu/open_access_pubs/6485

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Amy Heard Egbert, Denise E. Wilfley, Kamryn T. Eddy, Kerri N. Boutelle, Nancy Zucker, Carol B. Peterson, Angela Celio Doyle, Daniel Le Grange, and Andrea B. Goldschmidt

Attention-Deficit/Hyperactivity Disorder Symptoms Are Associated with Overeating with and without Loss of Control in Youth with Overweight/Obesity

Amy Heard Egbert, MA,¹ Denise E. Wilfley, PhD,² Kamryn T. Eddy, PhD,³
Kerri N. Boutelle, PhD,⁴ Nancy Zucker, PhD,⁵ Carol B. Peterson, PhD,⁶
Angela Celio Doyle, PhD,⁷ Daniel Le Grange, PhD,^{8,9} and Andrea B. Goldschmidt, PhD¹⁰

Abstract

Background: There is growing evidence that attention-deficit/hyperactivity disorder (ADHD) and loss of control (LOC) eating, both prevalent in children and adolescents, may be related to one another. However, the relationship between ADHD and overeating without LOC has been largely unexamined, thus precluding an understanding of the independent contributions of LOC and episode size in these associations. The current study sought to examine associations between ADHD symptoms and maladaptive eating by evaluating three different types of eating episodes characterized by the presence/absence of LOC and the amount of food consumed: objectively large LOC episodes [objective binge eating (OBE)], subjectively large binge episodes [subjective binge eating (SBE)], and objectively large overeating episodes without LOC [objective overeating (OO)].

Methods: Participants were 385 youth (M age = 10.89, SD = 2.25) drawn from five different research protocols at institutions across the United States. Participants and their parents completed questionnaires and semistructured interviews to assess ADHD symptoms, OBE, SBE, and OO.

Results: As hypothesized, negative binomial regressions revealed that ADHD symptoms were significantly associated with OBE, $\chi^2(1) = 16.61, p < 0.001$, and with OO, $\chi^2(1) = 10.64, p < 0.01$. Contrary to expectations, they were not associated with SBE.

Conclusions: These results indicate the need for future studies to explore possible shared mechanisms (e.g., impulsivity) underlying associations between ADHD symptoms, OBE, and OO. Clinical implications include support for considering ADHD symptoms in programs that target both prevention of LOC eating and obesity more generally.

Keywords: ADHD; adolescents; binge eating; children; loss of control eating

¹Department of Psychology, Loyola University Chicago, Chicago, IL.

²Department of Psychiatry, Washington University School of Medicine, St. Louis, MO.

³Department of Psychiatry, Massachusetts General Hospital and Harvard Medical School, Boston, MA.

⁴Department of Pediatrics and Psychiatry, University of California San Diego, La Jolla, CA.

⁵Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Durham, NC.

⁶Department of Psychiatry, University of Minnesota Medical Center, Minneapolis, MN.

⁷Eating Disorders Center, Evidence Based Treatment Centers of Seattle, Seattle, WA.

⁸Department of Psychiatry, University of California San Francisco, San Francisco, CA.

⁹Department of Psychiatry and Behavioral Neuroscience, The University of Chicago, Chicago, IL.

¹⁰Department of Psychiatry and Human Behavior, Alpert Medical School of Brown University, Weight Control and Diabetes Research Center/The Miriam Hospital, Providence, RI.

Introduction

Attention-deficit/hyperactivity disorder (ADHD) is a prevalent condition affecting 3–10% of school-aged children.^{1,2} ADHD is characterized by two categories of impairment, hyperactivity/impulsivity and inattention, and can be diagnosed in individuals who experience clinically significant symptoms in either one or both domains.³

In addition to the impairments that are characteristic of the disorder, ADHD has been linked to a range of adverse health outcomes, including poor sleep, diabetes, and obesity.⁴ In particular, an increasing number of cross-sectional studies have begun to examine the link between obesity and ADHD and have found a bidirectional relationship between the two conditions in children.⁵ Furthermore, there are now longitudinal studies suggesting that ADHD symptoms in childhood may play a causal role in the development of overweight in adolescence and adulthood.^{6,7}

Despite mounting evidence in support of a link between ADHD and obesity, the mechanisms underlying the association are less understood. ADHD symptoms have been positively linked to multiple dietary factors, such as sugar-sweetened beverage consumption and total caloric intake.^{8,9} Likewise, other mechanisms have been suggested, such as genetics and circadian rhythm disruption.⁵ Additionally, several studies have indicated that dysregulated eating may be a significant contributor to the increased rates of obesity in individuals with ADHD.⁵ More specifically, loss of control (LOC) eating has been identified as a potential mediator of the relationship between ADHD diagnosis and zBMI score in children.¹⁰

LOC eating is characterized by the subjective experience of being unable to control what or how much food one is eating.³ LOC, along with the consumption of an objectively large amount of food in a given period of time (*i.e.*, objective binge eating or OBE), is the central component of binge eating disorder (BED).³

Because it is difficult to classify what is considered an objectively large amount of food for a child, and because LOC eating in children is associated with significant distress and impairment regardless of whether an objectively large amount of food is consumed,¹¹ research in children has focused on the experience of LOC, irrespective of episode size. A recent review of the literature found similar levels of psychosocial impairment (*e.g.*, anxiety, depression, internalizing, and externalizing symptoms) and BMI in youth with OBE compared with youth who experienced LOC without overeating (*i.e.*, subjective binge eating, or SBE).¹²

There is also evidence that LOC may be critically linked to obesity. LOC eating occurs in youth with overweight at greater rates than their normal weight counterparts, the likelihood of reporting LOC eating appears to increase proportionally with weight,¹³ and LOC eating is a strong predictor of subsequent weight gain.¹⁴

In addition to obesity, the few studies that have assessed LOC eating and ADHD have also found associations between the two. Reinblatt et al. found that school-aged children with

an ADHD diagnosis showed a 12-fold increase in the odds of reporting LOC (with no distinction between SBE and OBE).¹⁰ Another study that looked only at OBE found a significant positive association between OBE and ADHD symptoms in young adults.¹⁵ However, few studies have evaluated associations between LOC and ADHD in children with obesity, despite links between obesity and both LOC and ADHD.

Although there is a growing body of literature supporting the importance of LOC in connection with both obesity and ADHD symptoms, fewer studies have assessed the impact of overeating without LOC (objective overeating; OO). Some of the research that does exist shows that OO and OBE may lie on a continuum such that individuals who report OO experience more distress than those who do not experience OO, and individuals who report OBE, in turn, experience more distress than those with OO.¹⁶ Other studies have found that individuals with both OBE and SBE experience more eating pathology (*e.g.*, shape and weight concerns) than those with OO.¹⁷

With regard to zBMI, findings are mixed, with some studies demonstrating that individuals with OO weigh significantly less than those with OBE and others finding no difference.¹² When studied longitudinally, Sonnevile et al. found that OBE was uniquely predictive of weight gain in adolescents and young adults, while OO was not¹⁸; however, this is the only study to date that has examined the predictive validity of OO.¹² The apparent lack of research on OO is problematic given the importance of teasing apart the effects of LOC and episode size to understand whether they are differentially associated with other negative outcomes, such as obesity and ADHD.

When considering ADHD specifically, the literature on OO is also limited. Sonnevile et al. found that ADHD symptoms were associated with overeating during late childhood; however, they assessed overeating by asking children if they had ever eaten so much that they were in pain or made themselves throw up.¹⁹ Therefore, overeating in this context is representative of more than just eating an objectively large amount of food as it also introduces the idea of compensatory behavior. Other studies have found that an ADHD diagnosis is associated with higher daily calorie consumption, but have not evaluated OO specifically.⁸

Although the literature points to the idea that both LOC and overeating may be associated with ADHD symptoms, more research must be conducted to test this hypothesis with well-established definitions and measures of overeating. Furthermore, the studies that have examined associations between LOC and ADHD symptoms have done so by including both OBE and SBE in the definition of LOC¹⁰ or by only evaluating OBE without SBE.¹⁵ However, links between ADHD symptoms and OBE, SBE, and OO should be examined separately to avoid confounding LOC with episode size. Likewise, continuing to tease apart the distress associated with LOC vs. episode size is also important as OBE is currently required to receive an eating disorder diagnosis.³

Finally, little attention has been given to how other demographic factors, such as age and gender, may impact the

relationship between ADHD symptoms and dysregulated eating (*i.e.*, OBE, SBE, or OO). Such factors may be important to consider given that ADHD is more often diagnosed in boys than in girls²⁰ and OBE appears to emerge in adolescence.¹⁸

The current study seeks to expand the literature by evaluating the association between ADHD symptoms and both LOC and eating episode size in youth with overweight. It examines ADHD symptoms in the context of (1) OBE, (2) SBE, and (3) OO to determine whether these conditions are differentially related to ADHD symptoms.

Many studies have examined LOC eating in children meeting criteria for an ADHD diagnosis, but have not evaluated prodromal ADHD symptoms, which have important implications for prevention or early intervention programs. This study utilizes ADHD symptoms in place of a formal ADHD diagnosis to evaluate the association between attention and impulsivity on LOC eating and overeating across a spectrum of severity levels. It also examines the frequency of OBE, SBE, and OO, as opposed to presence or absence, as this appears to be of importance in youth with overweight.²¹ Finally, this study investigates whether age and sex moderate the relationship between ADHD and dysregulated eating.

It is expected that ADHD symptoms will be positively associated with LOC eating episode frequency,

such that OBE and SBE frequency will each be significantly associated with ADHD symptoms. Likewise, we expect that ADHD symptoms will also be positively associated with OO frequency. Finally, we hypothesize that these associations will be stronger for older children than for younger children and stronger for boys than for girls.

Method

Participants

Participants included were 385 youth who were overweight or obese (M age = 10.89, SD = 2.25). The majority of participants were female (63%) and Caucasian (60%). On average, participants had a standardized BMI score of 2.26 (SD = 0.34), which corresponds to the 97th age- and sex-adjusted percentile. See Table 1 for full participant characteristics for both the total sample and by dysregulated eating status.

For the purposes of the current study, dysregulated eating is defined as reporting OBE, SBE, or OO. The current sample was drawn from a larger pool of individuals ($n = 526$) presenting for a number of different protocols at research institutions across the United States (Washington University School of Medicine,^{22,23} The University of

Table 1. Participant Characteristics by Dysregulated Eating Status

	Dysregulated eating ($n = 124$)	No dysregulated eating ($n = 261$)	Total ($n = 385$)	p
Age, n (SD)	11.49 (2.24)	10.60 (2.06)	10.89 (2.25)	<0.001 ^a
BMI z-score, n (SD)	2.33 (0.37)	2.23 (0.33)	2.26 (0.34)	0.01 ^a
Sex, n (%)				0.24 ^b
Male	46 (37)	81 (31)	127 (37)	
Female	78 (63)	180 (69)	258 (63)	
Race, n (%)				<0.001 ^b
White	75 (60)	201 (77)	276 (72)	
African American	43 (35)	42 (16)	85 (22)	
Other	6 (5)	18 (7)	24 (6)	
Ethnicity, n (%)				0.30 ^b
Hispanic	12 (9)	35 (13)	47 (12)	
Non-Hispanic	112 (90)	226 (87)	338 (88)	
ADHD symptoms, n (SD)	56.17 (8.26)	54.42 (6.18)	54.98 (6.59)	<0.05 ^a
OBE, n (SD)	1.07 (2.47)	—	0.35 (1.48)	<0.001 ^a
SBE, n (SD)	1.63 (3.59)	—	0.52 (2.17)	<0.001 ^a
OO, n (SD)	0.99 (2.08)	—	0.32 (1.26)	<0.001 ^a

Dysregulated eating = Report of objective binge eating episode, subjective binge eating episode, or objective overeating episode.

^aMultivariate analysis of variance (MANOVA), including age, zBMI, ADHD symptoms, and dysregulated eating in the model.

^bChi-square test of association.

ADHD, attention-deficit/hyperactivity disorder.

Table 2. Participant Characteristics by Study Site

	Reason for presentation	Demographics	Exclusion criteria
Washington University School of Medicine/ San Diego State University	Online weight loss intervention	Overweight and obese 12–17-year-olds	Past or current diagnosis of an eating disorder, medical conditions impacting weight or physical activity ability, taking medication significantly affecting weight
Washington University School of Medicine	Laboratory-based study examining the relationship between mood and eating	Overweight and obese 6–12-year-old girls	Medications or conditions impacting appetite, weight, or attention; severe restriction or limited preferences of food; current diagnosis of bulimia nervosa
The University of Chicago	Initial assessment at University of Chicago Eating and Weight Disorders Program	Overweight and obese 7–18-year-olds	Meeting DSM-IV criteria for anorexia nervosa or bulimia nervosa
Children's Hospital Boston	Behavioral weight loss treatment at Optimal Weight for Life Clinic	Overweight and obese 8–18-year-olds	Developmental/cognitive disorders, obesity-related disorders associated with intellectual disability, psychotic disorders
San Diego State University/University of California San Diego	Study of weight loss maintenance treatment for children	Overweight and obese 7–12-year-olds	Medical or psychiatric treatment limitations, use of medications affecting appetite or weight, involvement in other weight loss or psychological treatment
University of Minnesota	Treatment study for two interventions to reduce overeating	Overweight and obese 8–12-year-olds	Presence of medical or psychiatric disorder requiring physician supervision of diet and exercise, medication affecting weight or eating, food allergies and diet restrictions, involvement in other weight loss program

Chicago,²⁴ Children's Hospital Boston,²⁵ San Diego State University/University of California San Diego,²⁶ and University of Minnesota²⁷), including nonintervention studies and weight loss treatment studies, as well as for clinic-based treatment of eating disorders or obesity. See Table 2 for detailed inclusion criteria for each site.

In addition to the criteria listed in Table 2, participants were included in the current study if they completed the Child Behavior Checklist (CBCL), the Eating Disorder Examination (EDE), Child or Adult Version, and had recorded height and weight measures.

Procedure

Institutional Review Board approval was obtained for the research protocols at each respective site. Informed consent was provided by each participant's parent/guardian and written assent was obtained from each child before participation in the study. The current study only includes baseline data that were collected before participation in any treatment or experimental study.

Measures

Demographics. zBMI and BMI percentile scores were calculated based on standardized height and weight measurements for age. Race was categorized as Caucasian, African American, or Other, and ethnicity was categorized as Hispanic or Non-Hispanic.

Eating Disorder Examination. Participants completed the EDE or the child EDE (ChEDE).²⁸ The EDE and the ChEDE are semistructured interviews used to diagnose eating disorders and assess related symptomatology. The two interviews are nearly identical, but differences between the ChEDE and the EDE include modifications to simplify language and complex concepts (e.g., overvaluation of shape and weight) for children. Higher scores represent more severe eating disorder psychopathology.

The EDE and ChEDE define OBE as eating episodes characterized by both LOC and consumption of an objectively large amount of food according to clinical rating standards, and SBE as LOC eating episodes in which the respondent considers the amount of food eaten to be excessive, but that amount is not considered objectively large by clinical rating standards. OO is defined as consumption of an objectively large amount of food without LOC. OBE, SBE, and OO frequencies were recorded for the past 28 days. Both the EDE and ChEDE have good reliability and validity.^{28,29}

Child Behavior Checklist/6-18. The CBCL is a 113-item caregiver report measure used to assess problematic behavior in children.³⁰ The CBCL has several behavioral subscales, including an Attention Subscale, as well as a total score that represents global behavioral issues. Respondents report their agreement with how much each item describes their child now or in the past 6 months on the following three-point

scale: 0 [*Not True (as far as you know)*], 1 [*Somewhat or Sometimes True*], and 2 [*Very True or Often True*]. The 10-item Attention Subscale was used for the current study to assess for ADHD symptoms. Items include statements asking how much the child is “Impulsive or acts without thinking” and “Inattentive or easily distracted.” T-scores range from 0 to 100, and scores greater than 65 are considered to be in the clinical range. The CBCL has good reliability and validity,^{30,31} and the Attention Subscale has been found to have excellent discriminating power for detecting ADHD.³² Continuous t-scores were used in all analyses.

Statistical Analyses

All analyses were conducted using SPSS, Version 24.0. Preliminary analyses were conducted using analyses of variance to characterize the sample (Table 1). The primary study analyses were then conducted in two phases, adjusting all models for study site, as well as race, age, gender, and zBMI, as is customary in the literature.¹⁰

A series of negative binomial regression models were used to examine relationships between ADHD symptoms and dysregulated eating. Negative binomial regression was used due to the existence of count data for all dependent variables (OBE, SBE, and OO). Negative binomial regression differs from normal (ordinary least squares) regression, in that it does not allow for negative integers, accounts for a distribution skewed toward zero, as is often the case with count data, and allows for overdispersion of the model (*i.e.*, the variance exceeds the mean).³³ It is a generalized linear model in which predicted scores are expressed as natural logarithms. Before conducting the analysis, all variables were centered to avoid multicollinearity.³⁴

In the first model, OBE frequency was entered as the dependent variable. Study site was entered as a covariate to adjust for any confounding effects. In addition, race, age, gender, and zBMI were also entered into the model as covariates, and ADHD symptoms were entered in as the independent variable of interest. Additionally, the age \times ADHD symptom and

gender \times ADHD symptom moderators were entered. In the second and third models, respectively, SBE and OO frequencies were analyzed separately using the same approach.

Results

Descriptive Information

Means, standard deviations, and correlations are presented in Table 3. Overall, participants had an average CBCL Attention Problems score of 55.05 (SD = 7.01), which fell in the nonclinical range. Eleven percent of participants ($n = 45$) met criteria for clinically significant ADHD symptoms (CBCL scores greater than or equal to 65). As can be seen in Figure 1, about 1/3 of youth ($n = 125$) in our sample endorsed at least one dysregulated eating episode (OBE, SBE, or OO). Of those, 72% reported had at least one LOC episode ($n = 89$). Overlap between OBE, SBE, and OO was also examined (Fig. 1). Eighty percent of youth who had a dysregulated eating episode only endorsed one type. SBE was the most common, followed by OO, and then OBE. The largest amount of overlap was between OBE and SBE, with 11 participants endorsing both. Only 2% of all participants reported having all three forms of dysregulated eating.

ADHD Symptoms and Associations with Dysregulated Eating

ADHD symptoms were then specifically evaluated using negative binomial regression. Full results of the regression analysis are presented in Table 4. First, associations between ADHD symptoms and OBE were evaluated. The likelihood ratio for the full model was $\chi^2(9) = 129.44$, $p < 0.001$, indicating model significance. Adjusting for study site, race, age, gender, and zBMI, Wald chi-square analyses indicated that ADHD symptoms were positively associated with OBE frequency $\chi^2(1) = 16.61$, $p < 0.001$, such that there was a 6% increase in OBE frequency for every one-point increase in ADHD symptoms. Next, SBE frequency was evaluated. The likelihood ratio for the full model was

Table 3. Means and Correlations of Variables of Interest

		1	2	3	4	5	6
1	Age	—					
2	zBMI	0.02	—				
3	OBE episodes (past 28 days)	0.16**	0.10	—			
4	SBE episodes (past 28 days)	0.04	0.02	0.27**	—		
5	OO episodes (past 28 days)	0.09	0.02	0.25**	0.29**	—	
6	ADHD symptoms	0.08	0.03	0.17**	0.04	0.10*	—
	Mean	10.89	2.26	0.35	0.52	0.32	54.98
	Standard deviation	2.24	0.35	1.48	2.17	1.26	6.96
	Range	6.93–17.90	1.11–3.30	0–12	0–30	0–14	50–97

* $p < 0.05$; ** $p < 0.01$.




Figure 1. Venn diagram of prevalence rates of objective binge eating (OBE) episodes, subjective binge eating (SBE) episodes, and objective overeating (OO) episodes. Of 385 total respondents, 124 exhibit *at least one* OBE, SBE, or OO episode. Percentages shown in this diagram are based on $n = 124$ as the divisor.

$\chi^2(9) = 189.88$, $p < 0.001$, which indicated that the overall model was significant. However, Wald chi-square analyses revealed that ADHD symptoms were not significantly associated with SBE frequency $\chi^2(1) = 1.30$, $p = 0.25$.

Finally, OO frequency was evaluated. The likelihood ratio for the full model was $\chi^2(9) = 88.63$, $p < 0.001$, indicating model significance. Wald chi-square analyses indicated that adjusting for study site, race, age, gender, and zBMI, ADHD symptoms were also positively associated with OO frequency $\chi^2(1) = 10.64$, $p < 0.01$, such that there was a 5% increase in OO for every one-point increase in ADHD symptoms. Contrary to hypotheses, age and gender

were not significant moderators of the relationship between ADHD and OBE, SBE, or OO.

Discussion

The current study assessed the relationship between ADHD symptoms and three types of maladaptive eating, LOC with overeating, LOC without overeating, and overeating without LOC, in a sample of youth with overweight. This study sought to examine whether the relationship between ADHD symptoms and eating behavior differed by episode size (objectively or subjectively large) and by the presence/absence of LOC. We found that OBE and OO frequencies were both associated with ADHD symptoms. However, there was no significant association between ADHD symptoms and SBE frequency, indicating that associations between ADHD symptoms and dysregulated eating behavior may be better accounted for by variations in episode size, rather than the presence of LOC.

This study contributes to the literature in several ways. First, it builds upon previous findings that ADHD symptoms are significantly associated with LOC eating in youth.¹⁰ Moreover, this study also builds on the literature by showing that when OBE and SBE are considered separately, ADHD symptoms are only associated with OBE frequency. Although previous research has found few differences in psychopathology between youth with OBE and with SBE,¹² most studies have not examined ADHD symptoms specifically. It is possible that OBE and SBE are differentially related to some forms of psychopathology, but not others. This study also found that in addition to OBE, ADHD symptoms are associated with OO in youth with overweight.

One explanation for our findings may lie in shared mechanisms underlying ADHD symptoms, OBE, and OO. The literature has shown that children with ADHD have more difficulty inhibiting their desire for immediate small rewards in favor of long-term large rewards and that they make more

Table 4. Summary of Three Negative Binomial Regression Analyses for Variables Predicting Objective Binge Eating Episodes, Subjective Binge Eating Episodes, and Objective Overeating ($n = 385$)

Variable	OBE			SBE			OO		
	B	SE B	z	B	SE B	z	B	SE B	z
Race									
White	1.12	0.80	1.41	1.05*	0.49	2.13	2.99**	1.07	2.80
Black	1.40	0.80	1.76	0.73	0.50	1.45	2.26*	1.08	2.10
Age	0.20**	0.06	3.34	0.04	0.05	0.81	0.15*	0.06	2.49
Sex (female = 1)	0.36	0.28	-1.25	-0.48	0.27	1.75	0.62*	0.29	2.14
zBMI	0.87*	0.39	2.23	0.50	0.31	1.60	0.66	0.35	1.87
ADHD symptoms	0.05***	0.01	4.12	0.02	0.02	1.13	0.05**	0.01	3.28

All models adjusted for study site.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

impulsive errors on tasks measuring inhibitory skills.^{35,36} Likewise, problems with impulsivity have also been found in youth with and without overweight who also endorse LOC eating, using both objective and self-report measures.^{10,37,38}

Given these findings one hypothesis is that overeating (with or without LOC) may be a behavioral manifestation of the impulsivity that is also characteristic of ADHD. This is supported by the adult literature, which has found that individuals who experience OBE have significantly more problems with impulse control than those who only experience SBE.³⁹ Likewise, greater levels of impulsivity may explain why some individuals act upon feelings of LOC by overeating, while others can experience those feelings, but not engage in OBE. While no study to our knowledge has evaluated impulsivity in youth who experience OO, it is possible that these individuals also demonstrate diminished levels of inhibitory control. OO and OBE both share a central component—overeating.

While individuals with OO do not endorse losing control over how much they are eating, they may still be eating impulsively due to internal or external cues (e.g., boredom or environmental cues). However, this theory should be tested by empirical research on the mechanisms underlying OO.

In addition to theoretical implications, this study's evaluation of prodromal symptoms of ADHD has important implications for prevention, namely that OBE and OO may be problems even for children who do not meet full criteria for ADHD. While the inclusion of participants with a full spectrum of ADHD symptoms is a strength of the study, it is not possible to distinguish whether including children who were symptomatic for ADHD, but did not meet full criteria for the disorder, might have impacted the findings. It is possible that in children who have a diagnosis of ADHD, symptoms would be more strongly related to OBE than to OO. However, the results of this study show that at subclinical levels of ADHD, children with overweight may be susceptible not only to OBE but also to OO.

This study has several strengths, including a large sample comprising both treatment-seeking and nontreatment-seeking participants. As noted above, an additional strength of this study is the emphasis on prodromal symptoms of ADHD and implications in the context of prevention. Finally, this study used well-validated semistructured interviews to evaluate LOC eating and OO.

Limitations of the study include the cross-sectional nature of the data as well as the lack of diversity of the sample given that all participants were overweight, which precludes generalizing the findings to normal weight youth. The restricted zBMI range of the sample may explain our findings that zBMI was not significantly associated with OBE, SBE, or OO and that it was not correlated with ADHD symptoms. Additionally, ADHD symptoms were identified using a caregiver report questionnaire, without corroborating neuropsychological testing or direct observation. Such questionnaires may be subject to the bias of a single reporter, which could have impacted the results.

Finally, in the current sample, the mean age of participants was just under 11 years. Although (consistent with previous

literature) participants who endorsed dysregulated eating were older than those who did not (Table 1), the mean age was still under 12 years. Thus, the age of the sample may have impacted our findings as research has found that ADHD symptoms in early childhood are associated with overeating in middle childhood, and then finally OBE in adolescence.¹⁹ As such, it is possible that more participants would have endorsed OBE had the sample been older.

In summary, ADHD symptoms appear to be associated with overeating in youth both with and without LOC. However, LOC without overeating does not appear to be linked to ADHD symptoms. These findings suggest that both OBE and OO may be linked to impaired impulsivity/inattention. Future studies should examine potential underlying mechanisms explaining associations between OBE/OO and ADHD symptoms in children and explore possible associations between these forms of dysregulated eating and specific symptoms of ADHD (e.g., hyperactive vs. inattentive symptoms).

Future research should also be conducted to examine the relationship between OO/OBE and ADHD symptoms in a longitudinal manner to assess whether youth with subthreshold ADHD symptoms are more likely to develop BED in adulthood. Finally, the findings of this study demonstrate the importance of conducting longitudinal research on all types of overeating and LOC eating among youth to elucidate the prospective impact and overlap of these different conditions.

Acknowledgments

The authors wish to acknowledge the following funding sources: NIH grants T32-MH082761 (Dr. Le Grange), R01-HD036904 (Dr. Wilfley), K24-MH070446 (Dr. Wilfley), P30-DK50456 (Drs. Peterson, Zucker, and Boutelle), F31-MH071019 (Dr. Eddy), and K23-DK105234 (Dr. Goldschmidt); a National Science Foundation Graduate Research Fellowship (Ms. Egbert); an Academy for Eating Disorders Graduate Student Research Grant (Dr. Goldschmidt); an American Psychological Association's Division 38 Student Research Award (Dr. Celio Doyle); and an RGA/Washington University Longer Life Foundation Research Award (Dr. Wilfley).

Author Disclosure Statement

No competing financial interests exist.

References

1. Polanczyk GV, Salum GA, Sugaya LS, et al. Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *J Child Psychol Psychiatry* 2015;56:345–365.
2. Polanczyk GV, Willcutt EG, Salum GA, et al. ADHD prevalence estimates across three decades: An updated systematic review and meta-regression analysis. *Int J Epidemiol* 2014;43:434–442.
3. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub, Washington, DC, 2013.

4. Nigg J. Attention-deficit/hyperactivity disorder and adverse health outcomes. *Clin Psychol Rev* 2013;33:215–228.
5. Cortese S, Tessari L. Attention-deficit/hyperactivity disorder (ADHD) and obesity: Update 2016. *Curr Psychiatry Rep* 2017;19:4.
6. Aguirre Castaneda RL, Kumar S, et al. Childhood attention-deficit/hyperactivity disorder, sex, and obesity. *Mayo Clin Proc* 2016;91:352–361.
7. White B, Nicholls D, Christie D, et al. Childhood psychological function and obesity risk across the lifecourse: Findings from the 1970 British Cohort Study. *Int J Obes* 2012;36:511–516.
8. van Egmond-Fröhlich AW, Weghuber D, de Zwaan M. Association of symptoms of attention-deficit/hyperactivity disorder with physical activity, media time, and food intake in children and adolescents. *PLoS One* 2012;7:e49781.
9. Ptacek R, Kuzelova H, Stefano GB, et al. Disruptive patterns of eating behaviors and associated lifestyles in males with ADHD. *Med Sci Monit* 2014;20:608–613.
10. Reinblatt SP, Mahone EM, Tanofsky-Kraff M, et al. Pediatric loss of control eating syndrome: Association with attention-deficit/hyperactivity disorder and impulsivity. *Int J Eat Disord* 2015;48:580–588.
11. Morgan CM, Yanovski SZ, Nguyen TT, et al. Loss of control over eating, adiposity, and psychopathology in overweight children. *Int J Eat Disord* 2002;31:430–441.
12. Goldschmidt AB. Are loss of control while eating and overeating valid constructs? A critical review of the literature. *Obes Rev* 2017;18:412–449.
13. Goldschmidt AB, Aspen VP, Sinton MM, et al. Disordered eating attitudes and behaviors in overweight youth. *Obesity* 2008;16:257–264.
14. Tanofsky-Kraff M, Shomaker LB, Olsen C, et al. A prospective study of pediatric loss of control eating and psychological outcomes. *J Abnorm Psychol* 2011;120:108.
15. Steadman KM, Knouse LE. Is the relationship between ADHD symptoms and binge eating mediated by impulsivity? *J Atten Disord* 2016;20:907–912.
16. Ackard DM, Neumark-Sztainer D, Story M, Perry C. Overeating among adolescents: Prevalence and associations with weight-related characteristics and psychological health. *Pediatrics* 2003;111:67–74.
17. Shomaker LB, Tanofsky-Kraff M, Elliott C, et al. Salience of loss of control for pediatric binge episodes: Does size really matter? *Int J Eat Disord* 2010;43:707–716.
18. Sonnevile KR, Horton NJ, Micali N, et al. Longitudinal associations between binge eating and overeating and adverse outcomes among adolescents and young adults: Does loss of control matter? *JAMA Pediatr* 2013;167:149–155.
19. Sonnevile KR, Calzo JP, Horton NJ, et al. Childhood hyperactivity/inattention and eating disturbances predict binge eating in adolescence. *Psychol Med* 2015;45:2511–2520.
20. Froehlich TE, Lanphear BP, Epstein JN, et al. Prevalence, recognition, and treatment of attention-deficit/hyperactivity disorder in a national sample of US children. *Arch Pediatr Adolesc Med* 2007;161:857–864.
21. Glasofer DR, Tanofsky-Kraff M, Eddy KT, et al. Binge eating in overweight treatment-seeking adolescents. *J Pediatr Psychol* 2007;32:95–105.
22. Doyle AC, Goldschmidt A, Huang C, et al. Reduction of overweight and eating disorder symptoms via the internet in adolescents: A randomized controlled trial. *J Adolesc Health* 2008;43:172–179.
23. Goldschmidt AB, Tanofsky-Kraff M, Wilfley DE. A laboratory-based study of mood and binge eating behavior in overweight children. *Eat Behav* 2011;12:37–43.
24. Eddy KT, Celio-Doyle A, Rienecke R, et al. Eating disorder not otherwise specified in adolescents. *J Am Acad Child Adolesc Psychiatry* 2008;47:156–164.
25. Eddy KT, Tanofsky-Kraff M, Thompson-Brenner H, et al. Eating disorder pathology among overweight treatment-seeking youth: Clinical correlates and cross-sectional risk modeling. *Behav Res Ther* 2007;45:2360–2371.
26. Wilfley DE, Stein RI, Saelens BE, et al. Efficacy of maintenance treatment approaches for childhood overweight: A randomized controlled trial. *JAMA* 2007;298:1661–1673.
27. Boutelle KN, Zucker NL, Peterson CB, et al. Two novel treatments to reduce overeating in overweight children: A randomized controlled trial. *J Consult Clin Psychol* 2011;79:759–771.
28. Bryant-Waugh RJ, Cooper PJ, Taylor CL, Lask BD. The use of the eating disorder examination with children: A pilot study. *Int J Eat Disord* 1996;19:391–397.
29. Cooper Z, Cooper PJ, Fairburn CG. The validity of the eating disorder examination and its subscales. *Br J Psychiatry J Ment Sci* 1989;154:807–812.
30. Achenbach TM and Rescorla LA. *Manual for the ASEBA School Age Forms and Profiles*. University of Vermont Research Center for Children, Youth, and Families: Burlington, VT, 2001.
31. Achenbach TM, Elderbrock C. *Manual for the Child Behavior Checklist and Revised Child Behavior Profile*. University of Vermont Department of Psychiatry: Burlington, VT, 1991.
32. Chen WJ, Faraone SV, Biederman J, Tsuang MT. Diagnostic accuracy of the Child Behavior Checklist scales for attention-deficit hyperactivity disorder: A receiver-operating characteristic analysis. *J Consult Clin Psychol* 1994;62:1017–1025.
33. Cox S, West SG, Aiken LS. The analysis of count data: A gentle introduction to Poisson regression and its alternatives. *J Pers Assess* 2009;91:121–136.
34. Holmbeck GN. Post-hoc probing of significant moderational and mediational effects in studies of pediatric populations. *J Pediatr Psychol* 2002;27:87–96.
35. Patros CHG, Alderson RM, Kasper LJ, et al. Choice-impulsivity in children and adolescents with attention-deficit/hyperactivity disorder (ADHD): A meta-analytic review. *Clin Psychol Rev* 2016;43:162–174.
36. Wright L, Lipszyc J, Dupuis A, et al. Response inhibition and psychopathology: A meta-analysis of go/no-go task performance. *J Abnorm Psychol* 2014;123:429–439.
37. Lee-Winn AE, Townsend L, Reinblatt SP, Mendelson T. Associations of neuroticism and impulsivity with binge eating in a nationally representative sample of adolescents in the United States. *Personal Individ Differ* 2016;90:66–72.
38. Govey MA, Lim CS, Dutton GR, et al. Executive function and dysregulated eating behaviors in pediatric obesity. *J Pediatr Psychol*. [Epub ahead of print.]
39. Keel PK, Mayer SA, Harnden-Fischer J. Importance of size in defining binge eating episodes in bulimia nervosa. *Int J Eat Disord* 2001;29:294–301.

Address correspondence to:
 Amy Heard Egbert, MA
 Department of Psychology
 Loyola University Chicago
 1000 W. Sheridan Road
 Chicago, IL 60626
 E-mail: ahead2@luc.edu