

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2017

Limbic system damage in MS: MRI assessment and correlations with clinical testing

Jie Wen

Washington University School of Medicine in St. Louis

Dmitriy A. Yablonskiy

Washington University School of Medicine in St. Louis

Amber Salter

Washington University School of Medicine in St. Louis

Anne H. Cross

Washington University School of Medicine in St. Louis

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Wen, Jie; Yablonskiy, Dmitriy A.; Salter, Amber; and Cross, Anne H., "Limbic system damage in MS: MRI assessment and correlations with clinical testing." PLoS One. 12, 11. e0187915. (2017).

https://digitalcommons.wustl.edu/open_access_pubs/6521

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.


RESEARCH ARTICLE

Limbic system damage in MS: MRI assessment and correlations with clinical testing

Jie Wen¹, Dmitriy A. Yablonskiy¹, Amber Salter², Anne H. Cross^{3*}

1 Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, MO, United States of America, **2** Division of Biostatistics, Washington University School of Medicine, St. Louis, MO, United States of America, **3** Department of Neurology, Washington University School of Medicine, St. Louis, MO, United States of America

* crossa@wustl.edu



OPEN ACCESS

Citation: Wen J, Yablonskiy DA, Salter A, Cross AH (2017) Limbic system damage in MS: MRI assessment and correlations with clinical testing. PLoS ONE 12(11): e0187915. <https://doi.org/10.1371/journal.pone.0187915>

Editor: Stephen D. Ginsberg, Nathan S Kline Institute, UNITED STATES

Received: July 3, 2017

Accepted: October 27, 2017

Published: November 9, 2017

Copyright: © 2017 Wen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was supported by the National Multiple Sclerosis Society (<http://www.nationalmssociety.org/>) (RG4463 to AHC, G-1508-08577 to JW); by the Conrad N Hilton Foundation (<https://www.hiltonfoundation.org/>) (Marilyn Hilton Award for Innovation in MS Research to AHC); the Manny & Rosalyn Rosenthal – Dr. John L. Trotter MS Center Chair in Neuroimmunology of the Foundation for Barnes-Jewish Hospital (<https://www.barnes-jewish.org/>)

Abstract

Volume loss in some limbic region structures has been observed in multiple sclerosis (MS) patients. However, in vivo evaluation of existing tissue cellular microstructure integrity has received less attention. The goal of studies reported here was to quantitatively assess loss of limbic system volumes and tissue integrity, and to evaluate associations of these measures with cognitive and physical dysfunction in MS patients. Thirty-one healthy controls (HC) and 80 MS patients, including 32 relapsing remitting (RRMS), 32 secondary progressive (SPMS) and 16 primary progressive (PPMS), participated in this study. Tissue cellular integrity was evaluated by means of recently introduced tissue-specific parameter $R2t^*$ that was calculated from multi-gradient-echo MRI signals using a recently developed method that separates $R2t^*$ from BOLD (blood oxygen level dependent) contributions to GRE signal decay rate constant ($R2^*$), and accounting for physiological fluctuations and artifacts from background gradients. Volumes in limbic system regions, normalized to skull size (NV), were measured from standard MPRAGE images. MS patients had lower $R2t^*$ and smaller normalized volumes in the hippocampus, amygdala, and several other limbic system regions, compared to HC. Alterations in $R2t^*$ of several limbic system regions correlated with clinical and neurocognitive test scores in MS patients. In contrast, smaller normalized volumes in MS were only correlated with neurocognitive test scores in the hippocampus and amygdala. This study reports the novel finding that $R2t^*$, a measure that estimates tissue integrity, is more sensitive to tissue damage in limbic system structures than is atrophy. $R2t^*$ measurements can serve as a biomarker that is distinct from and complementary to volume measurements.

Introduction

Cognitive impairment, including impairments of memory, attention, information processing and executive functions, affects approximately 50% of multiple sclerosis (MS) patients [1]. Dysfunctions of episodic memory and information processing speed are the most common

www.barnesjewish.org/Giving/About-The-Foundation) (AHC).

Competing interests: Anne Cross has received payments for consulting for Abbvie, Bayer, Biogen, EMD Serono, Genentech, Genzyme, Mallinckrodt, Novartis, and Teva. This does not alter our adherence to all PLOS ONE policies on sharing data and materials.

types of dysfunctions reported [2]. The limbic system contains structures that play important roles in memory functions. These include not only the hippocampus but also the amygdala, which has roles in cognition and memory processing as well as in emotion [3,4]. Previous studies in MS observed atrophy in parts of the limbic system [5,6] which was associated with memory impairment [7]. Abnormalities in the white matter (WM) tracts connecting structures of the limbic system, such as the cingulum, fornix and uncinate fasciculus, have been demonstrated in MS using diffusion tensor imaging, and the abnormalities correlated with cognitive dysfunction [8–10].

We developed a technique called gradient echo plural contrast imaging (GEPCI), which is a post-processing technique that is based on a commercially-available gradient recalled echo (GRE) pulse sequence with multiple gradient echoes [11–13]. GEPCI can provide quantitative measurement of $R2^*$, which is the relaxation rate constant of the transverse magnetization which measures how fast the MR signal decays. We previously analyzed $R2^*$ to evaluate global and regional GM integrity in MS, finding that lower $R2^*$ in cortical GM correlated with worse neurocognitive test scores [14]. $R2^*$ and quantitative susceptibility mapping (QSM) have been found to be sensitive to iron deposition in basal ganglia and were correlated with inhibitory control in MS patients [15]. In studies reported in this paper we used MRI-based methods to measure tissue-specific relaxation properties of GRE MRI signal, i.e. the $R2^*$ ($1/T2^*$) relaxation rate constant and its sub-component, $R2t^*$, specifically related to tissue cellular structure [16,17]. The $R2^*$ was demonstrated to detect WM abnormalities in MS patients, progressively decreasing with myelin loss [18]. Besides tissue-specific contributions, susceptibility related effects, such as extravascular BOLD (blood-oxygen-level dependent) effect which is continually changing in response to neuronal activity [19], also contribute to $R2^*$ signal. Our group introduced a method [16] to separate tissue-specific ($R2t^*$) from BOLD [20] contributions to $R2^*$. The tissue-specific ($R2t^*$) MRI relaxation parameter depends on the cellular environment of water molecules (the main source of MRI signal). Higher concentrations of proteins, lipids, and other components of biological tissue and cellular constituents (sources of MR signal relaxation) lead to higher relaxation rate constants. For example, in pure water or CSF, the $R2t^*$ is about 1 s^{-1} , while in normal brain tissue it is about $15\text{--}20\text{ s}^{-1}$. Using this method [16] and acquisition plus post-processing techniques that minimize artifacts related to macroscopic magnetic field inhomogeneities [21], and physiological fluctuations [22], we previously demonstrated that $R2t^*$ measurements are sensitive to age-related changes in brain tissue cellular structure in healthy adults [17] and loss of tissue cellular integrity in Alzheimer Disease [23]. By the elimination of the variable contribution to $R2^*$ that is dependent on venous blood oxygenation level (BOLD), $R2t^*$ values provide more tissue specific information than $R2^*$ [17,23].

Here we measured $R2t^*$ and volumes of limbic system components—with a focus on hippocampus and amygdala—in 80 MS patients and 31 healthy controls (HC). We observed that not only did MS patients have smaller hippocampus volumes compared to HC, but $R2t^*$ in hippocampus and amygdala and several other limbic regions was also reduced, suggesting damage even in non-atrophied tissue. Moreover, alterations in $R2t^*$ and volume of several different regions in the limbic system of MS patients correlated with clinical and neurocognitive test scores.

Materials and methods

Participants

The study was approved by the Institutional Review Board of Washington University School of Medicine. Eighty MS patients with relapsing remitting (RRMS, $n = 32$), secondary progressive (SPMS, $n = 32$) or primary progressive (PPMS, $n = 16$) clinical courses were enrolled. Ten male and 21 female HC, age 23 to 85 years (mean age \pm SD: 49.53 ± 15.98), were enrolled to

Table 1. Participant characteristics.

		HC	RRMS	PPMS	SPMS
Number of participants	F	17	27	10	20
	M	9	5	6	12
Age	F	53.00±14.54	54.70±8.01	56.40±8.60	57.45±8.71
	M	42.23±17.17	51.80±16.27	53.50±10.43	56.67±10.79
EDSS	F	N/A	2.70±1.34	5.70±1.18	6.03±1.23
	M	N/A	2.00±0.71	5.75±1.04	5.29±1.48
25FTW	F	N/A	5.07±1.63	15.77±26.65	13.88±8.90
	M	N/A	3.87±0.93	12.62±10.66	9.35±5.77
9HPT (Dom)	F	N/A	22.58±7.11	28.82±19.40	28.30±16.59
	M	N/A	23.42±4.90	31.23±7.79	36.03±11.09
9HPT (Non Dom)	F	N/A	22.16±4.39	29.32±15.17	37.10±29.88
	M	N/A	20.96±2.84	47.21±49.91	39.97±19.60
PASAT (3s)	F	N/A	45.89±11.97	47.90±11.12	42.44±14.12
	M	N/A	49.40±6.80	51.00±5.73	42.00±9.81
PASAT (2s)	F	N/A	34.00±10.05	37.10±8.32	32.06±12.34
	M	N/A	38.60±5.37	42.67±9.63	30.33±8.69
SDMT	F	N/A	53.07±11.84	51.20±9.15	43.16±12.48
	M	N/A	55.00±4.53	51.17±14.34	40.75±13.85

Mean ± SD for each parameter was shown in the table. HC = Healthy Control; RRMS = Relapse Remitting Multiple Sclerosis; PPMS = Primary Progressive Multiple Sclerosis; SPMS = Secondary Progressive Multiple Sclerosis; EDSS = Expanded Disability Status Scale; 25FTW = Timed 25-Foot Walk; 9HPT (Dom) = 9-Hole Peg Test of the dominant hand; 9HPT (Non Dom) = 9-Hole Peg Test of the non-dominant hand; PASAT = Paced Auditory Serial Addition Test; SDMT = Symbol Digit Modalities Test.

<https://doi.org/10.1371/journal.pone.0187915.t001>

represent the age range of the MS patients (Table 1). Sixty out of 80 MS patients were on disease-modifying treatments, including beta-interferon (n = 20), dimethyl fumarate (12), fingolimod (4), glatiramer acetate (9), methotrexate (1), natalizumab (4), ocrelizumab (5) and teriflunomide (5). All subjects provided written consent.

MRI

Scans were performed using a 3T Trio MRI scanner (Siemens, Erlangen, Germany) with a 32-channel phased-array head coil. High resolution GEPCI [13] datasets of voxel size 1×1×2 mm³ were acquired using a three dimensional multi-gradient-echo sequence with flip angle of 30°, TR = 50 ms and total acquisition time of 12 min. For each acquisition, 10 echoes were collected with first echo time TE₁ = 4ms and echo spacing ΔTE = 4ms. Navigator echoes were collected to correct physiological fluctuation-induced artifacts [22]. A voxel spread function algorithm [21] was used to correct for artifacts related to macroscopic field inhomogeneity effects. High resolution clinical MPAGE [24] images (1 mm³) were also collected for tissue segmentation.

Image processing

Multi-channel images were combined using a published algorithm [13], and fitted by a model [16] on a voxel-by-voxel basis:

$$S(TE) = S_0 \cdot \exp[-R_2^* \cdot TE] \cdot F_{BOLD}(TE) \cdot F(TE) \quad (1)$$

where S_0 is the signal magnitude. Functions F_{BOLD} and F describe signal decay due to BOLD effect [16] and macroscopic field inhomogeneity [21], respectively. This model and data

acquisition method allow measurements of tissue-specific $R2t^*$ relaxation rate constant free of artifacts related to macroscopic field inhomogeneities and physiological fluctuations.

MPRAGE images were put into “FreeSurfer” (Laboratory for Computational Neuroimaging, Martinos Center for Biomedical Imaging) for brain segmentations, which were registered onto GEPCI maps using “FSL” (FMRIB, Oxford). Median $R2t^*$ values for hippocampus, amygdala, parahippocampal cortex, insula, entorhinal cortex, lateral and medial orbitofrontal cortices, caudal and rostral anterior cingulate cortices, isthmus of cingulate, and posterior cingulate were collected. Medians (rather than means) were used to help offset any segmentation errors. Volumes (V) of each limbic region were normalized to skull size using “Sienax” in “FSL” to derive normalized volume (NV) [22]. MPRAGE and FLAIR images were used to obtain lesion load (LL) of MS patients by using “lesion-TOADs” tool [25] in MIPAV [26].

Clinical neurological tests

The Expanded Disability Status Scale (EDSS), 25 foot timed walk (25FTW), upper extremity function (nine hole peg test - 9HPT), the PASAT, a test of auditory information processing speed and calculation ability, and SDMT, a test of visual processing speed, each validated for MS, were assessed on the day of MRI by examiners blinded to imaging results. Notably, new learning and memory abilities have been previously shown in MS patients to be primarily associated with processing speed [27]. PASAT and SDMT raw scores were converted into z-scores using published control data for age and education.

Statistics

Analyses were done in the statistical program R. ANCOVA for repeated measures with false discovery rate (FDR) correction, with age and gender as covariates, was used to compare $R2t^*$ and NV differences in hippocampus and amygdala between HC and MS patients of different clinical subtypes. $R2t^*$ and NV in limbic structures were examined using Pearson correlations with cognitive test scores (PASAT and SDMT), controlled for age, gender, LL and treatment. Spearman correlations, controlled for age, gender, LL and treatment, were used for correlations between $R2t^*$ and NV in hippocampus and amygdala, and non-normally distributed clinical results (EDSS, 25FTW and 9HPT). For determining correlations of brain imaging with 9HPT tests, only right-handed (73 of 80 patients) patients were included. 9HPT scores of dominant and non-dominant hands were correlated with MRI measurements in the left and right hemispheres, respectively. FDR with Benjamini-Hochberg procedure [28] was used to correct for multiple testing. After correction, $p < 0.05$ was considered significant.

Results

Gender effects on $R2t^*$ and volumes in the limbic system

The limbic system, with focus on the hippocampus and amygdala, was examined using $R2t^*$ and regional volumes (Fig 1). Some prior studies reported gender differences in hippocampus size [29,30] and larger amygdala in men [31]. Our data are in agreement with these studies, however these differences do not exist in HC if volumes are normalized to the skull size.

In HC, we found no significant gender differences in $R2t^*$ of hippocampus, amygdala and other limbic cortical regions. In our MS cohort, there were no significant differences in NV between genders except that normalized hippocampal volume was larger in females with MS (Fig 2). Our data do not support a gender difference in normalized size of amygdala, nor do they indicate any gender differences in $R2t^*$.




Fig 1. Examples of GEPCI S_0 (T1w) images and $R2t^*$ maps selected from a healthy control (HC: 58 yr old, female) and MS patient (MS: 60 yr old, female). Hippocampus and amygdala regions are marked by yellow and green contours, respectively. Despite no dramatic volumetric changes between HC and MS in these two structures, the MS patient showed significant lower $R2t^*$ in the anterior hippocampus region.

<https://doi.org/10.1371/journal.pone.0187915.g001>

Comparisons of $R2t^*$ and NV between MS patients and HC

Reduced NV within regions of the limbic system has previously been reported in MS [5–7]. Consistent with those reports, we observed lower NV in MS than HC in both hippocampus and amygdala (Fig 3). MS patients also had lower $R2t^*$ in the right hippocampus (Fig 3).

MS patients had been categorized as RRMS, SPMS, or PPMS prior to entering the study. We compared $R2t^*$ and NV among the clinical subtypes and compared each subtype with HC. All subtypes had reduced $R2t^*$ compared to HC in the right hippocampus (Fig 4). $R2t^*$ and NV were reduced in SPMS patients in both hippocampus and amygdala compared to HC, except $R2t^*$ in left amygdala. PPMS patients had smaller NV in bilateral hippocampus. $R2t^*$ of right amygdala and NV of bilateral hippocampus were reduced in SPMS compared to RRMS. (Fig 4).

Associations of $R2t^*$ and NV in hippocampus and amygdala with cognitive test performance

$R2t^*$ and NV of limbic regions were examined for associations with performance on cognitive tests. Reduced $R2t^*$ of the right hippocampus and right amygdala each correlated moderately




Fig 2. Gender comparisons for V and NV of hippocampus and amygdala in each hemisphere in HC (A–H) and MS (I–P) groups. Panels with p-values (with FDR correction) ≤ 0.05 and > 0.01 are marked with “*”. Panels with p-values (with FDR correction) ≤ 0.01 are marked with “**”. F = Female; M = Male; LH = Left Hemisphere; RH = Right Hemisphere.

<https://doi.org/10.1371/journal.pone.0187915.g002>

with worse SDMT scores (Fig 5R and 5T). Smaller NV of the hippocampus and amygdala of each hemisphere correlated moderately with worse SDMT performance (Fig 5U–5X). Reduced $R2t^*$ in left amygdala correlated moderately with worse cognitive test performance on the 2s PASAT (Fig 5K), however, NV showed no significant correlations with PASAT scores (Fig 5E–5H and 5M–5P).




Fig 3. HC versus MS group comparisons of $R2_t^*$ (A–D) and NV (E–H) of hippocampus and amygdala for each hemisphere. Panels with p-values (with FDR correction) ≤ 0.05 and > 0.01 are marked with “**”. Panels with p-values (with FDR correction) ≤ 0.01 are marked with “*”. LH = Left Hemisphere; RH = Right Hemisphere.

<https://doi.org/10.1371/journal.pone.0187915.g003>

Associations of $R2_t^*$ and NV with tests of physical impairment

Reduced $R2_t^*$ of right amygdala showed modest correlation with worse EDSS score ($r = -0.29$, $p = 0.05$) (Fig 6). Smaller NV of right hippocampus displayed a nonsignificant trend to be correlated with worse non-dominant hand performance on 9HPT ($r = -0.24$, $p = 0.09$).

Other limbic system structures

We also examined $R2_t^*$ and NV in other cortical GM regions in the limbic system, including the parahippocampus, insula, and entorhinal, orbitofrontal and cingulate cortices. No significant gender differences in $R2_t^*$ or NV were found in these structures for HC or MS. Compared to HC, our cohort of MS patients had significantly lower $R2_t^*$ in the right insula ($p < 0.01$) and left lateral orbitofrontal cortex ($p < 0.01$). In contrast, no differences were found for NV between HC and MS groups in any of these structures. Considering left and right regions as distinct, lower $R2_t^*$ of 11 limbic cortex regions correlated with worse performance on cognitive tests (bolded, Table 2), whereas no correlations were found for NV. FDR was used to correct for multiple comparisons in all assessments.




Fig 4. HC versus different MS sub-groups comparisons of $R2t^*$ (top) and NV (bottom) of hippocampus and amygdala for each hemisphere. Significant p-values (with FDR correction) are listed in each panel. LH = Left Hemisphere; RH = Right Hemisphere.

<https://doi.org/10.1371/journal.pone.0187915.g004>

Discussion

MRI techniques have been successfully used to detect MS pathology for decades. Current standard clinical sequences include T1-weighted (T1W), T2-weighted (T2W), and fluid attenuated inversion recovery (FLAIR). However, these techniques are not quantitative. In this study, we measured quantitative parameters of $R2t^*$ and volume of limbic system components in 31 HC and 80 MS patients. Lower $R2t^*$ values and smaller normalized volumes were found in the hippocampus, amygdala and several other limbic system regions in MS patients, compared to HC. Both $R2t^*$ and volumetric measures correlated with clinical cognitive tests measuring processing speed, working memory, attention and concentration. Notably, correlations of $R2t^*$ were stronger and involved more regions, compared to volumetric measures. Thus, compared with volumetrics, $R2t^*$ may detect abnormalities earlier or with greater sensitivity. A quantitative tool to sensitively detect changes in the limbic system and other parts of the brain could be useful to guide clinical decisions regarding the effectiveness of a given MS treatment in an individual patient, and might also be useful in trials of potential new therapies.

MRI plays an important role in diagnosis and management of MS. Conventional T1W and T2W imaging techniques allow visualization of many MS lesions in WM but do not correlate well with MS disability. The image contrast in standard “weighted” MR sequences depends on




Fig 5. Correlations of R2t* and NV with clinical cognitive scores in hippocampus and amygdala of MS patients. Correlations of R2t* are shown on the left panels. Correlations of NV are shown on the right panels. Correlations with three clinical cognitive tests (3s PASAT, 2s PASAT and SDMT) are presented in 3 rows, respectively. The r-values and p-values (with FDR correction) are provided in each panel. LH = Left Hemisphere; RH = Right Hemisphere.

<https://doi.org/10.1371/journal.pone.0187915.g005>

only on MR relaxation time constants of the tissue but also on parameters of the pulse sequences, making it intrinsically non-quantitative; this likely contributes to weak correlations with disability measures.

Tissue specific (i.e. free of artifacts related to magnetic field inhomogeneities and physiological fluctuations) $R2^*$ ($1/T2^*$) relaxation rate constant is a quantitative measurement of the transverse relaxation rate of MRI signal in tissue. It can be considered as a quantitative measure reflecting the reciprocal of the degree of hyperintensity of MS lesions on T2W images. Measuring tissue T2 and T1 relaxation times has correlated well with MS pathology seen at autopsy [32]. Such studies suggested that more quantitative tissue-specific relaxation times might provide more accurate and sensitive reflections of pathology than measuring lesion size on “weighted” images.

Alterations of $R2^*$ have been shown to reflect MS pathology, including demyelination [33]. We previously reported that $R2^*$ is reduced in MS in CNS WM [18] and cortical GM [14], and that $R2^*$ measurements in some cortical GM regions correlated with worse cognitive test scores in MS. Because $R2^*$ includes contributions due to neuronal activity—BOLD effect [34], we developed methods [16] to separate the tissue-specific part of $R2^*$ (which we call $R2t^*$) from the total $R2^*$ values. Previously, we demonstrated that changes in $R2t^*$ correlated with age-related changes in cellular density in healthy human brain [16,17] and correlated better than $R2^*$ with cognitive tests in Alzheimer’s Disease patients [23]. In studies reported in this paper, we compared regional $R2t^*$ to NV for ability to detect and measure tissue abnormalities in limbic GM tissues of healthy and MS subjects, and examined relationships of $R2t^*$ and NV with physical and cognitive test scores.

$R2t^*$ and NV of the hippocampus and amygdala were compared between female and male MS and HC. In HC, no significant gender differences were found for NV or for $R2t^*$. However,




Fig 6. Correlations of R2t* and NV with clinical disability scores (EDSS, 9HPT of dominant and non-dominant hands) in hippocampus and amygdala of MS patients. The correlations (r) and p-values (with FDR correction) are provided in each panel. LH = Left Hemisphere; RH = Right Hemisphere.

<https://doi.org/10.1371/journal.pone.0187915.g006>

Table 2. Correlations between clinical cognitive tests and R2t* and NV in limbic cortex.

		R2t*						NV					
		3s PASAT		2s PASAT		SDMT		3s PASAT		2s PASAT		SDMT	
		r	p	r	p	r	p	r	p	r	p	r	p
Parahippocampal	LH	0.01	0.91	0.06	0.63	0.13	0.34	0.07	0.77	-0.02	0.98	0.30	0.09
	RH	0.32	0.03	0.22	0.11	0.24	0.07	0.15	0.77	0.06	0.98	0.33	0.08
Insula	LH	0.31	0.03	0.27	0.05	0.37	0.01	0.15	0.77	0.12	0.97	0.14	0.31
	RH	0.23	0.08	0.23	0.11	0.35	0.01	0.07	0.77	0.00	0.98	0.12	0.35
Entorhinal	LH	-0.17	0.22	-0.16	0.19	-0.07	0.61	0.11	0.77	0.08	0.98	0.23	0.11
	RH	-0.13	0.34	-0.19	0.13	-0.07	0.61	0.05	0.81	0.02	0.98	0.15	0.31
Lateral orbitofrontal	LH	0.21	0.11	0.19	0.14	0.19	0.15	0.10	0.77	0.00	0.98	0.21	0.15
	RH	0.31	0.03	0.35	0.01	0.34	0.01	0.09	0.77	0.04	0.98	0.17	0.24
Medial orbitofrontal	LH	0.16	0.22	0.15	0.23	0.14	0.31	0.12	0.77	0.19	0.91	0.12	0.35
	RH	0.09	0.47	0.19	0.13	0.01	0.93	0.02	0.90	0.10	0.98	0.02	0.88
Caudal anterior cingulate	LH	0.36	0.01	0.42	<0.01	0.42	<0.01	0.00	0.97	-0.02	0.98	0.19	0.20
	RH	0.36	0.01	0.49	<0.01	0.33	0.01	0.26	0.51	0.15	0.91	0.24	0.11
Isthmus of cingulate	LH	0.29	0.03	0.33	0.01	0.29	0.02	0.02	0.90	0.08	0.98	0.27	0.10
	RH	0.29	0.03	0.33	0.01	0.40	<0.01	0.09	0.77	0.12	0.97	0.13	0.32
Posterior cingulate	LH	0.26	0.04	0.19	0.13	0.35	0.01	-0.09	0.77	-0.02	0.98	0.02	0.88
	RH	0.30	0.03	0.27	0.05	0.32	0.01	0.05	0.81	0.15	0.91	0.24	0.11
Rostral anterior cingulate	LH	0.29	0.03	0.40	<0.01	0.22	0.10	0.12	0.77	0.16	0.91	0.15	0.31
	RH	0.11	0.38	0.19	0.13	-0.11	0.43	0.08	0.77	0.06	0.98	0.25	0.11

Lower R2t* and NV each correlate with worse performance on clinical cognitive tests scores in several regions of the limbic system (significant values are bolded). All p-values are after multiple comparison correction (FDR).

<https://doi.org/10.1371/journal.pone.0187915.t002>

in MS patients in our study, males tended to have smaller hippocampi (NV) than females. A previous study [35] has shown that subcortical gray matter atrophy is more significant in male than in female, although that study did not report atrophy in hippocampus. However, hippocampal atrophy has been reported in MS [5–7].

A novel finding in the present study was that MS patients had lower R2t* values in several regions of the limbic system compared to HC, and lower R2t* values significantly correlated with worse performance on cognitive tests relevant to MS. Abnormally low R2t* was seen in right hippocampus, right insula and left lateral orbitofrontal. In our MS cohort, smaller NV in MS than HC was only noted in the hippocampus and amygdala, and no correlations with cognitive scores were found for volumetric measures. Together, these findings suggest that MS damage is widespread within the limbic system, and that R2t* measurements are more sensitive to tissue damage in these structures than is atrophy. Importantly, our results might also indicate that R2t* can detect changes earlier than volume changes, but longitudinal studies will be needed to confirm this. Since volumetric measurements are the current standard in the field to measure tissue damage related to neurodegeneration in MS, this study will provide a quantitative tool with considerable potential to detect abnormalities in the limbic system with more sensitivity and/or earlier.

The amygdala and hippocampus are located in the medial temporal lobe, a region which is involved in recognition of sounds [4]. We observed that reductions of R2t* in both hippocampus and amygdala in MS patients correlated with worse results on the PASAT, a test of auditory information processing speed, calculation ability, and working memory. Correlations with 2s PASAT ($r = 0.36$, $p = 0.007$) were stronger than with 3s PASAT ($r = 0.28$, $p = 0.07$) in left amygdala. This might reflect the greater difficulty of the 2s PASAT, making the latter more

sensitive to mild abnormalities. The amygdala is well known to function in emotional learning and memory [36]. The 2s PASAT is considered more stressful than the 3s PASAT, perhaps evoking an emotional response in the test-taker that is sensitive to amygdala abnormalities. In contrast, NV of the hippocampus and amygdala only correlated with SDMT, but not with 3s or 2s PASAT scores. Interestingly, regional correlations of NV and $R2t^*$ with cognitive test results did not always overlap. These disparate results indicate that $R2t^*$ can contribute information on tissue integrity that volume does not. We found weak or no correlations between $R2t^*$ in limbic system and motor functions tested by 9HPT and 25TW, respectively. This was not unexpected. We suspect any weak correlations with 9HPT are indirect—that is, more a result of the global level of CNS pathology in a given patient than to a direct structure-function correlation.

There are several limitations in the current study. The number of participants was relatively small, which might have limited the ability to detect correlations. Extensive tests of memory, learning and emotion were not included in our study design. We did find that $R2t^*$ and volumetric measures showed significant correlations with tests that are sensitive to common MS cognitive deficits (PASAT and SDMT), including processing speed and working memory. Future studies will include more thorough testing of the primary hippocampus and amygdala functions of memory and learning.

Conclusions

This study reports several novel findings, as well as expanding on existing literature. Limbic system abnormalities have received relatively little prior attention in MS. We found reduced $R2t^*$, a measure that reflects tissue integrity, in several cortical limbic system regions that were not abnormal using volumetrics. $R2t^*$ was also more sensitive to changes in cognitive test performance than NV. High resolution MR sequences to measure $R2t^*$ require less than 15 minutes, and can be readily added to scanning protocols on clinical scanners. This and prior studies indicate that $R2t^*$ can be used as a quantitative tissue biomarker, providing information that is distinct from and complementary to volume.

Acknowledgments

Anne Cross holds the Manny & Rosalyn Rosenthal–Dr. John L. Trotter MS Center Chair in Neuroimmunology of the Foundation for Barnes-Jewish Hospital. Jie Wen is a former post-doctoral fellow of National MS Society USA.

Author Contributions

Conceptualization: Jie Wen, Dmitriy A. Yablonskiy, Anne H. Cross.

Data curation: Jie Wen, Amber Salter.

Formal analysis: Amber Salter.

Funding acquisition: Anne H. Cross.

Investigation: Jie Wen, Dmitriy A. Yablonskiy, Anne H. Cross.

Methodology: Jie Wen, Dmitriy A. Yablonskiy, Amber Salter.

Project administration: Anne H. Cross.

Resources: Dmitriy A. Yablonskiy.

Supervision: Dmitriy A. Yablonskiy, Anne H. Cross.

Validation: Amber Salter.

Writing – original draft: Jie Wen.

Writing – review & editing: Dmitriy A. Yablonskiy, Amber Salter, Anne H. Cross.

References

1. Rao SM, Leo GJ, Bernardin L, Unverzagt F (1991) Cognitive dysfunction in multiple sclerosis. I. Frequency, patterns, and prediction. *Neurology* 41: 685–691. PMID: [2027484](#)
2. Chiaravalloti ND, DeLuca J (2008) Cognitive impairment in multiple sclerosis. *Lancet Neurol* 7: 1139–1151. [https://doi.org/10.1016/S1474-4422\(08\)70259-X](https://doi.org/10.1016/S1474-4422(08)70259-X) PMID: [19007738](#)
3. Amunts K, Kedo O, Kindler M, Pieperhoff P, Mohlberg H, Shah NJ, et al. (2005) Cytoarchitectonic mapping of the human amygdala, hippocampal region and entorhinal cortex: intersubject variability and probability maps. *Anat Embryol (Berl)* 210: 343–352.
4. LeDoux JE (2000) Emotion circuits in the brain. *Annu Rev Neurosci* 23: 155–184. <https://doi.org/10.1146/annurev.neuro.23.1.155> PMID: [10845062](#)
5. Sicotte NL, Kern KC, Giesser BS, Arshanapalli A, Schultz A, Montag M, et al. (2008) Regional hippocampal atrophy in multiple sclerosis. *Brain* 131: 1134–1141. <https://doi.org/10.1093/brain/awn030> PMID: [18375977](#)
6. Audoin B, Zaaraoui W, Reuter F, Rico A, Malikova I, Confort-Gouny S, et al. (2010) Atrophy mainly affects the limbic system and the deep grey matter at the first stage of multiple sclerosis. *Journal of Neurology Neurosurgery and Psychiatry* 81: 690–695.
7. Benedict RH, Ramasamy D, Munschauer F, Weinstock-Guttman B, Zivadinov R (2009) Memory impairment in multiple sclerosis: correlation with deep grey matter and mesial temporal atrophy. *J Neurol Neurosurg Psychiatry* 80: 201–206. <https://doi.org/10.1136/jnnp.2008.148403> PMID: [18829629](#)
8. Koenig KA, Sakaie KE, Lowe MJ, Lin J, Stone L, Bermel RA, et al. (2015) The relationship between cognitive function and high-resolution diffusion tensor MRI of the cingulum bundle in multiple sclerosis. *Multiple Sclerosis Journal* 21: 1794–1801. <https://doi.org/10.1177/1352458515576983> PMID: [26106010](#)
9. Meijer KA, Muhler N, Cercignani M, Sethi V, Ron MA, Thompson AJ, et al. (2016) White matter tract abnormalities are associated with cognitive dysfunction in secondary progressive multiple sclerosis. *Multiple Sclerosis Journal* 22: 1429–1437. <https://doi.org/10.1177/1352458515622694> PMID: [26733423](#)
10. Keser Z, Hasan KM, Mwangi B, Gabr RE, Steinberg JL, Wilken J, et al. (2017) Limbic Pathway Correlates of Cognitive Impairment in Multiple Sclerosis. *Journal of Neuroimaging* 27: 37–42. <https://doi.org/10.1111/jon.12381> PMID: [27541485](#)
11. Yablonskiy DA (2000) Gradient echo plural contrast imaging (GEPC)—New fast magnetic resonance imaging technique for simultaneous acquisition of T2, T1 (or spin density) and T2*-weighted images. *Radiology* 217: 204–204.
12. Sati P, Cross AH, Luo J, Hildebolt CF, Yablonskiy DA (2010) In vivo quantitative evaluation of brain tissue damage in multiple sclerosis using gradient echo plural contrast imaging technique. *NeuroImage* 51: 1089–1097. <https://doi.org/10.1016/j.neuroimage.2010.03.045> PMID: [20338247](#)
13. Luo J, Jagadeesan BD, Cross AH, Yablonskiy DA (2012) Gradient echo plural contrast imaging—signal model and derived contrasts: T2*, T1, phase, SWI, T1f, FST2* and T2*-SWI. *NeuroImage* 60: 1073–1082. <https://doi.org/10.1016/j.neuroimage.2012.01.108> PMID: [22305993](#)
14. Wen J, Yablonskiy DA, Luo J, Lancia S, Hildebolt C, Cross AH, et al. (2015) Detection and quantification of regional cortical gray matter damage in multiple sclerosis utilizing gradient echo MRI. *NeuroImage Clin* 9: 164–175. <https://doi.org/10.1016/j.nicl.2015.08.003> PMID: [27330979](#)
15. Schmalbrock P, Prakash RS, Schirda B, Janssen A, Yang GK, Russell M, et al. (2016) Basal Ganglia Iron in Patients with Multiple Sclerosis Measured with 7T Quantitative Susceptibility Mapping Correlates with Inhibitory Control. *American Journal of Neuroradiology* 37: 439–446. <https://doi.org/10.3174/ajnr.A4599> PMID: [26611996](#)
16. Ulrich X, Yablonskiy DA (2016) Separation of cellular and BOLD contributions to T2* signal relaxation. *Magn Reson Med* 75: 606–615. <https://doi.org/10.1002/mrm.25610> PMID: [25754288](#)
17. Zhao Y, Wen J, Cross AH, Yablonskiy DA (2016) On the relationship between cellular and hemodynamic properties of the human brain cortex throughout adult lifespan. *NeuroImage* 133: 417–429. <https://doi.org/10.1016/j.neuroimage.2016.03.022> PMID: [26997360](#)
18. Sati P, Cross AH, Luo J, Hildebolt CF, Yablonskiy DA (2010) In vivo quantitative evaluation of brain tissue damage in multiple sclerosis using gradient echo plural contrast imaging technique. *NeuroImage* 51: 1089–1097. <https://doi.org/10.1016/j.neuroimage.2010.03.045> PMID: [20338247](#)

19. Ogawa S, Lee TM, Kay AR, Tank DW (1990) Brain Magnetic-Resonance-Imaging with Contrast Dependent on Blood Oxygenation. *Proc Natl Acad Sci U S A* 87: 9868–9872. PMID: [2124706](#)
20. Ogawa S, Lee TM, Kay AR, Tank DW (1990) Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc Natl Acad Sci U S A* 87: 9868–9872. PMID: [2124706](#)
21. Yablonskiy DA, Sukstanskii AL, Luo J, Wang X (2013) Voxel spread function method for correction of magnetic field inhomogeneity effects in quantitative gradient-echo-based MRI. *Magn Reson Med* 70: 1283–1292. <https://doi.org/10.1002/mrm.24585> PMID: [23233445](#)
22. Wen J, Cross AH, Yablonskiy DA (2015) On the role of physiological fluctuations in quantitative gradient echo MRI: implications for GEPCI, QSM, and SWI. *Magn Reson Med* 73: 195–203. <https://doi.org/10.1002/mrm.25114> PMID: [24482009](#)
23. Zhao Y, Raichle ME, Wen J, Benzinger TL, Fagan AM, Hassenstab J, et al. (2017) In vivo detection of microstructural correlates of brain pathology in preclinical and early Alzheimer Disease with magnetic resonance imaging. *Neuroimage* 148: 296–304. <https://doi.org/10.1016/j.neuroimage.2016.12.026> PMID: [27989773](#)
24. Mugler JP, Brookeman JR (1990) Three-dimensional magnetization-prepared rapid gradient-echo imaging (3D MP RAGE). *Magn Reson Med* 15: 152–157. PMID: [2374495](#)
25. Shiee N, Bazin PL, Ozturk A, Reich DS, Calabresi PA, Pham DL, et al. (2010) A topology-preserving approach to the segmentation of brain images with multiple sclerosis lesions. *Neuroimage* 49: 1524–1535. <https://doi.org/10.1016/j.neuroimage.2009.09.005> PMID: [19766196](#)
26. McAuliffe MJ, Lalonde FM, McGarry D, Gandler W, Csaky K, Trus BL, et al. Medical Image Processing, Analysis and Visualization in clinical research; 2001 2001. pp. 381–386.
27. Chiaravalloti ND, Stojanovic-Radic J, DeLuca J (2013) The role of speed versus working memory in predicting learning new information in multiple sclerosis. *J Clin Exp Neuropsychol* 35: 180–191. <https://doi.org/10.1080/13803395.2012.760537> PMID: [23350959](#)
28. Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate—a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Methodological* 57: 289–300.
29. Szabo CA, Lancaster JL, Xiong J, Cook C, Fox P (2003) MR imaging volumetry of subcortical structures and cerebellar hemispheres in normal persons. *AJNR Am J Neuroradiol* 24: 644–647. PMID: [12695196](#)
30. Filipek PA, Richelme C, Kennedy DN, Caviness VS Jr. (1994) The young adult human brain: an MRI-based morphometric analysis. *Cereb Cortex* 4: 344–360. PMID: [7950308](#)
31. Goldstein JM, Seidman LJ, Horton NJ, Makris N, Kennedy DN, Caviness VS Jr, et al. (2001) Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. *Cereb Cortex* 11: 490–497. PMID: [11375910](#)
32. Seewann A, Vrenken H, van der Valk P, Blezer EL, Knol DL, Castelijns JA, et al. (2009) Diffusely abnormal white matter in chronic multiple sclerosis: imaging and histopathologic analysis. *Arch Neurol* 66: 601–609. <https://doi.org/10.1001/archneurol.2009.57> PMID: [19433660](#)
33. Patel K, Luo J, Alvarez E, Piccio L, Schmidt R, Yablonskiy DA, et al. (2015) Detection of cortical lesions in multiple sclerosis: A new imaging approach. *Mult Scler J: Exp Transl Clin* (in press).
34. Yablonskiy DA, Haacke EM (1994) Theory of NMR signal behavior in magnetically inhomogeneous tissues: the static dephasing regime. *Magn Reson Med* 32: 749–763. PMID: [7869897](#)
35. Schoonheim MM, Popescu V, Rueda Lopes FC, Wiebenga OT, Vrenken H, Douw L, et al. (2012) Subcortical atrophy and cognition: sex effects in multiple sclerosis. *Neurology* 79: 1754–1761. <https://doi.org/10.1212/WNL.0b013e3182703f46> PMID: [23019265](#)
36. Phelps EA (2004) Human emotion and memory: interactions of the amygdala and hippocampal complex. *Curr Opin Neurobiol* 14: 198–202. <https://doi.org/10.1016/j.conb.2004.03.015> PMID: [15082325](#)