

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2017

Cerebrospinal fluid A β 42/40 corresponds better than A β 42 to amyloid PET in Alzheimer's disease

Piotr Lewczuk

Universitätsklinikum Erlangen, and Friedrich-Alexander-Universität

Anja Matzen

IBL International GmbH

Kaj Blennow

Sahlgrenska Academy at the University of Gothenburg

Lucilla Parnetti

University of Perugia

Jose Luis Molinuevo

University of Barcelona

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Lewczuk, Piotr; Matzen, Anja; Blennow, Kaj; Parnetti, Lucilla; Molinuevo, Jose Luis; Eusebi, Paolo; Kornhuber, Johannes; Morris, John C.; and Fagan, Anne M., "Cerebrospinal fluid A β 42/40 corresponds better than A β 42 to amyloid PET in Alzheimer's disease." *Journal of Alzheimer's Disease*. 55, 2. 813-822. (2017).

https://digitalcommons.wustl.edu/open_access_pubs/6725

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Piotr Lewczuk, Anja Matzen, Kaj Blennow, Lucilla Parnetti, Jose Luis Molinuevo, Paolo Eusebi, Johannes Kornhuber, John C. Morris, and Anne M. Fagan

Cerebrospinal Fluid A $\beta_{42/40}$ Corresponds Better than A β_{42} to Amyloid PET in Alzheimer's Disease

Piotr Lewczuk^{a,b,1,*}, Anja Matzen^{c,1}, Kaj Blennow^d, Lucilla Parnetti^e, Jose Luis Molinuevo^{f,g}, Paolo Eusebi^e, Johannes Kornhuber^a, John C. Morris^h and Anne M. Fagan^h

^a*Department of Psychiatry and Psychotherapy, Universitätsklinikum Erlangen, and Friedrich-Alexander-Universität, Erlangen-Nürnberg, Erlangen, Germany*

^b*Department of Neurodegeneration Diagnostics, Medical University of Białystok, and Department of Biochemical Diagnostics, University Hospital of Białystok, Białystok, Poland*

^c*IBL International GmbH, Hamburg, Germany*

^d*Clinical Neurochemistry Laboratory, Institute of Neuroscience and Physiology, Department of Psychiatry and Neurochemistry, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden*

^e*Section of Neurology, Center for Memory Disturbances, University of Perugia, Italy*

^f*Alzheimer's disease and other cognitive disorders unit, Neurology Service, Hospital Clínic de Barcelona – Universitat de Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain*

^g*Barcelonaβeta Brain Research Center, Pasqual Maragall Foundation, Barcelona, Spain*

^h*The Knight Alzheimer's Disease Research Center, Department of Neurology, Washington University School of Medicine, St. Louis, MO, USA*

Handling Associate Editor: Sylvain Lehmann

Accepted 31 August 2016

Abstract.

Background: Decreased concentrations of amyloid- β 1-42 (A β_{42}) in cerebrospinal fluid (CSF) and increased retention of A β tracers in the brain on positron emission tomography (PET) are considered the earliest biomarkers of Alzheimer's disease (AD). However, a proportion of cases show discrepancies between the results of the two biomarker modalities which may reflect inter-individual differences in A β metabolism. The CSF A $\beta_{42/40}$ ratio seems to be a more accurate biomarker of clinical AD than CSF A β_{42} alone.

Objective: We tested whether CSF A β_{42} alone or the A $\beta_{42/40}$ ratio corresponds better with amyloid PET status and analyzed the distribution of cases with discordant CSF-PET results.

Methods: CSF obtained from a mixed cohort ($n = 200$) of cognitively normal and abnormal research participants who had undergone amyloid PET within 12 months ($n = 150$ PET-negative, $n = 50$ PET-positive according to a previously published cut-off) was assayed for A β_{42} and A β_{40} using two recently developed immunoassays. Optimal CSF cut-offs for amyloid positivity were calculated, and concordance was tested by comparison of the areas under receiver operating characteristic (ROC) curves (AUC) and McNemar's test for paired proportions.

¹These authors contributed equally to the study.

*Correspondence to: Prof. Dr. med. Piotr Lewczuk, Lab for Clinical Neurochemistry and Neurochemical Dementia Diagnostics, Department of Psychiatry and Psychotherapy,

Schwabachanlage 6, 91054 Erlangen, Germany. Tel.: +49 9131 85 34324; Fax: +49 9131 85 34 238; E-mail: Piotr.Lewczuk@uk-erlangen.de.

Results: CSF A $\beta_{42/40}$ corresponded better than A β_{42} with PET results, with a larger proportion of concordant cases (89.4% versus 74.9%, respectively, $p < 0.0001$) and a larger AUC (0.936 versus 0.814, respectively, $p < 0.0001$) associated with the ratio. For both CSF biomarkers, the percentage of CSF-abnormal/PET-normal cases was larger than that of CSF-normal/PET-abnormal cases.

Conclusion: The CSF A $\beta_{42/40}$ ratio is superior to A β_{42} alone as a marker of amyloid-positivity by PET. We hypothesize that this increase in performance reflects the ratio compensating for general between-individual variations in CSF total A β .

Keywords: Alzheimer's disease, amyloid- β , biomarker, cerebrospinal fluid, positron emission tomography

INTRODUCTION

Reduced concentrations of amyloid- β (A β) 1-42 in the cerebrospinal fluid (CSF) are associated with increased retention of A β tracers in the brain via positron emission tomography (PET) [1-5] and both are considered the earliest biomarkers of Alzheimer's disease (AD) [6-10]. However, several published studies show that results of the two measures are discordant in a proportion of individuals in the cohorts evaluated to date, most notably as CSF A β_{42} -abnormal/amyloid PET-normal (reviewed in [11]). Such discordance may reflect assay methodological variability and/or imprecision, or these biomarker modalities may reflect partly different pathophysiological events [12].

Results from several groups demonstrate that normalization of the CSF A β_{42} concentration to the level of total A β peptides (or the most abundant isoform, A β 1-40 [A β_{40}]) in the CSF by means of an A $\beta_{42/40}$ ratio improves the accuracy of clinical AD diagnosis [13-17] by "controlling for" inter-individual differences in overall levels of CSF A β . This led us to hypothesize that the CSF concentration of A β_{42} depends not only on the pathophysiological status of a given individual, hence suggesting the presence or absence of AD pathology, but also on the total amount of A β peptides in the CSF, i.e., the total production of A β in the brain. At least two additional arguments, derived from experience regarding pre-analytical sample handling and data interpretation, further support the application of the A $\beta_{42/40}$ ratio for AD diagnostic purposes: (a) A $\beta_{42/40}$ is less prone to the error of misinterpretation when inappropriate test tubes to collect CSF are used [18] (likely due to the similar amount of absorption of both A β species); and (b) in two studies that applied the A $\beta_{42/40}$ ratio measured with the same ELISA assays to predict AD in mild cognitive impairment (MCI) subjects [15] or in early symptomatic AD [19], virtually identical A $\beta_{42/40}$ cut-offs were obtained despite differences in absolute A β_{42} concentrations.

Considering these data, we hypothesized that the CSF A $\beta_{42/40}$ ratio might reflect amyloid PET status (positive versus negative) better than the A β_{42} concentration alone. To test this hypothesis, we used dichotomized PET results to calculate the optimal cut-offs for A β_{42} concentrations and A $\beta_{42/40}$ ratios measured with two novel, recently validated ELISAs [14], and then compared the concordance of each of the two CSF biomarkers with the results of the amyloid PET. We also compared the distribution of the discordant results, i.e., the percentage of cases with pathologic CSF findings and normal PET results versus the percentage with normal CSF and pathologic PET results. We proposed that the results from these analyses might provide insight into the possible mechanism(s) for the observed known discordance between amyloid imaging and levels of CSF A β_{42} and, thus, may be informative for pathologic disease staging.

MATERIALS AND METHODS

Participants

All procedures were approved by the Institutional Review Board and the Washington University Human Research Protection Office, and written informed consent was obtained from each participant. Participants were community-dwelling volunteers enrolled in studies of normal aging and dementia at the Knight Alzheimer's Disease Research Center (ADRC) at Washington University in St. Louis. Participants were 45-83 years of age at baseline and had no neurological, psychiatric or systemic medical illness that might compromise longitudinal study participation, nor medical contraindication to lumbar puncture (LP) or PET. All participants underwent cognitive assessments (including the Mini-Mental State Examination (MMSE) [20]), neurological evaluations and clinical assessments that included the Clinical Dementia Rating (CDR) [21].

To be included in the present analysis, baseline clinical and cognitive assessment, and LP and amyloid PET (via Pittsburgh Compound B, PiB) within 12 months of each other had to be available (mean \pm SD, 67 ± 78 days). Two-hundred CSF samples were selected *a priori* based on cortical amyloid load via PET (25% amyloid PET-positive and 75% amyloid PET-negative, see below), independent of and blinded to participant demographics and clinical status. Given published reports of greater CSF-PET discordance in amyloid PET-negative individuals, greater numbers of amyloid-negative participant samples were evaluated so to enrich for possible and more common discordant (PiB-negative/A β_{42} -positive) cases. APOE genotype was obtained from the Knight ADRC Genetics Core [22].

CSF collection, processing, and analysis

CSF was collected and processed according to standardized protocols. Samples (20–30 mL) were collected using LP procedures (22 G \times 3 1/2 inch atraumatic Sprotte spinal needle, Pajunk, #001151-30 C) via gravity drip into a 50 mL conical polypropylene tube (Corning #430829) at 8:00am after overnight fasting, gently inverted to avoid possible rostro-caudal gradient effects and immediately put on wet ice. Within 1 h of collection, samples were briefly centrifuged at low speed (2000 \times g, 5 min at 4°C) to pellet any cellular debris. All but the bottom 500 μ L was transferred to a new 50 mL polypropylene tube, aliquoted (500 μ L) into sterile 2 mL polypropylene screw-cap tubes with O-rings (Corning #430915), and immediately frozen at -80°C until analysis. The interval between sample collection and batch analysis varied from 1–9 years, with equal representation of storage intervals among the different amyloid PET groups (PiB+ versus PiB-). Samples underwent a single freeze-thaw cycle prior to assay, were thawed on wet ice (\sim 3 h) followed by vortexing prior to blinded analysis, and were run in duplicate. A β_{42} and A β_{40} concentrations were measured with immunoassays from IBL International (Hamburg, Germany) according to the manufacturer's protocols. To be included in the statistical analyses, the results had to pass quality control (QC) criteria consistent with the Alzheimer's Disease Neuroimaging Initiative (ADNI) study [23], including coefficients of variability (CV) of duplicates $\leq 25\%$ and the results of the kit's run validation samples ("kit controls") within the acceptable range as defined by the manufacturer. Samples also had data available for A β_{42} , Tau, and

pTau181 using INNOTEST ELISA kits (Fujirebio, Ghent, Belgium).

In vivo amyloid imaging

Participants received a single intravenous bolus injection of ^{11}C PiB (between 4.7 and 19.0 mCi). PET imaging [24] was conducted with a Siemens 962 HR+ ECAT PET or Biograph 40 scanner (Siemens/CTI, Knoxville KY), and magnetic resonance imaging (MRI) using MPRAGE T1-weighted images (1 mm \times 1 mm \times 1.25 mm) was obtained for anatomic reference as described previously [25, 26]. Participants underwent a 60-min dynamic scan with ^{11}C PiB, and PET data was processed using regions of interest (ROIs) derived from structural MRIs using Freesurfer [25, 26]. Regional time-activity curves were extracted for each ROI, and binding potentials were calculated using Logan graphical analysis [27] with cerebellar gray matter as the reference region. The mean cortical binding potential (MCPB) was calculated based on a set of selected cortical regions known to have high levels of deposition in AD [28], including precuneus, rostral middle frontal cortex, superior frontal cortex, superior temporal cortex, middle temporal cortex, lateral orbitofrontal cortex, and medial orbitofrontal cortex [25]. A MCPB cutoff of >0.18 for positivity was chosen to be consistent with prior studies [9, 28–30].

Statistical analyses

Sample size was calculated with the negative/positive ratio, power, and significance set to 3:1, 0.9, and 0.05, respectively, and the areas under the receiver operating characteristic (ROC) curves (AUCs) of 0.974 (A $\beta_{42/40}$) and 0.827 (A β_{42}) [14]. Unless otherwise stated, results are presented as means \pm standard deviations (SD) or 95% confidence intervals (CI). CSF A β_{42} and A $\beta_{42/40}$ cut-offs, and the corresponding sensitivities and specificities for PiB PET-positivity, were calculated based on the dichotomous PET grouping at the maximal Youden Indices, using a cut-off for PiB positivity of 0.18 [28]. Comparisons of continuous variables were performed with Student's *t*-test or Welch test in cases of unequal variances; dichotomous variables were compared with chi-squared test. AUCs of the ROC curves were compared with DeLong test [32]. McNemar's test for paired proportions was used for testing differences between the proportions of CSF-PiB concordant cases, and sensitivities and specificities. All

tests were two-tailed. Logistic regression models were applied with logit of probability of PET-positivity (logit ($\pi_{\text{PET}+}$)) as the outcome, and the CSF biomarkers and age as continuous, and APOE $\epsilon 4$ status (presence of $\epsilon 4$ allele = 1) and sex (males = 1) as dichotomous predictors. Covariates were selected either for epidemiological reasons (age, sex, APOE) or due to observed significances between PET+ and PET- groups (A β_{40} concentrations). Mann-Whitney U test was used to compare LP/PET intervals between “CSF/PET concordant” and “CSF/PET discordant” groups separately for A β_{42} and A $\beta_{42/40}$. MedCalc 13.1 (MedCalc Software bvba, Ostend, Belgium) and Stata 14.1 (StatCorp, College Station, TX) were used for statistical analyses. Significance was defined as $p < 0.05$.

RESULTS

Demographic data and CSF biomarkers

As expected given the 3 : 1 (PiB-:PiB+) selection criterion based on amyloid PET positivity, the cohort was comprised of 88% cognitively normal (CDR 0) and 12% cognitively abnormal (CDR >0 ; $n = 19$ CDR 0.5, $n = 3$ CDR 1, $n = 1$ CDR 2) individuals. The cohort demographics and biomarker data are shown in Table 1. One sample did not pass QC so was excluded from statistical analysis. Not unexpectedly, the amyloid-positive (PiB+) group was older than the amyloid-negative (PiB-) group ($p < 0.001$), contained a higher percentage of APOE $\epsilon 4$ carriers ($p < 0.01$), and had a lower mean score on the MMSE ($p = 0.02$). The PiB- group contained a greater percentage of women than the PiB+ group (65% versus 43%, respectively), but there were no differences in the years of education ($p = 0.28$). Median absolute intervals between the time of CSF collection and PiB

PET scan did not differ significantly between the two groups (24 versus 33 days, $U = 3408$, $p = 0.45$), and in both groups PET was performed before LP in similar proportions of the cases (36.7% and 38.7% of PET+ and PET- participants, respectively, $\chi^2(1) = 0.058$, $p = 0.81$).

Inter-assay CVs, tested with two kit control samples, were 2.2% and 7.1% for A β_{40} , and 2.3% and 6.5% for A β_{42} . Median range-to-averages of the duplicate CSF samples measurements (expressing intra-assay variation) were 2.3% for A β_{40} and 2.5% for A β_{42} .

By definition, the amount of cortical amyloid deposition (MCBP, mean \pm SD) was higher in the PiB+ compared to the PiB- group ($p < 0.001$). As expected, PiB+ participants had significantly lower concentrations of CSF A β_{42} ($p < 0.001$) and a lower mean A $\beta_{42/40}$ ratio ($p < 0.001$), as well as significantly higher concentrations of CSF Tau ($p < 0.001$) and pTau181 ($p < 0.001$). Interestingly, the PiB+ group exhibited a modest but significantly higher mean concentration of CSF A β_{40} compared to the PiB- group ($p = 0.047$). Differences in LP/PET intervals between “CSF/PET concordant” and “CSF/PET discordant” groups were not significant ($p > 0.1$ for both CSF biomarkers).

Concordance between amyloid PET and CSF biomarkers

The relationships between CSF analyte values and cortical amyloid load (PiB MCBP) are shown in Fig. 1. The optimal CSF cut-offs that best differentiated the dichotomized PiB+ and PiB- groups were 735 pg/mL for A β_{42} , 11,600 pg/mL for A β_{40} , and 0.05 for the A $\beta_{42/40}$ ratio. At these cut-offs, the A β_{40} , A β_{42} concentrations and the A $\beta_{42/40}$ ratio yielded sensitivities (with dichotomized PiB PET status defined as the reference) of 85.7% (CI: 72.8–94.1),




Fig. 1. Scatterplots of cortical amyloid PET load using [^{11}C] PiB and concentrations of CSF A β_{42} (A), the A $\beta_{42/40}$ ratio (B), and A β_{40} (C). Solid vertical line represents the *a priori* dichotomous cut-off for PiB positivity as published previously [28]. Dashed horizontal lines on panels A and B represent the best-performing cut-offs of the respective CSF biomarkers calculated in the present study.

Table 1
Demographic and biomarker data

Characteristic	PiB+	PiB-	<i>p</i>
N	49	150	NA
Female, %	43%	65%	<0.01*
Age at LP, y	72.4 \pm 7.3	64.2 \pm 9.6	<0.001
% <i>APOE</i> ϵ 4 carriers	58%	31%	<0.01*
Education, y	15.4 \pm 2.9	15.9 \pm 2.5	0.283
MMSE	28.00 \pm 3.03	29.07 \pm 1.21	0.02
Median LP-PiB PET interval (IQR), days	24 (16 – 64)	33 (12–94)	0.45 [†]
PiB MCBP	0.5275 \pm 0.2454	0.0473 \pm 0.0502	<0.001
CSF A β_{40} , pg/mL	15 280 \pm 4 160	13 890 \pm 4 260	0.047
CSF A β_{42} , pg/mL	577 \pm 184	887 \pm 292	<0.001
CSF A $\beta_{42/40}$	0.038 \pm 0.009	0.065 \pm 0.013	<0.001
CSF Tau, pg/mL	492 \pm 392	234 \pm 125	<0.001
CSF pTau181, pg/mL	75.8 \pm 35.3	47.2 \pm 19.1	<0.001

* χ^2 test. [†]Mann-Whitney test. *APOE*, apolipoprotein E; CSF, cerebrospinal fluid; IQR, interquartile range; LP, lumbar puncture; MCBP, mean cortical binding potential; MMSE, Mini-Mental State Examination (0–30, with 30 a perfect score) at clinical assessment closest in time to LP; NA, not applicable; PET, positron emission tomography; PiB, Pittsburgh Compound B. Unless otherwise indicated, values represent the mean \pm SD.

81.6% (CI: 68.0–91.2) and 95.9% (CI: 86.0–99.5), respectively, with the corresponding specificities of 34.0% (CI: 26.5–42.2), 72.0% (CI: 64.8–79.6), and 88.0% (CI: 81.7–92.7), corresponding to accuracies of 46.7% for A β_{40} , 74.9% for A β_{42} , and 89.4% for the A $\beta_{42/40}$ ratio.

For both A β_{42} and A $\beta_{42/40}$, the percentage of cases with abnormal CSF findings but normal PET results (false positives: 20.6% and 9.0% for A β_{42} and A $\beta_{42/40}$, respectively), clustering in the lower left quadrant of Fig. 1, was larger than the percentage of cases with normal CSF findings and abnormal PET (false negatives: 4.5% and 1.5% for A β_{42} and A $\beta_{42/40}$, respectively), clustering in the upper right quadrant of Fig. 1. Detailed results are presented in Table 2.

McNemar test for paired proportions revealed a highly significantly larger proportion of A $\beta_{42/40}$ -PET concordant cases compared to A β_{42} -PET concordant cases ($\chi^2(1) = 19.56$, $p < 0.001$), with an odds ratio of discordant cases of 5.14 (CI: 2.26 to 13.69).

ROC curves yielded AUC's of 0.599 for A β_{40} , 0.814 for A β_{42} and 0.936 for the A $\beta_{42/40}$ ratio (Fig. 2), with the A $\beta_{42/40}$ ratio demonstrating significantly higher correspondence with PiB status compared to the concentrations of A β_{42} ($p < 0.001$) and A β_{40} ($p < 0.001$) (Table 2).

Logistic regression models

Logistic regression models are presented in Supplementary Table 1. In both models (with A β_{42} and A $\beta_{42/40}$, respectively), when adjusted for A β_{40} ,




Fig. 2. Receiver operating characteristic (ROC) curves of the CSF A β_{42} concentration and A $\beta_{42/40}$ ratio. Values for areas under the ROC curves (AUC) correspond to mean \pm 95% CI. Open circles represent the combinations of the sensitivities and the specificities at the maximal Youden Indices.

age, *APOE* ϵ 4 status, and sex, the effects of A β_{42} and A $\beta_{42/40}$ remained highly significant ($p < 0.001$). Consistent with increased A β_{40} in the PiB+ group, the effect of A β_{40} was positive and significant ($p < 0.001$) in the A β_{42} model, but became non-significant in the A $\beta_{42/40}$ model ($p = 0.36$). In both models, the effects of age were positive and significant. Male sex was a positive predictor but only with borderline significance, and presence of at least one *APOE* ϵ 4 allele was a non-significant covariate in both models ($p > 0.05$).

Table 2
Analyses of concordance of CSF A β_{42} and the A $\beta_{42/40}$ ratio with amyloid PET status

Parameter	A β_{42}	A $\beta_{42/40}$
Cut-off at the max. YI	735 pg/mL	0.05
Sensitivity at the max. YI (95% CI)	81.6% (68.0–91.2)	95.9% (86.0–99.5)
Specificity at the max. YI (95% CI)	72.7% (64.8–79.6)	88.0% (81.7–92.7)
Positive PV (95% CI)	49.4% (38.1–60.7)	72.3% (59.8–82.7)
Negative PV (95% CI)	92.4% (86.0–96.5)	98.5% (94.7–99.8)
Accuracy (concordance proportion)	74.9%	89.4%*
%False Pos. (N)/%False Neg. (N)	20.6% (41)/4.5% (9)	9.0% (18)/1.5% (3)
Area Under the ROC curve (95% CI)	0.814 (0.752–0.865)	0.936 [†] (0.892–0.966)

* $p < 0.001$ compared to the A β_{42} concentrations (McNemar test for paired proportions); [†] $p < 0.001$ compared to the A β_{42} concentrations (DeLong test for differences in AUC's); False positive, abnormal CSF/normal PET; false negative, normal CSF/abnormal PET; CSF, cerebrospinal fluid; PET, positron emission tomography; PV, predictive value; ROC, receiver operating characteristic; YI, Youden index.

DISCUSSION

The main finding in this study is significantly better concordance of the CSF A $\beta_{42/40}$ ratio, compared to CSF A β_{42} alone, with cortical A β deposition as measured by [^{11}C] PiB PET. This finding is consistent with our [14] and others' [16, 17] reports of better diagnostic accuracy of the A $\beta_{42/40}$ ratio compared to A β_{42} in early AD, as well as with CSF biomarkers of neurodegeneration, Tau and pTau181 [13]. We hypothesize that the concentration of A β_{42} in the CSF depends not only on the physiological amyloid status of a given individual (presence or absence of amyloid pathology) but also on the total amount of A β peptides that are present. By normalizing to the concentration of A β_{40} , the most abundant A β species in the CSF, the ratio removes the potential confound of differences in overall A β concentration and provides a better index of underlying amyloid-related pathology. Two experimental observations further support the application of the A $\beta_{42/40}$ ratio for AD diagnostics: (a) A $\beta_{42/40}$ is less prone to the error of misinterpretation when non-polypropylene test tubes (as are often found in standard clinical LP trays) are used to collect CSF [18] likely due to the fact that both A β isoforms seem to absorb to the tube surface to a similar extent; and (b) two studies using a different set of ELISAs (produced by a currently non-existing vendor and based on N-terminally non-specific antibodies) that have applied the A $\beta_{42/40}$ ratio to predict AD in MCI subjects [15] and discriminate early symptomatic AD from controls [19] reported almost identical A $\beta_{42/40}$ cut-offs (0.095 and 0.098, respectively), whereas the corresponding cut-offs for A β_{42} alone differed by more than 15% (640 pg/mL and 550 pg/mL, respectively).

The analyses of CSF A β isoforms in the current study were performed with two novel assays that

were validated recently by our group and approved for human *in vitro* diagnostic (IVD) use in Europe [14]. The assays exhibit very good analytical performance, both apply the same protocol for sample dilution, incubation, and washing steps, and require only a small volume (25 μL) of CSF for both biomarkers measured in duplicate. Interestingly, we observed small but significantly ($p = 0.047$) elevated levels of CSF A β_{40} in the PiB+ group compared to the PiB-group. This is consistent with similarly slight, but significant, increases observed in the MCI/AD group compared to controls when measured with the same assay [14]. Also, in a recent study by Dorey et al. [33] investigating the performance of these markers for correspondence with clinical diagnosis, CSF A β_{40} concentrations were higher in AD than non-AD patients, and inclusion of CSF A β_{40} as an A $\beta_{42/40}$ ratio corrected 76.2% of misinterpreted cases in AD patients with normal CSF A β_{42} concentrations and 94.7% of cases when A β_{40} was used alone. When we adjusted for the A $\beta_{42/40}$ ratio, but not for A β_{42} alone, the A β_{40} concentration became insignificant as a predictor of amyloid PET status in the logistic regression models, whereas both A β_{42} and the A $\beta_{42/40}$ ratio remained highly significant as predictors of amyloid PET after adjustment for A β_{40} , age, *APOE* $\epsilon 4$ carriage, and sex. After adjustment for other covariates, older age remained a significant predictor of cortical amyloid-PET positivity, consistent with age being the strongest risk factor for AD. Odds ratio of A β_{42} -PET versus A $\beta_{42/40}$ -PET discordant cases, resulting from McNemar test, indicates that the odds to falsely predict the PET outcome is five-fold lower with the A $\beta_{42/40}$ ratio compared to A β_{42} alone.

Both biomarker modalities (CSF analysis and amyloid PET) have their practical advantages and disadvantages in terms of cost (actual and reimbursable), perceived invasiveness, and availability in clinical as

well as research settings [34], and our results do not endorse one or the other as a “gold standard” for amyloid biomarker positivity. From a scientific point of view, the discordance between CSF A β_{42} and amyloid PET (typically observed as false positives, i.e., abnormal/low CSF A β_{42} in the absence of amyloid-PET positivity) has been hypothesized to be due to possible methodological variability/limitations in the CSF assays, differences in the temporal trajectories of processes identified by the two biomarker modalities (e.g., CSF levels of soluble A β_{42} becoming abnormal prior to aggregated A β_{42} [amyloid] becoming detectable by PET), and/or inter-individual differences in overall levels of A β peptides [9, 12]. These possible scenarios are not mutually exclusive. Our use of well-performing assays that utilize the same dilution factor for both A β_{42} and A β_{40} minimizes methodological variability as a source of discordance between CSF A β_{42} and amyloid PET results and increases the confidence of the ratio as being an accurate measure of the proportion of each of these species in CSF. In so doing, the increase in concordance that we observe for the A $\beta_{42/40}$ ratio suggests much of the “false positives” as defined by CSF A β_{42} is due to lower overall levels of A β in these individuals, whereas the “false negatives” can be attributed in large part to higher overall CSF A β levels. Indeed, the negative predictive value of the A $\beta_{42/40}$ ratio in this study (98%) approached 100%, thus supporting its use for ruling out the presence of underlying amyloid pathology. While use of the ratio also significantly increased the positive predictive value for cortical amyloid PET positivity, 9% of individuals in this cohort were still classified as amyloid-positive by CSF A $\beta_{42/40}$ but amyloid-negative by PET (false negative), consistent with CSF A β_{42} levels dropping prior to cortical amyloid being detectable by PiB PET. That all but one participant in this discordant group (whether CSF abnormality was defined by A β_{42} alone or the A $\beta_{42/40}$ ratio) were cognitively normal is consistent with findings from a recent study in which low CSF A β_{42} but normal amyloid PET was much more frequent in cognitively normal individuals and early MCI cases than in the late MCI stage or AD dementia [12]. Longitudinal CSF and amyloid PET biomarker evaluation will be required to test whether levels of CSF A β_{42} indeed become abnormal (drop) before amyloid-positivity is detected with PET as has been suggested by a recent study by Vlassenko and colleagues [30]. That study, however, did not assess CSF A β_{40} so comparison of the two CSF measures was not possible. Results of a recent

study by Palmqvist et al. also suggest that decreases in CSF A β_{42} precede alterations in amyloid PET [35]. Interestingly, another recent study using different ELISAs also reported better correspondence of the A $\beta_{42/40}$ ratio with A β PET, however, with a higher proportion of A β -/PET+ individuals compared to the present study [36]. This discrepancy is likely due to the differences in the cohorts (higher numbers of cognitively normal individuals expected to be in the preclinical period in the current cohort) and/or different methodological issues. Another study reported no significant differences in concordances between A $\beta_{42/40}$ and A β_{42} with amyloid PET, although the AUC for A $\beta_{42/40}$ was slightly larger than that for A β_{42} alone [37]. This lack of significance might be attributed to the lower statistical power (a much smaller cohort, $n=38$) and/or the use of different ELISAs and PET tracer (^{18}F -flutemetamol) in that study. Similarly, another smaller study concluded that the A $\beta_{42/40}$ ratio had a “large” effect size (defined as Cohen’s $f^2=0.35$), compared to a “medium” effect size (defined as Cohen’s $f^2=0.15$), for A β_{42} , in predicting concurrent PiB burden [38]. Aggregation of soluble A β into diffuse, non-fibrillar plaques that are not identifiable by amyloid PET imaging may also contribute to the “false negative” discordant cases [39]. If confirmed, a finding of CSF alterations preceding those in PET would support the CSF A $\beta_{42/40}$ ratio (and perhaps even A β_{42} alone, albeit with less accuracy) as the earliest detectable marker of β -amyloidosis.

Our study is not without limitations. Although amyloid PET in this study is considered to be the reference to define the presence of amyloid, positivity was defined according to dichotomous, research-based thresholds in a defined set of cortical regions. Also, since samples were specifically selected to over-represent possible CSF/PET discordance (by evaluating 25% PET-positive and 75% PET-negative cases), the cohort does not reflect the general distribution of amyloid PET positivity in the general population. Furthermore, because of this strategy, the cohort was significantly skewed toward cognitively normal individuals (88% cognitively normal, 12% cognitively abnormal). Additional study is needed in larger cohorts comprised of greater numbers of cognitively abnormal individuals, as well as those clinically diagnosed with non-AD dementias in order to better evaluate the CSF/PET concordance in non-AD populations. Finally, we performed statistical analyses of concordance after calculating optimal cut-off values for both CSF biomarkers. As this was designed to be a

biomarker concordance study, independent of clinical status, the diagnostic value of any of the biomarkers and modalities studied here remains to be evaluated.

In summary, the CSF A β _{42/40} ratio performs better, compared to CSF A β ₄₂ alone, in identifying individuals with versus without underlying cortical amyloid as defined by PiB PET. The ratio outperforms the single A β ₄₂ measure in all metrics, including sensitivity, specificity, positive and negative predictive values, and overall accuracy. Of special note is its better concordance in individuals classified as amyloid-negative by PET, suggesting that the higher discordance between CSF A β ₄₂ alone and amyloid PET-positivity may reflect overall lower levels of A β ₄₂ in some individuals as opposed to its aggregation into plaques (or oligomeric forms that are likely not detected with the current assays [40]. The CSF A β _{42/40} ratio may, therefore, be useful as a proxy for amyloid status in AD clinical trials and eventually in clinical care settings.

ACKNOWLEDGMENTS

This study was supported by IBL International GmbH, Hamburg, Germany, and grants P50 AG05681, P01 AG03991 and P01 AG026276 from the National Institute on Aging/National Institute of Health (NIA/NIH) (JC Morris, PI). The sponsor and all coauthors participated in the design of the study, and in the collection, analysis and interpretation of the data; the sponsor had no influence on the decision to submit the paper for publication. The research leading to these results has received support from the Innovative Medicines Initiative Joint Undertaking under grant agreement nr. 115372, resources of which are composed of financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in kind contribution. PL was supported by the German Bundesministerium für Bildung und Forschung (grant 01ED1203D) within the BiomarkAPD Project of the JPND. We cordially thank Michael Habig and Oliver Schmidt of IBL International for the development of the assays and extremely helpful discussions. We thank the research participants at the Knight Alzheimer's Disease Research Center (ADRC) and their families, as well as the members of the Knight ADRC Clinical, Biomarker, Genetics and Imaging Cores for providing data.

Authors' disclosures available online (<http://j-alz.com/manuscript-disclosures/16-0722r1>).

SUPPLEMENTARY MATERIAL

The supplementary material is available in the electronic version of this article: <http://dx.doi.org/10.3233/JAD-160722>.

REFERENCES

- [1] Fagan AM, Mintun MA, Mach RH, Lee SY, Dence CS, Shah AR, LaRossa GN, Spinner ML, Klunk WE, Mathis CA, DeKosky ST, Morris JC, Holtzman DM (2006) Inverse relation between in vivo amyloid imaging load and cerebrospinal fluid Abeta42 in humans. *Ann Neurol* **59**, 512-519.
- [2] Forsberg A, Engler H, Almkvist O, Blomquist G, Hagman G, Wall A, Ringheim A, Langstrom B, Nordberg A (2008) PET imaging of amyloid deposition in patients with mild cognitive impairment. *Neurobiol Aging* **29**, 1456-1465.
- [3] Grimmer T, Riemenschneider M, Forstl H, Henriksen G, Klunk WE, Mathis CA, Shiga T, Wester HJ, Kurz A, Drzezga A (2009) Beta amyloid in Alzheimer's disease: Increased deposition in brain is reflected in reduced concentration in cerebrospinal fluid. *Biol Psychiatry* **65**, 927-934.
- [4] Tolboom N, van der Flier WM, Yaqub M, Boellaard R, Verwey NA, Blankenstein MA, Windhorst AD, Scheltens P, Lammertsma AA, van Berckel BN (2009) Relationship of cerebrospinal fluid markers to 11C-PiB and 18F-FDDNP binding. *J Nucl Med* **50**, 1464-1470.
- [5] Jagust WJ, Landau SM, Shaw LM, Trojanowski JQ, Koeppe RA, Reiman EM, Foster NL, Petersen RC, Weiner MW, Price JC, Mathis CA (2009) Relationships between biomarkers in aging and dementia. *Neurology* **73**, 1193-1199.
- [6] Bateman RJ, Xiong C, Benzinger TL, Fagan AM, Goate A, Fox NC, Marcus DS, Cairns NJ, Xie X, Blazey TM, Holtzman DM, Santacruz A, Buckles V, Oliver A, Moulder K, Aisen PS, Ghetti B, Klunk WE, McDade E, Martins RN, Masters CL, Mayeux R, Ringman JM, Rossor MN, Schofield PR, Sperling RA, Salloway S, Morris JC (2012) Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med* **367**, 795-804.
- [7] Buchhave P, Minthon L, Zetterberg H, Wallin AK, Blennow K, Hansson O (2012) Cerebrospinal fluid levels of beta-amyloid 1-42, but not of tau, are fully changed already 5 to 10 years before the onset of Alzheimer dementia. *Arch Gen Psychiatry* **69**, 98-106.
- [8] Sutphen CL, Jasielec MS, Shah AR, Macy EM, Xiong C, Vlassenko AG, Benzinger TL, Stoops EE, Vanderstichele HM, Brix B, Darby HD, Vandijck ML, Ladenson JH, Morris JC, Holtzman DM, Fagan AM (2015) Longitudinal cerebrospinal fluid biomarker changes in preclinical Alzheimer disease during middle age. *JAMA Neurol* **72**, 1029-1042.
- [9] Fagan AM, Mintun MA, Shah AR, Aldea P, Roe CM, Mach RH, Marcus D, Morris JC, Holtzman DM (2009) Cerebrospinal fluid tau and ptau(181) increase with cortical amyloid deposition in cognitively normal individuals: Implications for future clinical trials of Alzheimer's disease. *EMBO Mol Med* **1**, 371-380.
- [10] Morris JC, Roe CM, Xiong C, Fagan AM, Goate AM, Holtzman DM, Mintun MA (2010) APOE predicts amyloid-beta but not tau Alzheimer pathology in cognitively normal aging. *Ann Neurol* **67**, 122-131.

- [11] Blennow K, Mattsson N, Scholl M, Hansson O, Zetterberg H (2015) Amyloid biomarkers in Alzheimer's disease. *Trends Pharmacol Sci* **36**, 297-309.
- [12] Mattsson N, Insel PS, Donohue M, Landau S, Jagust WJ, Shaw LM, Trojanowski JQ, Zetterberg H, Blennow K, Weiner MW, Alzheimer's Disease Neuroimaging Initiative (2015) Independent information from cerebrospinal fluid amyloid-beta and florbetapir imaging in Alzheimer's disease. *Brain* **138**, 772-783.
- [13] Wiltfang J, Esselmann H, Bibl M, Hüll M, Hampel H, Kessler H, Frölich L, Schröder J, Peters O, Jessen F, Luckhaus C, Perneczky R, Jahn H, Fiszer M, Maler JM, Zimmermann R, Bruckmoser R, Kornhuber J, Lewczuk P (2007) Amyloid beta peptide ratio 42/40 but not A beta 42 correlates with phospho-Tau in patients with low- and high-CSF A beta 40 load. *J Neurochem* **101**, 1053-1059.
- [14] Lewczuk P, Leleental N, Spitzer P, Maler JM, Kornhuber J (2015) Amyloid β 42/40 CSF concentration ratio in the diagnostics of Alzheimer's disease: Validation of two novel assays. *J Alzheimers Dis* **43**, 183-191.
- [15] Hansson O, Zetterberg H, Buchhave P, Andreasson U, Londos E, Minthon L, Blennow K (2007) Prediction of Alzheimer's disease using the CSF Abeta42/Abeta40 ratio in patients with mild cognitive impairment. *Dement Geriatr Cogn Disord* **23**, 316-320.
- [16] Slaets S, Le Bastard N, Martin JJ, Sleegers K, Van Broeckhoven C, De Deyn PP, Engelborghs S (2013) Cerebrospinal fluid Abeta1-40 improves differential dementia diagnosis in patients with intermediate P-tau181P levels. *J Alzheimers Dis* **36**, 759-767.
- [17] Beaufils E, Dufour-Rainfray D, Hommet C, Brault F, Cottier JP, Ribeiro MJ, Mondon K, Guilloteau D (2013) Confirmation of the amyloidogenic process in posterior cortical atrophy: Value of the Abeta42/Abeta40 ratio. *J Alzheimers Dis* **33**, 775-780.
- [18] Lewczuk P, Beck G, Esselmann H, Bruckmoser R, Zimmermann R, Fiszer M, Bibl M, Maler JM, Kornhuber J, Wiltfang J (2006) Effect of sample collection tubes on cerebrospinal fluid concentrations of tau proteins and amyloid beta peptides. *Clin Chem* **52**, 332-334.
- [19] Lewczuk P, Esselmann H, Otto M, Maler JM, Henkel AW, Henkel MK, Eikenberg O, Antz C, Krause WR, Reulbach U, Kornhuber J, Wiltfang J (2004) Neurochemical diagnosis of Alzheimer's dementia by CSF A β 42, A β 42/A β 40 ratio and total tau. *Neurobiol Aging* **25**, 273-281.
- [20] Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* **12**, 189-198.
- [21] Morris JC (1993) The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology* **43**, 2412-2414.
- [22] Pastor P, Roe CM, Villegas A, Bedoya G, Chakraverty S, Garcia G, Tirado V, Norton J, Rios S, Martinez M, Kosik KS, Lopera F, Goate AM (2003) Apolipoprotein Eepsilon4 modifies Alzheimer's disease onset in an E280A PS1 kindred. *Ann Neurol* **54**, 163-169.
- [23] Shaw LM, Vanderstichele H, Knapik-Czajka M, Figurski M, Coart E, Blennow K, Soares H, Simon AJ, Lewczuk P, Dean RA, Siemers E, Potter W, Lee VM, Trojanowski JQ (2011) Qualification of the analytical and clinical performance of CSF biomarker analyses in ADNI. *Acta Neuropathol* **121**, 597-609.
- [24] Klunk WE, Engler H, Nordberg A, Wang Y, Blomqvist G, Holt DP, Bergstrom M, Savitcheva I, Huang GF, Estrada S, Aussen B, Debnath ML, Barletta J, Price JC, Sandell J, Lopresti BJ, Wall A, Koivisto P, Antoni G, Mathis CA, Langstrom B (2004) Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann Neurol* **55**, 306-319.
- [25] Su Y, D'Angelo GM, Vlassenko AG, Zhou G, Snyder AZ, Marcus DS, Blazey TM, Christensen JJ, Vora S, Morris JC, Mintun MA, Benzinger TL (2013) Quantitative analysis of PiB-PET with FreeSurfer ROIs. *PLoS One* **8**, e73377.
- [26] Su Y, Blazey TM, Snyder AZ, Raichle ME, Marcus DS, Ances BM, Bateman RJ, Cairns NJ, Aldea P, Cash L, Christensen JJ, Friedrichsen K, Hornbeck RC, Farrar AM, Owen CJ, Mayeux R, Brickman AM, Klunk W, Price JC, Thompson PM, Ghetti B, Saykin AJ, Sperling RA, Johnson KA, Schofield PR, Buckles V, Morris JC, Benzinger TL, Dominantly Inherited Alzheimer N (2015) Partial volume correction in quantitative amyloid imaging. *Neuroimage* **107**, 55-64.
- [27] Logan J, Fowler JS, Volkow ND, Wolf AP, Dewey SL, Schlyer DJ, MacGregor RR, Hitzemann R, Bendriem B, Gatley SJ, et al. (1990) Graphical analysis of reversible radioligand binding from time-activity measurements applied to [N-11C-methyl]-(-)-cocaine PET studies in human subjects. *J Cereb Blood Flow Metab* **10**, 740-747.
- [28] Mintun MA, Larossa GN, Sheline YI, Dence CS, Lee SY, Mach RH, Klunk WE, Mathis CA, DeKosky ST, Morris JC (2006) [11C]PIB in a nondemented population: Potential antecedent marker of Alzheimer disease. *Neurology* **67**, 446-452.
- [29] Vlassenko AG, Mintun MA, Xiong C, Sheline YI, Goate AM, Benzinger TL, Morris JC (2011) Amyloid-beta plaque growth in cognitively normal adults: Longitudinal [11C]Pittsburgh compound B data. *Ann Neurol* **70**, 857-861.
- [30] Vlassenko AG, McCue L, Jasielec MS, Su Y, Gordon BA, Xiong C, Holtzman DM, Benzinger TL, Morris JC, Fagan AM (2016) Imaging and cerebrospinal fluid biomarkers in early preclinical Alzheimer disease. *Ann Neurol* **80**, 379-387.
- [31] Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, Busa E, Seidman LJ, Goldstein J, Kennedy D, Caviness V, Makris N, Rosen B, Dale AM (2004) Automatically parcellating the human cerebral cortex. *Cereb Cortex* **14**, 11-22.
- [32] DeLong ER, DeLong DM, Clarke-Pearson DL (1988) Comparing the areas under two or more correlated receiver operating characteristic curves: A nonparametric approach. *Biometrics* **44**, 837-845.
- [33] Dorey A, Perret-Liaudet A, Tholance Y, Fourier A, Quadrio I (2015) Cerebrospinal fluid Abeta40 improves the interpretation of Abeta42 concentration for diagnosing Alzheimer's disease. *Front Neurol* **6**, 247.
- [34] Lewczuk P, Kornhuber J (2016) Do we still need positron emission tomography for early Alzheimer's disease diagnosis? *Brain*, doi: 10.1093/brain/aww168
- [35] Palmqvist S, Mattsson N, Hansson O, Alzheimer's Disease Neuroimaging Initiative (2016) Cerebrospinal fluid analysis detects cerebral amyloid-beta accumulation earlier than positron emission tomography. *Brain* **139**(Pt 4), 1226-1236.
- [36] Janelidze S, Zetterberg H, Mattsson N, Palmqvist S, Vanderstichele H, Lindberg O, van Westen D, Stomrud E, Minthon L, Blennow K, Swedish BioFINDER study group, Hansson O (2016) CSF Abeta42/Abeta40 and Abeta42/Abeta38 ratios: Better diagnostic markers of Alzheimer disease. *Ann Clin Transl Neurol* **3**, 154-165.
- [37] Adamczuk K, Schaeverbeke J, Vanderstichele HM, Lilja J, Nelissen N, Van Laere K, Dupont P, Hilven K, Poesen K, Vandenbergh R (2015) Diagnostic value of cerebrospinal

- fluid Abeta ratios in preclinical Alzheimer's disease. *Alzheimers Res Ther* **7**, 75.
- [38] Racine AM, Kosciak RL, Nicholas CR, Clark LR, Okonkwo OC, Oh JM, Hillmer AT, Murali D, Barnhart TE, Betthauser TJ, Gallagher CL, Rowley HA, Dowling NM, Asthana S, Bendlin BB, Blennow K, Zetterberg H, Carlsson CM, Christian BT, Johnson SC (2016) Cerebrospinal fluid ratios with Abeta42 predict preclinical brain beta-amyloid accumulation. *Alzheimers Dement (Amst)* **2**, 27-38.
- [39] Cairns NJ, Ikonomic MD, Benzinger T, Storandt M, Fagan AM, Shah AR, Reinwald LT, Carter D, Felton A, Holtzman DM, Mintun MA, Klunk WE, Morris JC (2009) Absence of Pittsburgh compound B detection of cerebral amyloid beta in a patient with clinical, cognitive, and cerebrospinal fluid markers of Alzheimer disease: A case report. *Arch Neurol* **66**, 1557-1562.
- [40] Englund H, Degerman Gunnarsson M, Brundin RM, Hedlund M, Kilander L, Lannfelt L, Pettersson FE (2009) Oligomerization partially explains the lowering of Abeta42 in Alzheimer's disease cerebrospinal fluid. *Neurodegener Dis* **6**, 139-147.