

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2017

Genetic association of MMP10, MMP14, and MMP16 with dental caries

D. D. Lewis
University of Pittsburgh

J. R. Shaffer
University of Pittsburgh

E. Feingold
University of Pittsburgh

M. Cooper
University of Pittsburgh

M. M. Vanyukov
University of Pittsburgh

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Lewis, D. D.; Shaffer, J. R.; Feingold, E.; Cooper, M.; Vanyukov, M. M.; Maher, B. S.; Slayton, R. L.; Willing, M. C.; Reis, S. E.; McNeil, D. W.; Crout, R. J.; Weyant, R. J.; Levy, S. M.; Vieira, A. R.; and Marazita, M. L., "Genetic association of MMP10, MMP14, and MMP16 with dental caries." *International Journal of Dentistry*. 2017, 8465125. 1-7. (2017).
https://digitalcommons.wustl.edu/open_access_pubs/6792

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

D. D. Lewis, J. R. Shaffer, E. Feingold, M. Cooper, M. M. Vanyukov, B. S. Maher, R. L. Slayton, M. C. Willing, S. E. Reis, D. W. McNeil, R. J. Crout, R. J. Weyant, S. M. Levy, A. R. Vieira, and M. L. Marazita

Research Article

Genetic Association of *MMP10*, *MMP14*, and *MMP16* with Dental Caries

**D. D. Lewis,¹ J. R. Shaffer,¹ E. Feingold,^{1,2} M. Cooper,^{3,4} M. M. Vanyukov,^{1,5,6}
B. S. Maher,⁷ R. L. Slayton,⁸ M. C. Willing,⁹ S. E. Reis,^{10,11} D. W. McNeil,¹² R. J. Crout,¹³
R. J. Weyant,¹⁴ S. M. Levy,^{15,16} A. R. Vieira,^{3,4} and M. L. Marazita^{1,3,4,6,11}**

¹ Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA

² Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA

³ Center for Craniofacial and Dental Genetics, School of Dental Medicine, University of Pittsburgh, Pittsburgh, PA, USA

⁴ Department of Oral Biology, School of Dental Medicine, University of Pittsburgh, Pittsburgh, PA, USA

⁵ Department of Pharmaceutical Sciences, School of Pharmacy, University of Pittsburgh, Pittsburgh, PA, USA

⁶ Department of Psychiatry, School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA

⁷ Department of Mental Health, Johns Hopkins Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA

⁸ Department of Pediatric Dentistry, School of Dentistry, University of Washington, Seattle, WA, USA

⁹ Division of Genetics and Genomic Medicine, Department of Pediatrics, School of Medicine, Washington University at St. Louis, St. Louis, MO, USA

¹⁰ Department of Medicine, School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA

¹¹ Clinical and Translational Science Institute, School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA

¹² Psychology, Dental Practice and Rural Health, West Virginia University, Morgantown, WV, USA

¹³ Department of Periodontics, School of Dentistry, West Virginia University, Morgantown, WV, USA

¹⁴ Department of Dental Public Health and Information Management, School of Dental Medicine, University of Pittsburgh, Pittsburgh, PA, USA

¹⁵ Department of Preventive and Community Dentistry, University of Iowa College of Dentistry, Iowa City, IA, USA

¹⁶ Department of Epidemiology, University of Iowa College of Public Health, Iowa City, IA, USA

Correspondence should be addressed to D. D. Lewis; ddl8@pitt.edu

Received 22 May 2016; Accepted 6 February 2017; Published 28 February 2017

Academic Editor: Tommaso Lombardi

Copyright © 2017 D. D. Lewis et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Matrix metalloproteinases (MMPs), which degrade extracellular proteins as part of a variety of physiological processes, and their inhibitors have been implicated in the dental caries process. Here we investigated 28 genetic variants spanning the *MMP10*, *MMP14*, and *MMP16* genes to detect association with dental caries experience in 13 age- and race-stratified ($n = 3,587$) samples from 6 parent studies. Analyses were performed separately for each sample, and results were combined across samples by meta-analysis. Two SNPs (rs2046315 and rs10429371) upstream of *MMP16* were significantly associated with caries in an individual sample of white adults and via meta-analysis across 8 adult samples after gene-wise adjustment for multiple comparisons. Noteworthy is SNP rs2046315 ($p = 8.14 \times 10^{-8}$) association with caries in white adults. This SNP was originally nominated in a genome-wide-association study (GWAS) of dental caries in a sample of white adults and yielded associations in a subsequent GWAS of surface level caries in white adults as well. Therefore, in our study, we were able to recapture the association between rs2046315 and dental caries in white adults. Although we did not strengthen evidence that *MMPs* 10, 14, and 16 influence caries risk, *MMP16* is still a likely candidate gene to pursue.

1. Introduction

Despite significant improvements in oral health in the U.S, dental caries still remains the most prevalent chronic disease among children. The etiology of caries is multifactorial, involving a number of environmental factors, including microbial flora and fluoride exposure. Although these environmental factors substantially contribute to the disease itself, the impact of genetic factors plays a considerable role that has been long recognized and studied [1].

Evidence of genetic contribution to variation in liability to caries has been detected in studies showing heritability estimates between 40% and 60% [2–4]. Furthermore, over the past decade, there have been several studies that have nominated candidate genes based on their known biological functions in oral health such as genes involved in enamel formation [5–8], tooth development, [8, 9], taste preference [4, 10, 11], and host defense [12–15].

Matrix metalloproteinases (MMPs) are a well-studied multigene family that belongs to the metalloproteinase class of endopeptidases which are responsible for the remodeling and degradation of extracellular matrix molecules (ECM) [16]. ECM macromolecules are essential for maintaining a cellular environment for biological processes such as embryonic development, tissue remodeling, wound healing, and angiogenesis [17]. MMPs are involved in several diseases such as cancer [18], arthritis [19], tissue ulceration [20], periodontitis [21, 22], early tooth development [23, 24], and dental caries [16, 25, 26].

Studies have demonstrated that MMPs are involved in dental caries lesion progression by their presence in both dentin and saliva and their active role in the dentin matrix degradation [24, 27–29]. The initiation of caries is a dynamic process of which the mineral part of the dentin is dissolved (demineralization), exposing the organic matrix to degradation by bacterial and host derived enzymes such as MMPs [30]. Subsequent to demineralization, the breakdown of the collagenous organic matrix in the dentin occurs, which is necessary for caries formation [31]. Members of the metalloproteinase family such as *MMP10*, also referred to as stromelysin-2, are capable of degrading all components of the extracellular matrix [32]. Even more, a preliminary analysis of GWAS data suggested that variation in *MMP10* may increase the risk for dental caries [8, 33], thus providing compelling evidence for further investigation.

Several members of the MMP family have been observed in cells and tissues of the dentine-pulp complex and are thought to be involved in physiological processes during early tooth development [29, 34]. Correspondingly, *MMP14* is a membrane-type 1 matrix metalloproteinase (MT1-MMP), a membrane-bound member of the MMP gene family that has been identified to be expressed on the cell surface of ameloblasts and odontoblasts of the developing tooth [35]. Additionally, studies have demonstrated delayed tooth eruption in *mmp14* deficient mice [36] and have shown that *MMP14* plays a role in the activation of the enamel matrix protein *MMP20* [37]. Likewise, *MMP16*, a member of the type I transmembrane group as well, has been suggested to be involved in tooth development by regulating ameloblast

maturation and enamel formation [38]. The rationale for the selection of *MMP10*, *MMP14*, and *MMP16* was based on their biological role in tooth formation and plausible effects on caries etiology. Therefore, the aim of this study is to determine if variants spanning the regions of MMPs 10, 14, and 16 are associated with dental caries.

2. Methods

2.1. Samples and Data Collection. Study participants were drawn from 6 parent studies in this investigation: The Center for Oral Health Research in Appalachia cohort 1 [COHRA1, *N* = 1,769 [39]], Iowa Head Start [IHS, *N* = 64 [6]], Iowa Fluoride Study [IFS, *N* = 136 [40, 41]], Dental Strategies Concentrating on Risk Evaluation [Dental SCORE, *N* = 502 [42, 43]], the Dental Registry and DNA Repository [DRDR, *N* = 875 [8]], and the Center for Education and Drug Abuse Research [CEDAR, *N* = 241 [44]] (Table 1).

All study protocols were approved by the Institutional Review Boards of the corresponding universities. Details of the participant recruitment protocol and study design for each parent study have been previously reported and summarized and further described in the Appendix (see Supplementary Material available online at <https://doi.org/10.1155/2017/8465125>). The 6 parent studies were stratified into 13 samples by age and race (8 adults and 5 children samples).

Dental caries assessments were performed by trained dental professionals (dentists or dental hygienists) via intraoral examination. Intraclass correlation coefficient (ICC) analysis was applied to measure the intra- and interexaminer reproducibility of the caries assessment. High concordance rates were observed for both interexaminer reliability (ICC = 0.86–0.99) and intraexaminer reliability (ICC > 0.99) [39]. Each tooth was identified as either permanent or primary and each surface on each tooth was scored for evidence of decay from which traditional DMFT and dft indices were generated. DMFT was defined as the number of decayed, missing due to decay, or restored (filled) teeth of the permanent dentition, excluding third molars. Correspondingly, dft was defined as the number of decayed or restored teeth of the primary dentition.

3. Genotypes

Genotyping for a custom panel of single nucleotide polymorphisms (SNPs) was performed by the Center for Inherited Disease Research (CIDR) using the Illumina GoldenGate platform (San Diego, USA). The majority of this panel was chosen to follow up results from GWAS scans of dental caries. Additionally, we also included SNPs such as those in and near MMP genes, based on our specific interest in strong candidate genes. For this study, we investigated three MMP genes: *MMP10*, *MMP14*, and *MMP16* (Table 2). These genes were selected based on their known roles in tooth development [32, 35, 38, 45] and previous investigations with dental caries [33]. The following procedure was used to select 28 tag SNPs within these genes for genotyping. We began with a list of all SNPs within a gene and filtered out variants with minor allele frequencies less than 2% because

TABLE 1: Characteristics of the samples: mean (range) or percentage, %.

Sample	N	Female sex	Age, years	dft/DMFT ¹
Children				
COHRA whites	608	46.7%	7.3 (3.0–12.0)	2.3 (0–17)
COHRA blacks	81	46.9%	7.6 (3.2–11.8)	1.8 (0–8)
IHS whites	41	58.5%	4.1 (3.2–5.3)	6.3 (0–20)
IHS blacks	23	52.2%	4.3 (3.4–5.6)	5.7 (0–17)
IFS whites	136	48.5%	5.2 (4.4–6.8)	1.2 (0–16)
Adults				
COHRA whites	994	62.8%	34.3 (18.0–75.0)	10.5 (0–28)
COHRA blacks	86	70.9%	36.2 (18.2–60.8)	9.3 (9–28)
Dental SCORE whites	277	63.2%	64.0 (48.0–78.0)	16.4 (2–28)
Dental SCORE blacks	225	72.9%	61.6 (47.0–79.0)	14.8 (1–28)
DRDR whites	702	50.0%	43.0 (18.0–74.8)	16.6 (0–28)
DRDR blacks	173	57.8%	44.5 (18.0–74.4)	16.5 (0–28)
CEDAR whites	173	31.2%	20.4 (15.7–28.6)	5.4 (0–21)
CEDAR blacks	68	44.3%	20.2 (15.6–27.8)	6.4 (0–16)

¹ dft was the measure of caries experience of the primary dentition in children samples; DMFT was the measure of caries experience of the permanent dentition in adult samples.

TABLE 2: Genetic variants in MMPs 10, 14, and 16.

SNP	Chromosome	Position ^a	MAF (COHRA1) ^b	Base change	Location/functionality
<i>MMP10</i>					
rs7948454	11	102641196	0.06	C-T	Downstream
rs12272341	11	102644601	0.13	A-G	Intronic
rs470154	11	102647310	0.06	C-G	Intronic
rs17293607	11	102650389	0.12	C-T	Missense Gly65Arg
rs559518	11	102656079	0.36	A-G	Intronic
<i>MMP14</i>					
rs8003217	14	23304416	0.16	A-C	Downstream
rs762052	14	23308986	0.14	A-G	Intronic
rs10133740	14	23310131	0.15	C-T	Intronic
rs17243048	14	23311480	0.17	A-G	Intronic
rs12893368	14	23312208	0.17	C-G	Intronic
<i>MMP16</i>					
rs17718917	8	89030490	0.08	A-G	Intergenic
rs1477907	8	89033615	0.09	A-G	Intergenic
rs16876790	8	89035664	0.38	A-T	Intergenic
rs2664368	8	89045674	0.20	C-T	Intergenic
rs10103111	8	89075226	0.23	C-T	Intronic
rs1824717	8	89075979	0.49	A-G	Intronic
rs17719876	8	89083319	0.08	C-T	Intronic
rs2616487	8	89084284	0.34	A-G	Intronic
rs6469206	8	89084691	0.45	G-T	Intronic
rs7826929	8	89084837	0.13	A-G	Intronic
rs2054415	8	89087358	0.05	G-T	Intronic
rs1551893	8	89102366	0.07	A-T	Intronic
rs1382104	8	89103325	0.44	C-T	Intronic
rs17720688	8	89104241	0.13	C-T	Intronic
rs10089111	8	89119305	0.38	G-T	Intronic
rs16878625	8	89125990	0.09	C-T	Intronic
rs10429371	8	89993488	0.22	C-T	Intergenic
rs2046315	8	90211100	0.13	A-G	Intergenic

^aBased on Build 37, ^bMAF = minor allele frequency in the COHRA1 sample.




FIGURE 1: Genetic association in adults and children samples for 3 MMP genes. Genetic association in samples of adults (a) and children (b) for 3 MMP genes. Negative log₁₀ transformed p values are shown. Childhood samples: Center for Oral Health in Appalachia (COHRA1 (red)), Iowa Head Start (IHS (blue)), and Iowa Fluoride Study (IFS (purple)). Adult samples: Center for Oral Health in Appalachia (COHRA1 (red)), Dental Strategies Concentrating on Risk Evaluation (Dental SCORE (green)), Dental Registry and DNA Repository (DRDR (orange)), and Center for Education and Drug Abuse Research (CEDAR (yellow)). Circles represent white samples, and squares represent black samples. For children samples, white diamonds represent meta-analysis across all white childhood samples, and gray diamonds represent meta-analysis across all black and white childhood samples combined. Similarly, for adult samples, white diamonds represent meta-analysis across all white adult samples and gray diamonds represent meta-analysis across all black and white adult samples, combined. The dotted lines represent the p threshold after adjustment for the number of independent single nucleotide polymorphisms within a gene. The physical location and directions of the genes are denoted by the blue arrows.

testing low-frequency polymorphisms was not feasible in our cohorts. We also filtered out variants with low “designability scores,” which is a measure of the predicted success of genotyping on the GoldenGate genotyping platform; we used 0.8 as the threshold in order to be reasonably certain that SNPs could be successfully genotyped. For the SNPs passing these filters we chose tag SNPs by pruning out SNPs with redundant information defined by multiple $R^2 < 0.95$ in the reference CEU data (from the International HapMap

Project), using a sliding window of up to 50 SNPs advancing one SNP at a time. The goal of this procedure was to capture much of the information on the genetic variation in a gene while genotyping the fewest number of individual SNPs. Note that technical limitations of the GoldenGate genotyping technology required that all SNPs be at least 60 base pairs away from each other. Overall, this SNP selection process yielded a set of low-correlated variants that were included based on their informativeness rather than their putative

biological functions. Details regarding the design of the genotype panel are available elsewhere [46].

4. Statistical Analysis

Analyses of dental caries experience were performed separately in each sample for children 3–12 years of age for the primary dentition only (dft) and adults ≥ 18 years of age for the permanent dentition only (DMFT). Note, in children with mixed dentition, only the primary teeth present were included in the caries assessments because the goal was to identify risk factors influencing caries in the primary teeth in the child cohorts. The analyses were performed separately in self-reported non-Hispanics whites and blacks in order to guard against population stratification. Examinations of self-reported biological relationships among the participants were confirmed by genetic relatedness via 96 ancestry markers. Our CEDAR sample included adolescents > 15 years of age and for the purposes of this study was considered an adult sample. Linear regression analysis using PLINK software [47] was used to test genetic association between DMFT/dft and each SNP under the additive model while adjusting for age and sex. To guard against confounding due to admixture, we adjusted for the first 4 principal components for analyses of blacks. Further discussion of statistical analyses is presented in the Appendix.

Results were combined across samples using Stouffer's inverse variance weighted method of meta-analysis using METAL software [48]. This method is appropriate because it takes into consideration the nonrandom heterogeneity that is exhibited by the cohorts. Meta-analyses were performed for whites only and for all participants. Given the multiple comparisons, we used the method by Li and Ji [49] to determine the effective number of independent tests, which is less than or equal to the total number of tests due to linkage disequilibrium (LD; i.e., correlation) among SNPs. The threshold for declaring statistical significance was set to p value of 0.05, divided by the number of independent tests.

5. Results

Characteristics of the 13 samples are shown in Table 1. There were noticeable differences in dental caries experience, which were expected because of the differences in age and demography within the samples. Figure 1 shows the results of tests of genetic association for three matrix metalloproteinases genes: *MMP14*, *MMP16*, and *MMP10*. Negative \log_{10} -transformed p values are shown for tests of association in individual samples and combined. Detailed association results for select SNPs from these genes are shown in the Appendix Table.

In adults, the strongest evidence of genetic association was detected for rs2046315, upstream of *MMP16* for COHRA1 whites ($p = 8.14 \times 10^{-8}$). In addition, meta-analysis across 8 white adult samples for this SNP yielded significant association as well ($p = 0.002$). SNP rs10429371 was significantly associated with caries for COHRA1 white adults ($p = 0.0002$) and via meta-analysis across black and white adults and children samples combined ($p = 0.0005$). Though not meeting the threshold after gene-wise adjustment for

multiple comparisons, meta-analysis across white adults for rs10429371 showed nominal evidence of association ($p = 0.004$). Another convincing evidence of association that did not meet gene-wise threshold was for a SNP in *MMP10* for COHRA1 white adults (rs17293607, $p = 0.01$). There were no significant associations observed in any of the children samples.

6. Discussion

We investigated 28 SNPs in or near three *MMP* genes (*MMP10*, *MMP14*, and *MMP16*) for evidence of association with dental caries experience in 13 race- and age-stratified samples from 6 independent studies ($n = 3600$). Significant evidence of association was seen between two SNPs upstream of *MMP16* with dental caries in white adults and via meta-analysis across 8 adulthood samples.

Regarding the two SNPs upstream of *MMP16* associated in this study, noteworthy is SNP rs2046315. Our present work provided evidence that variant rs2046315, our strongest association ($p = 8.14 \times 10^{-8}$), is associated with dental caries in the permanent dentition for white adults. rs2046315 is located on chromosome 8q21.3 and is 870 kb upstream from *MMP16* and 560 kb away from the nearest gene *RIKP2* (receptor-interacting serine-threonine kinase 2). *RIKP2* has no known role in caries etiology, though it has been detected to be involved in apoptosis and is expressed in both deciduous and permanent tooth pulp cells [50]. Nearby rs10429371, also upstream of *MMP16*, showed evidence of association across multiple samples, though it was likewise driven by the COHRA1 white adults. *MMP16*, a member of the type I transmembrane proteins, has been suggested to be involved in early tooth development [38] and was relatively far (>100 kb) from our top SNP (rs2046315), and its role in caries etiology is unknown.

SNP rs2046315 was originally nominated in a GWAS scan of dental caries in a mostly overlapping sample of COHRA1 white adults [8]. In a subsequent GWAS of smooth surface caries in the permanent dentition, rs2046315 was also associated for increased caries susceptibility in COHRA1 adults ($p = 3.08 \times 10^{-8}$) [45]. Consistent with the previous findings, this SNP was suggestively associated with caries for the pit-and-fissures surface in the permanent dentition as well [45]. Therefore, in our study, we recapitulated the strong association observed in COHRA1 white adults but did not observe evidence of association in other samples for *MMP16*. Although *MMP16* is still a logical candidate to pursue, we did not strengthen evidence for its role in dental caries. In this study, we did not observe any associations for *MMP10* and *MMP14* in either adults or children.

Potential reasons why we did not find any significant associations in samples other than COHRA1 whites was that heterogeneity was observed across our samples, such as associations that were specific to individual samples (i.e., COHRA1 white adults). This could be due to differences between populations, such as interactions with environmental factors that are present in specific populations but not in the others. Overall, matrix metalloproteinases genes are involved in numerous physiological processes and diseases

such as dental caries. Though evidence of genetic association was not detected for *MMP10* and *MMP14*, results from our study suggest *MMP16* SNP rs2046315 may contribute to caries susceptibility in our individual sample of white adults. Even though rs2046315 was not proven to be a functional SNP, the multifactorial disorder of dental caries can involve several influential genes that potentially play a significant, but not an extensive role in the disease outcome. However, further investigation is needed to make this conclusion, and if truly associated, additional work is needed to determine the casual allele(s) and the mechanism impacting risk of caries.

Competing Interests

The authors declare no conflict of interests.

Acknowledgments

The authors would like to thank the participants and researchers of the 6 parent studies whose contributions made this work possible. Funding support was provided by the following National Institutes of Health Grants U01-DE018903, R01-DE014899, R01-DE009551, R01-DE012101, R01-DE018914, R03-DE024264, P50-DA005605, and R01-DA019157, as well as the National Science Foundation/Department of Defense Grant DBI-1263020. The Dental Registry and DNA Repository is supported by the University of Pittsburgh School of Dental Medicine. The Dental SCORE sample is partially supported by the Commonwealth of Pennsylvania Department of Health Grant ME-02-384.

References

- [1] G. C. Townsend, M. J. Aldred, and P. Mark Bartold, "Genetic aspects of dental disorders," *Australian Dental Journal*, vol. 43, no. 4, pp. 269–286, 1998.
- [2] J. C. Boraas, L. B. Messer, and M. J. Till, "A genetic contribution to dental caries, occlusion, and morphology as demonstrated by twins reared apart," *Journal of Dental Research*, vol. 67, no. 9, pp. 1150–1155, 1988.
- [3] X. Wang, J. R. Shaffer, R. J. Weyant et al., "Genes and their effects on dental caries may differ between primary and permanent dentitions," *Caries Research*, vol. 44, no. 3, pp. 277–284, 2010.
- [4] S. Wendell, X. Wang, M. Brown et al., "Taste genes associated with dental caries," *Journal of Dental Research*, vol. 89, no. 11, pp. 1198–1202, 2010.
- [5] K. Deeley, A. Letra, E. K. Rose et al., "Possible association of amelogenin to high caries experience in a Guatemalan-Mayan population," *Caries Research*, vol. 42, no. 1, pp. 8–13, 2008.
- [6] R. L. Slayton, M. E. Cooper, and M. L. Marazita, "Tuftelin, mutans streptococci, and dental caries susceptibility," *Journal of Dental Research*, vol. 84, no. 8, pp. 711–714, 2005.
- [7] A. Patir, F. Seymen, M. Yildirim et al., "Enamel formation genes are associated with high caries experience in Turkish children," *Caries Research*, vol. 42, no. 5, pp. 394–400, 2008.
- [8] X. Wang, J. R. Shaffer, Z. Zeng et al., "Genome-wide association scan of dental caries in the permanent dentition," *BMC Oral Health*, vol. 12, no. 1, article no. 57, 2012.
- [9] P. N. Tannure, E. C. Küchler, A. Lips et al., "Genetic variation in MMP20 contributes to higher caries experience," *Journal of Dentistry*, vol. 40, no. 5, pp. 381–386, 2012.
- [10] R. Pidamale, B. Sowmya, A. Thomas et al., "Association between early childhood caries, streptococcus mutans level and genetic sensitivity levels to the bitter taste of, 6-N propylthiouracil among the children below 71 months of age," *Dental Research Journal (Isfahan)*, vol. 9, no. 6, pp. 730–734, 2012.
- [11] G. V. Kulkarni, T. Chng, K. M. Eny, D. Nielsen, C. Wessman, and A. El-Sohemy, "Association of GLUT2 and TAS1R2 genotypes with risk for dental caries," *Caries Research*, vol. 47, no. 3, pp. 219–225, 2013.
- [12] R. T. Acton, A. P. Dasanayake, R. A. Harrison et al., "Associations of MHC genes with levels of caries-inducing organisms and caries severity in African-American women," *Human Immunology*, vol. 60, no. 10, pp. 984–989, 1999.
- [13] A. Ozturk, P. Famili, and A. R. Vieira, "The antimicrobial peptide DEFB1 is associated with caries," *Journal of Dental Research*, vol. 89, no. 6, pp. 631–636, 2010.
- [14] N. Valarini, S. M. MacIel, S. K. Moura, and R. C. Poli-Frederico, "Association of dental caries with HLA class II allele in Brazilian adolescents," *Caries Research*, vol. 46, no. 6, pp. 530–535, 2012.
- [15] J. Briseño-Ruiz, T. Shimizu, K. Deeley et al., "Role of TRAV locus in low caries experience," *Human Genetics*, vol. 132, no. 9, pp. 1015–1025, 2013.
- [16] C. Chaussain-Miller, F. Fioretti, M. Goldberg, and S. Menashi, "The role of matrix metalloproteinases (MMPs) in human caries," *Journal of Dental Research*, vol. 85, no. 1, pp. 22–32, 2006.
- [17] R. Visse and H. Nagase, "Matrix metalloproteinases and tissue inhibitors of metalloproteinases: structure, function, and biochemistry," *Circulation Research*, vol. 92, no. 8, pp. 827–839, 2003.
- [18] C. Gialeli, A. D. Theocharis, and N. K. Karamanos, "Roles of matrix metalloproteinases in cancer progression and their pharmacological targeting," *FEBS Journal*, vol. 278, no. 1, pp. 16–27, 2011.
- [19] P. S. Burrage, K. S. Mix, and C. E. Brinckerhoff, "Matrix metalloproteinases: role in arthritis," *Frontiers in Bioscience*, vol. 11, no. 1, pp. 529–543, 2006.
- [20] B. C. Nwomeh, H.-X. Liang, I. K. Cohen, and D. R. Yager, "MMP-8 is the predominant collagenase in healing wounds and nonhealing ulcers," *Journal of Surgical Research*, vol. 81, no. 2, pp. 189–195, 1999.
- [21] M. Mäkelä, T. Salo, V.-J. Uitto, and H. Larjava, "Matrix metalloproteinases (MMP-2 and MMP-9) of the oral cavity: cellular origin and relationship to periodontal status," *Journal of Dental Research*, vol. 73, no. 8, pp. 1397–1406, 1994.
- [22] H. Birkedal-Hansen, "Role of matrix metalloproteinases in human periodontal diseases," *Journal of Periodontology*, vol. 64, no. 5, pp. 474–484, 1993.
- [23] H. Palosaari, J. Wahlgren, M. Larinas et al., "The expression of mmp-8 in human odontoblasts and dental pulp cells is down-regulated by tgf- β 1," *Journal of Dental Research*, vol. 79, no. 1, pp. 77–84, 2000.
- [24] L. Tjäderhane, T. Salo, H. Larjava, M. Larinas, and C. M. Overall, "A novel organ culture method to study the function of human odontoblasts in vitro: gelatinase expression by odontoblasts is differentially regulated by TGF- β 1," *Journal of Dental Research*, vol. 77, no. 7, pp. 1486–1496, 1998.
- [25] A. Mazzoni, L. Tjäderhane, V. Checchi et al., "Role of dentin MMPs in caries progression and bond stability," *Journal of Dental Research*, vol. 94, no. 2, pp. 241–251, 2015.

- [26] T. Sorsa, L. Tjäderhane, and T. Salo, "Matrix metalloproteinases (MMPs) in oral diseases," *Oral Diseases*, vol. 10, no. 6, pp. 311–318, 2004.
- [27] L. Tjäderhane, M. Sulkala, T. Sorsa, O. Teronen, M. Larmas, and T. Salo, "The effect of MMP inhibitor metostat on fissure caries progression in rats," *Annals of the New York Academy of Sciences*, vol. 878, pp. 686–688, 1999.
- [28] T. Sorsa, Y. L. Ding, T. Ingman et al., "Cellular source, activation and inhibition of dental plaque collagenase," *Journal of Clinical Periodontology*, vol. 22, no. 9, pp. 709–717, 1995.
- [29] S. Martin-De Las Heras, A. Valenzuela, and C. M. Overall, "The matrix metalloproteinase gelatinase A in human dentine," *Archives of Oral Biology*, vol. 45, no. 9, pp. 757–765, 2000.
- [30] M. Toledano, M. Yamauti, E. Osorio, and R. Osorio, "Zinc-inhibited MMP-mediated collagen degradation after different dentine demineralization procedures," *Caries Research*, vol. 46, no. 3, pp. 201–207, 2012.
- [31] R. M. Frank, P. Steuer, and J. Hemmerle, "Ultrastructural study on human root caries," *Caries Research*, vol. 23, no. 4, pp. 209–217, 1989.
- [32] V.-M. Kähäri and U. Saarialho-Kere, "Matrix metalloproteinases and their inhibitors in tumour growth and invasion," *Annals of Medicine*, vol. 31, no. 1, pp. 34–45, 1999.
- [33] S. B. Commander, *Investigation of Genetic Association of MMP10 with Dental Caries*, Univeristy of Pittsburgh, Pittsburgh, Pa, USA, 2011.
- [34] R. Hall, D. Septier, G. Embery, and M. Goldberg, "Stromelysin-1 (MMP-3) in forming enamel and predentine in rat incisor—coordinated distribution with proteoglycans suggests a functional role," *Histochemical Journal*, vol. 31, no. 12, pp. 761–770, 1999.
- [35] C. Caron, J. Xue, and J. D. Bartlett, "Expression and localization of membrane type 1 matrix metalloproteinase in tooth tissues," *Matrix Biology*, vol. 17, no. 7, pp. 501–511, 1998.
- [36] J. D. Bartlett, Z. Zhou, Z. Skobe, J. M. Dobeck, and K. Trygvason, "Delayed tooth eruption in membrane type-1 matrix metalloproteinase deficient mice," *Connective Tissue Research*, vol. 44, no. 1, pp. 300–304, 2003.
- [37] J. D. Bartlett, "Dental enamel development: proteinases and their enamel matrix substrates," *ISRN Dentistry*, vol. 2013, Article ID 684607, 24 pages, 2013.
- [38] C. Liu, Y. Niu, X. Zhou et al., "Cell cycle control, DNA damage repair, and apoptosis-related pathways control pre-ameloblasts differentiation during tooth development," *BMC Genomics*, vol. 16, no. 1, article no. 592, 2015.
- [39] D. E. Polk, R. J. Weyant, R. J. Crout et al., "Study protocol of the Center for Oral Health Research in Appalachia (COHRA) etiology study," *BMC Oral Health*, vol. 8, no. 1, article 18, 2008.
- [40] X. Wang, M. C. Willing, M. L. Marazita et al., "Genetic and environmental factors associated with dental caries in children: The Iowa Fluoride Study," *Caries Research*, vol. 46, no. 3, pp. 177–184, 2012.
- [41] T. A. Marshall, J. M. Eichenberger Gilmore, B. Broffitt, P. J. Stumbo, and S. M. Levy, "Relative validity of the Iowa fluoride study targeted nutrient semi-quantitative questionnaire and the block kids' food questionnaire for estimating beverage, calcium, and vitamin d intakes by children," *Journal of the American Dietetic Association*, vol. 108, no. 3, pp. 465–472, 2008.
- [42] A. N. Aiyer, K. E. Kip, O. C. Marroquin, S. R. Mulukutla, D. Edmundowicz, and S. E. Reis, "Racial differences in coronary artery calcification are not attributed to differences in lipoprotein particle sizes: the Heart Strategies Concentrating on Risk Evaluation (Heart SCORE) Study," *American Heart Journal*, vol. 153, no. 2, pp. 328–334, 2007.
- [43] A. N. Aiyer, K. E. Kip, S. R. Mulukutla, O. C. Marroquin, L. Hipps Jr., and S. E. Reis, "Predictors of significant short-term increases in blood pressure in a community-based population," *The American Journal of Medicine*, vol. 120, no. 11, pp. 960–967, 2007.
- [44] M. M. Vanyukov, B. S. Maher, B. Devlin et al., "Haplotypes of the monoamine oxidase genes and the risk for substance use disorders," *American Journal of Medical Genetics—Neuropsychiatric Genetics*, vol. 125, no. 1, pp. 120–125, 2004.
- [45] Z. Zeng, J. R. Shaffer, X. Wang et al., "Genome-wide association studies of pit-and-fissure- and smooth-surface caries in permanent dentition," *Journal of Dental Research*, vol. 92, no. 5, pp. 432–437, 2013.
- [46] B. O. C. Stanley, E. Feingold, M. Cooper et al., "Genetic association of MPPED2 and ACTN2 with dental caries," *Journal of Dental Research*, vol. 93, no. 7, pp. 626–632, 2014.
- [47] S. Purcell, B. Neale, K. Todd-Brown et al., "PLINK: a tool set for whole-genome association and population-based linkage analyses," *The American Journal of Human Genetics*, vol. 81, no. 3, pp. 559–575, 2007.
- [48] C. J. Willer, Y. Li, and G. R. Abecasis, "METAL: fast and efficient meta-analysis of genomewide association scans," *Bioinformatics*, vol. 26, no. 17, pp. 2190–2191, 2010.
- [49] J. Li and L. Ji, "Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix," *Heredity*, vol. 95, no. 3, pp. 221–227, 2005.
- [50] Y. Okai, K. Harada, K. Ohura, and K. Arita, "Study on apoptosis in human deciduous tooth pulp cells," *Journal of Hard Tissue Biology*, vol. 21, no. 4, pp. 413–420, 2012.

