

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2018

Draft genome sequence of a white spot syndrome virus isolate obtained in Ecuador

Leda Restrepo

Universidad de Los Andes - Colombia

Alejandro Reyes

Washington University School of Medicine in St. Louis

Leandro Bajaña

Centro Nacional de Acuicultura e Investigaciones Marinas, CENAIM

Irma Betancourt

Centro Nacional de Acuicultura e Investigaciones Marinas, CENAIM

Bonny Bayot

Centro Nacional de Acuicultura e Investigaciones Marinas, CENAIM

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Restrepo, Leda; Reyes, Alejandro; Bajaña, Leandro; Betancourt, Irma; and Bayot, Bonny, "Draft genome sequence of a white spot syndrome virus isolate obtained in Ecuador." *Genome Announcements*. 6, 26. e00605-18. (2018).

https://digitalcommons.wustl.edu/open_access_pubs/7043

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.



Draft Genome Sequence of a White Spot Syndrome Virus Isolate Obtained in Ecuador

 Leda Restrepo,^{a,b,c} Alejandro Reyes,^{b,c,d} Leandro Bajaña,^a Irma Betancourt,^a Bonny Bayot^{a,e}

^aEscuela Superior Politécnica del Litoral (ESPOL), Centro Nacional de Acuicultura e Investigaciones Marinas, CENAIM, Guayaquil, Ecuador

^bDepartment of Biological Sciences, Universidad de los Andes, Bogotá, Colombia

^cMax Planck Tandem Group in Computational Biology, Universidad de los Andes, Bogotá, Colombia

^dCenter for Genome Sciences and Systems Biology, Department of Pathology and Immunology, Washington University in Saint Louis, St. Louis, Missouri, USA

^eEscuela Superior Politécnica del Litoral (ESPOL), Facultad de Ingeniería Marítima, Ciencias Biológicas, Oceánicas y Recursos Naturales, FIMCBOR, Guayaquil, Ecuador

ABSTRACT White spot syndrome virus (WSSV) is the most devastating viral disease affecting cultivated shrimp around the world. Currently, there is no reported genetic information on WSSV affecting *Penaeus vannamei* in Ecuador. Therefore, we determined the genome of a WSSV isolate from shrimp and compared it with the genomes of isolates from other geographical locations.

Aquaculture is one of the most important economic activities in Ecuador, with *Penaeus vannamei* shrimp being the second highest nonoil export product, accounting for 95% of the country's total aquaculture production (1). The presence of the pathogen white spot syndrome virus (WSSV) in Ecuador was confirmed in 1999 (2). In subsequent years, the disease became an epidemic, resulting in the virus being labeled as the most devastating pathogen in the aquacultural history of shrimp farming. A monitoring effort was carried out along the Ecuadorian coast to evaluate the spread of the virus. Genotyping of the circulating WSSV types was performed to characterize the strain diversity within the country, leading to the sequencing of the complete genome of the most prevalent strain in this territory.

A whole-genome sequencing (WGS) strategy was performed on an Illumina HiSeq sequencer using DNA extracted from infected shrimp containing strain WSSV-EC-15098. In total, 20,649,280 paired-end 2 × 101-bp reads were obtained, which generated 2.085 Gb of total sequenced bases. The reads were mapped to the reference genomes of WSSV strain CN01 (3) (GenBank accession no. KT995472) and WSSV K-LV1 (4) (GenBank accession no. JX515788) using Bowtie 2 (5). They were then used for *de novo* assembly with SPAdes (6). The initial assembly generated 13 contigs. Reads mapping to the edges of the contigs were extracted and assembled using CAP3 (7), and the resulting contigs were used to close the assembly, obtaining a single contig of 288,997 bp. Open reading frames (ORFs) were predicted using the GeneMarkS software, with default parameters (8). The predicted protein sequences were searched against the GenBank database and the Clusters of Orthologous Groups (COGs) protein database using BLASTp (9).

The draft genome of WSSV-EC-15098 consists of 288,997 bp, with a mean G+C content of 40.92%. A total of 162 coding sequences (CDS) were predicted. The genomic architectures were compared with strains from China (3), Thailand (10), South Korea (4), and Mexico (11) using Mauve (12). A complete synteny among the assembled genomes was observed. Two major deletions were observed in the WSSV-EC-15098 genome. The deletions were verified by paired-end reads located at each side of the deletion and

Received 28 May 2018 Accepted 30 May 2018 Published 28 June 2018

Citation Restrepo L, Reyes A, Bajaña L, Betancourt I, Bayot B. 2018. Draft genome sequence of a white spot syndrome virus isolate obtained in Ecuador. *Genome Announc* 6:e00605-18. <https://doi.org/10.1128/genomeA.00605-18>.

Copyright © 2018 Restrepo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Leda Restrepo, ledarestrepoc@gmail.com.

mapping to the reference genome without any read mapping to any part of the deleted segments. The deletions are a 3.24-kb fragment containing the ORFs VP22 and VP25, all annotated as envelope proteins, and a 17.04-kb fragment including the ORF VP35, annotated as a capsid protein. The deletion of proteins related to the envelope is very interesting due to its potential role in modifying the way the virus interacts with the host in terms of recognition of the host cell and by the immune machinery. Additionally, it has been observed that strain WSSV-EC-15098 is widely spread but causes the lowest shrimp mortality rate.

The average nucleotide identity (ANI) between WSSV-EC-15098 and other genomes is between 99.67% and 99.94%, with the WSSV-China genome being the most closely related and WSSV-Korea genome being the most distantly related. The implementation of molecular epidemiology for the control of diseases could help in the development of strategies for the prevention and control of emerging and reemerging aquaculture-related diseases.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [MH090824](https://www.ncbi.nlm.nih.gov/nuclseq/MH090824).

ACKNOWLEDGMENTS

We thank Ana Elena Ayong for her technical help in this study and Stanislaus Sonnenholzner for his kind reviewing of the manuscript and suggestions. Additionally, we acknowledge the High-Performance Computing Service at the Universidad de los Andes in Bogotá, Colombia, (<http://hpc.uniandes.edu.co>) for providing high-performance computing (HPC) resources that have contributed to the research results reported in this work.

This work was funded by the Secretaria de Educación Superior, Ciencia, Tecnología e Innovación (SENESCYT) in the framework of the PIC-14-CENAIM-003 project “Desarrollo e implementación de métodos de control y prevención de enfermedades en especies acuáticas de uso comercial y uso potencial en maricultura y repoblación.”

REFERENCES

- Ministerio de Comercio Exterior. 2017. Pro Ecuador instituto de promoción de exportaciones e inversiones. Ministerio de Comercio Exterior, Guayaquil, Ecuador. <https://www.americaeconomia.com/economia-mercados/comercio/exportaciones-no-petroleras-de-ecuador-crecieron-77-y-suman-us-10093m>. (In Spanish.)
- Calderón J, Bayot B, Betancourt I, de Graindorge V. 1999. Monitoreo del virus de la mancha blanca en Ecuador. *El Mundo Acuicola* 5:211–214.
- Yang F, He J, Lin X, Li Q, Pan D, Zhang X, Xu X. 2001. Complete genome sequence of the shrimp white spot bacilliform virus. *J Virol* 75:11811–11820. <https://doi.org/10.1128/JVI.75.23.11811-11820.2001>.
- Chai C, Yoon J, Lee Y, Kim Y, Choi T. 2013. Analysis of the complete nucleotide sequence of a white spot syndrome virus isolated from Pacific white shrimp. *J Microbiol* 51:695–699. <https://doi.org/10.1007/s12275-013-3171-0>.
- Langmead B, Salzberg S. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
- Bankevich A, Nurk S, Antipov D, Gurevich A, Dvorkin M, Kulikov A, Lesin V, Nikolenko S, Pham S, Prjibelski A, Pyshkin A, Sirotkin A, Vyahhi N, Tesler G, Alekseyev M, Pevzner P. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Huang X, Madan A. 1999. CAP3: a DNA sequence assembly program. *Genome Res* 9:868–877. <https://doi.org/10.1101/gr.9.9.868>.
- Besemer J, Borodovsky M. 2005. GeneMark: Web software for gene finding in prokaryotes, eukaryotes and viruses. *Nucleic Acids Res* 33:W451–W454. <https://doi.org/10.1093/nar/gki487>.
- Tatusov R, Galperin M, Natale D, Koonin E. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* 28:33–36. <https://doi.org/10.1093/nar/28.1.33>.
- Van Hulten M, Witteveldt J, Peters S, Kloosterboer N, Tarchini R, Fiers M, Sandbrink H, Lankhorst R, Vlask J. 2001. The white spot syndrome virus DNA genome sequence. *Virology* 286:7–22. <https://doi.org/10.1006/viro.2001.1002>.
- Rodriguez-Anaya LZ, Gonzalez-Galaviza JR, Casillas-Hernandez R, Lares-Villaa F, Estrada K, Ibarra-Gameza JC, Sanchez-Flores A. 2016. Draft genome sequence of white spot syndrome virus isolated from cultured *Litopenaeus vannamei* in Mexico. *Genome Announc* 4:e01674–15. <https://doi.org/10.1128/genomeA.01674-15>.
- Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.