

2018

Epidemiology of lumbar punctures in hospitalized patients in the United States

Adrienne Vickers

University of South Alabama

John P. Donnelly

University of Alabama, Birmingham

Justin Xavier Moore

Washington University School of Medicine in St. Louis

Scott R. Barnum

CNine Biosolutions, Birmingham

Theresa N. Schein

CNine Biosolutions, Birmingham

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Recommended Citation

Vickers, Adrienne; Donnelly, John P.; Moore, Justin Xavier; Barnum, Scott R.; Schein, Theresa N.; and Wang, Henry E., "Epidemiology of lumbar punctures in hospitalized patients in the United States." *PLoS One*.13,12. e0208622. (2018).
https://digitalcommons.wustl.edu/open_access_pubs/7464

Authors

Adrienne Vickers, John P. Donnelly, Justin Xavier Moore, Scott R. Barnum, Theresa N. Schein, and Henry E. Wang

RESEARCH ARTICLE

Epidemiology of lumbar punctures in hospitalized patients in the United States

Adrienne Vickers¹, John P. Donnelly^{2,3}, Justin Xavier Moore^{2,4}, Scott R. Barnum⁵, Theresa N. Schein⁵, Henry E. Wang^{2,6*}

1 University of South Alabama School of Medicine, Mobile, Alabama, United States of America, **2** Department of Emergency Medicine, University of Alabama School of Medicine, Birmingham, Alabama, United States of America, **3** Department of Learning Health Sciences, University of Michigan Medical School, Ann Arbor, MI, United States of America, **4** Department of Surgery, Washington University School of Medicine, Saint Louis, MO, United States of America, **5** CNine Biosolutions, LLC., Birmingham, Alabama, United States of America, **6** Department of Emergency Medicine, University of Texas Health Science Center at Houston, Houston, Texas, United States of America

* henry.e.wang@uth.tmc.edu



OPEN ACCESS

Citation: Vickers A, Donnelly JP, Moore JX, Barnum SR, Schein TN, Wang HE (2018) Epidemiology of lumbar punctures in hospitalized patients in the United States. PLoS ONE 13(12): e0208622. <https://doi.org/10.1371/journal.pone.0208622>

Editor: Pedro Gonzalez-Alegre, University of Pennsylvania Perelman School of Medicine, UNITED STATES

Received: June 23, 2018

Accepted: November 20, 2018

Published: December 13, 2018

Copyright: © 2018 Vickers et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data underlying this study are from the Nationwide Emergency Department Sample (NEDS), which is publicly available at: <http://www.hcup-us.ahrq.gov/nedsoverview.jsp>. The data set used in this study is from 2010 and is available at the following link: <https://www.distributor.hcup-us.ahrq.gov/Databases.aspx>. The authors did not have special access privileges.

Abstract

Objectives

Lumbar puncture (LP) is an important technique for assessing and treating neurological symptoms. The objective of this study was to describe the characteristics of diagnostic lumbar punctures performed on hospitalized patients in the United States.

Methods

We analyzed data from the 2010 National Inpatient Sample (NIS) and the National Emergency Department Survey (NEDS). We included patients treated in the Emergency Department (ED) as well as those admitted to an inpatient bed through the ED. We identified patients undergoing LPs from ICD-9 procedural code 03.31 and CPT code 62270. We generated nationally weighted estimates of the total number of LPs. We also assessed patient and hospital characteristics of cases undergoing LP.

Results

Of an estimated 135 million hospitalizations (ED + admission, or ED only), there were an estimated 362,718 LPs (331,248–394,188), including 273,612 (251,850–295,375) among adults and 89,106 (71,870–106,342) among children (<18 years old). Of the 362,718 LPs, 136,764 (122,117–151,410) were performed in the ED without admission. The most common conditions associated with LP among children were fever of unknown origin, meningitis, seizures and other perinatal conditions. The most common conditions associated with LP among adults were headache and meningitis.

Conclusions

Lumbar Puncture remains an important procedure for diagnostic and therapeutic uses in United States Hospitals.

Funding: Supported by a grant from the Emergency Medicine Foundation. CNine Biosolutions provided support in the form of salaries for authors SB and TS but had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

Competing interests: We have the following interests. Scott R Barnum and Theresa N. Schein are employed by CNine Biosolutions, LLC. There are no patents, products in development or marketed products to declare. This does not alter our adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors.

Introduction

The lumbar puncture was first introduced by Heinrich Quinke in 1891 to relieve elevated intracranial pressure in cases of meningitis.[1, 2] Since its innovation, the role of the LP has expanded to encompass diagnostic roles, including the identification of life-threatening conditions such as meningitis and hydrocephalus. The role of the LP is particularly important in the diagnosis of acute bacterial meningitis, which requires rapid diagnosis to prevent morbidity and mortality.[3, 4] Although more recent advances in imaging enhanced the diagnosis of neurologic emergencies, these modalities may miss conditions identifiable from cerebrospinal fluid analysis. For example, computed tomography imaging cannot capture microscopic bleeding from cerebral aneurysms. More importantly, radiologic imaging cannot prove the presence or absence of meningitis. Thus, the procurement of CSF from an LP remains an important tool for diagnosing many important neurologic emergencies.

However, the LP is an invasive procedure and has important pitfalls. A meta-analysis found that 1 in 3 patients undergoing the procedure may develop a post-LP headache.[5] Cerebral herniation occurs in 5% of patients with acute bacterial meningitis, with a strong temporal association with the performance of the LP even after a negative CT scan.[6, 7] Anticoagulant use (including novel oral anticoagulants) is a contraindication to LP, and the use of these agents in routine outpatient care is increasing.[8–10] Additionally, successfully performing a LP may be more difficult in obese patients and those with abnormal spinal anatomy.[11]

There have been no national descriptions of the epidemiology of LPs in United States (US) Emergency Departments (EDs); for example, the number of patients undergoing diagnostic LP, and their characteristics, diagnoses and outcomes. An awareness of these patterns is important for several reasons. The total number of cases would indicate equipment and training resources needed to adequately provide this procedure. These figures would indicate the number of patients vulnerable to LP adverse events, aiding in the planning for rescue therapies and opportunities for more advanced techniques such as ultrasound or fluoroscopically-guided LP. Novel biomarkers have been developed to help differentiate bacterial from viral CSF infections; epidemiologic descriptions of LPs could indicate the number of patients potentially benefitting from these novel assays.

In this study, we sought to describe the epidemiology of diagnostic LPs among Emergency Department-treated patients the United States.

Methods

Study design

We analyzed data from the Nationwide Emergency Department Sample (NEDS). This study was classified as not human subjects research by the Institutional Review Board of the University of Alabama at Birmingham.

Survey content and administration

A component of the Healthcare Cost Utilization Project (HCUP), NEDS is the largest all-payer ED database in the US. NEDS combines national information from State Inpatient Databases (SIDs) and State Emergency Department Databases (SEDDs) and hence reflects patients treated in the ED and discharged or admitted to the hospital through an ED visit.[12] NEDS is an approximately 20% stratified sample of US hospital-based EDs entailing data from 945 hospitals in 33 states.[13] The NEDS data correspond to roughly 135 million ED visits annually. The NEDS database contains more than 100 clinical and nonclinical variables for each hospital visit or stay, including ED diagnoses, procedures, injury information, admission

and discharge status, patient demographics, payer source, ED charges, and hospital characteristics.

NEDS encompasses ED-treated patients, including a) patients treated in the ED and subsequently admitted to the hospital, and b) patients treated in and subsequently discharged or transferred. We used NEDS data for the year 2010 because it provides the largest sample population and, therefore, is most appropriate for estimating total regional and national populations.

Selection of patients

We identified all NEDS visits associated with the performance of a diagnostic LP. We identified LPs using ICD-9-CM procedure code 03.31 and Current Procedural Terminology code and Healthcare Common Procedure Coding System (CPT/HCPCS) procedure code 62270. We focused on diagnostic LPs because of our interest in patients treated in the ED for neurologic emergencies such as meningitis or subarachnoid hemorrhage. We excluded therapeutic LPs (CPT 62272) because these interventions are less common among ED-treated patients and are usually limited to patients with chronic neurologic conditions such as hydrocephalus.

It is important to note that the structure of the NEDS did not indicate the hospital location of procedures. Therefore, for patients initially treated in the ED and subsequently admitted, we could not differentiate if the LP was performed in the ED or in the inpatient setting. In contrast, LPs performed on patients who were treated only in ED could be more clearly attributed to the ED setting.

Data analysis

Using descriptive statistical techniques, we calculated the number and incidence of LPs performed in the United States Emergency Departments in 2010. Following standard NEDS weighting procedures, we produced nationally weighted estimates of LPs and their corresponding 95% confidence intervals. For age-related estimates, we categorized patients as children (<18 years) or adults (\geq 18 years). In sensitivity analyses we further stratified the pediatric population to the age categories 0 to 5 years, 6 to 10 years, and 11 to 17 years.

We characterized all hospitalized patients by age, sex, and region of residence, along with household income quartile based on the residential zip code. We further differentiated LP patients by their primary diagnosis, obtained through classification based on Clinical Classifications Software (CCS) code at discharge. We characterized hospitals according to geographic region, teaching status, trauma center designation, and population setting (urban versus rural, defined by population estimates). In order to account for any seasonal variation in ED use, we also characterized ED visits by season. ED patient outcomes were categorized as routine discharge, inpatient admission, death, transferred to another hospital, or "other", which includes transfers to nursing homes or other locations and left against medical advice. We also classified inpatient outcomes for patients admitted to the hospital of ED presentation using similar categories.

All analyses were performed using Stata v. 14.1 (Stata, Inc., College Station, TX).

Results

During the study period, in the NEDS data set there were an estimated 135 million ED visits in the United States. There were a total of 362,718 LPs (incidence 2.7 per 1,000 ED visits), including 273,612 (75%) among adults and 89,106 (25%) among children. (Table 1) Among children, the majority were performed on patient \leq 5 years old. Among the performed LPs, 136,764 (38%) were known to have occurred in the ED.

Table 1. National estimates of lumbar punctures performed for Emergency Department-treated patients, 2010.

Patient Group	Emergency Department Admitted Inpatients ¹		Emergency Department Only Patients ²	
	No. of Raw Observations	Estimated Number of Lumbar Punctures (95% CI)	No. of Raw Observations	Estimated Number of Lumbar Punctures (95% CI)
All Patients	79,697	362,718 (331,248–394,188)	30,045	136,764 (122,117–151,410)
Adult Patients (Age ≥18 years)	60,016	273,612 (251,850–295,375)	22,913	105,015 (93,268–116,762)
Children				
0–5 years	14,360	64,849 (50,703–78,995)	4,104	18,096 (15,047–21,144)
6–10 years	1,707	7,762 (6,321–9,203)	833	3,736 (3,181–4,294)
11–17 years	3,614	16,495 (14,357–18,633)	2,195	9,915 (8,723–11,108)

¹Includes patients who were treated in the ED and subsequently admitted to the hospital.

²Includes only patients who treated in and subsequently discharged from the ED

<https://doi.org/10.1371/journal.pone.0208622.t001>

Among children, patients undergoing LP were more likely to be male than female. (Table 2) In contrast, gender distributions were similar for adults. For both age groups, LPs were more common for urban than rural patients. The incidence of lumbar puncture was also higher among patients from lower income areas. Among adults, LPs were most common in hospitals in the south census region. Though this result was not seen for all children, it was appreciated in our sensitivity analysis for younger children ≤5 years old. (S1 Appendix) Among all children, LPs were more common in trauma centers. Regardless of age, LPs were more common in urban and teaching hospitals and more common during summer months.

Among children, fever of unknown origin, other perinatal conditions, meningitis, and seizures were the more common conditions associated with LP. (Table 3) Among adults, headache and meningitis accounted for more than half of the hospitalizations associated with an LP. Most children receiving LPs were discharged after being admitted, while a large majority of children not receiving LPs were discharged home from the ED. (Table 4) However, similar proportions of adults were routinely discharged directly from the ED and routinely discharged after admission. Among patients receiving LPs, a larger proportion of adults died following admission as compared with children, but causes of death are unknown. Results of outcomes comparisons were similar when stratifying by age for pediatric patients. (S2 Appendix)

Discussion

Our results indicate that out of an estimated 135 million Emergency Department hospitalizations reported in the 2010 NEDS data, approximately 362,718 were associated with a diagnostic LP. These figures suggest that approximately 3 of every 1,000 ED treatments may be associated with a diagnostic LP in the ED or subsequent inpatient course. Furthermore, acknowledging the limitations of the data set, at least 37% of the total estimated LPs were performed in the ED. Though the majority of LPs were performed on adults, an estimated 64,849 (18% of total) were performed on children ≤5 years old. These observations characterize the frequency and burden of LPs in contemporary hospital practice.

There have been few prior descriptions of LP epidemiology among either ED or hospitalized patients. Shah, et al. examined the incidence of traumatic lumbar punctures in the ED, but only included patients from a single tertiary hospital ED.[14] Kroll, et al. evaluated US LP trends by medical specialty over span of 20 years but did not include detailed patient or institutional information from the ED.[15] Our study provides one of the first national descriptions of LP performance among patients receiving initial treatment in the ED.

Table 2. Patient and hospital characteristics of Emergency Department-treated patients by age and lumbar puncture status, 2010.

Characteristic	Patients <18 Years Old		P-value	Patients ≥18 Years Old		P-value
	Lumbar Puncture (N = 89,106)	No Lumbar Puncture (N = 25,421,964)		Lumbar Puncture (N = 273,612)	No Lumbar Puncture (N = 103,176,378)	
Patient Sex			<0.001			0.21
Male	55.0 (54.3–55.7)	52.7 (52.6–52.9)		42.2(41.6–42.9)	42.6 (42.3–42.9)	
Female	45.0 (44.3–45.7)	47.3 (47.1–47.4)		57.8(57.1–58.4)	57.4 (57.1–57.7)	
Patient Residence			<0.001			<0.001
Urban	88.6 (86.4–90.4)	79.7 (78.0–81.2)		87.1(85.5–88.6)	80.1 (78.8–81.3)	
Rural	11.4 (9.6–13.6)	20.3 (18.8–22.0)		12.9 (11.4–14.5)	19.9 (18.7–21.2)	
Patient Median Income			0.004			<0.001
\$1–40,999	28.5 (23.9–33.6)	32.9 (30.1–35.8)		26.6 (24.1–29.1)	33.2 (30.9–35.6)	
\$41,000–\$50,999	27.7 (25.3–30.3)	29.0 (26.9–31.2)		27.1 (25.3–29.0)	27.8 (26.1–29.4)	
\$51,000–\$66,999	24.8 (21.7–28.2)	22.0 (20.23–23.9)		25.0 (22.6–27.6)	21.7 (20.3–23.3)	
\$67,000 or more	19.0 (16.5–21.9)	16.0 (14.2–18.1)		21.3 (19.0–23.7)	17.3 (15.6–19.1)	
Hospital Census Region			<0.001			<0.001
Northeast	13.4 (9.6–18.4)	18.1 (15.8–20.7)		16.2 (13.4–19.6)	19.5 (17.9–21.1)	
Midwest	15.7 (11.1–21.8)	22.4 (19.9–25.2)		21.5 (18.7–24.6)	23.8 (21.9–25.8)	
South	40.1 (30.8–50.2)	39.6 (36.3–43.1)		37.3 (33.6–41.2)	39.4 (37.4–41.4)	
West	30.8 (22.2–41.1)	19.9 (17.3–22.8)		25.0 (21.5–28.7)	17.4 (16.0–18.8)	
Hospital Population Setting			<0.001			<0.001
Urban	93.2 (91.3–94.8)	80.7 (79.0–82.3)		91.6 (90.1–92.9)	81.8 (80.5–83.0)	
Rural	6.8 (5.3–8.7)	19.3 (17.8–21.1)		8.4 (7.2–9.9)	18.2 (17.0–19.5)	
Hospital Teaching Status			<0.001			<0.001
Metropolitan non-teaching	28.1 (22.4–34.6)	40.6 (37.6–43.7)		38.7(35.3–42.3)	42.4 (40.4–44.4)	
Metropolitan teaching	65.1 (57.7–71.9)	40.0 (36.3–43.9)		52.9 (49.0–56.7)	39.4 (37.2–41.6)	
Non-metropolitan teaching and non-teaching	6.8 (5.3–8.7)	19.3 (17.8–21.1)		8.4 (7.2–9.9)	18.2 (17.0–19.5)	
Hospital Trauma Center			<0.001			<0.001
Trauma center	59.1 (49.5–68.0)	38.4 (34.9–42.1)		48.5 (44.5–52.5)	35.7 (33.6–37.4)	
Not trauma center	41.0 (32.1–50.5)	61.6 (57.9–65.1)		51.5 (47.5–55.5)	64.4 (62.2–66.4)	
Admission by Season			<0.001			0.001
Spring	23.8 (23.0–24.7)	26.2 (26.1–26.3)		24.8 (24.3–25.3)	25.3 (25.2–25.5)	
Summer	27.9 (26.7–29.0)	23.9 (23.8–24.1)		26.8 (26.3–27.3)	26.3 (26.2–26.4)	
Fall	25.1 (24.3–26.0)	24.9 (24.7–25.1)		25.4 (24.8–26.0)	24.7 (24.6–24.8)	
Winter	23.2 (22.3–24.1)	24.9 (24.8–25.1)		23.0 (22.6–23.5)	23.7 (23.6–23.8)	

Table cells reflect column totals and 95% confidence intervals.

<https://doi.org/10.1371/journal.pone.0208622.t002>

An interesting observation was the disproportionately higher number of younger pediatric and adult LPs in the Southern US census region, which may be due to regional variations in disease presentation or practice. For example, the incidence of suspected meningitis may be higher in the South census region. Wang, et al. observed that sepsis mortality was highest in the Southeast US, potentially explaining the higher frequency of LPs.[16] We also observed a higher proportion of LPs among low income patients. We do not know whether these disparities are due to differences in disease prevalence or clinical practices. More targeted epidemiological studies of these specific factors could be warranted to evaluate whether this is a concern of public health or clinical practice.

Table 3. Primary diagnosis groupings among Emergency Department-treated patients undergoing lumbar puncture by age, 2010.

Diagnoses (CCS Codes)	Patients <18 Years (N = 89,106)	Patients ≥18 Years (N = 273,612)
Headache; including migraine 84	8.6 (7.7–9.7)	39.9 (38.2–41.6)
Meningitis (excluding tuberculosis or sexually transmitted diseases) 76	13.6 (12.4–14.8)	12.7 (12.2–13.3)
Fever of unknown origin 246	20.4 (19.1–21.8)	4.7 (4.3–5.0)
Epilepsy; convulsions 83	11.2 (10.3–12.1)	5.4 (5.1–5.8)
Septicemia 2	5.3 (4.7–5.9)	7.5 (7.0–8.1)
Other nervous system disorders 95	2.1 (1.8–2.5)	8.8 (8.2–9.5)
Other perinatal conditions 224	17.9 (15.8–20.3)	-
Viral Infection 7	7.0 (6.2–7.8)	3.4 (3.2–3.6)
Urinary tract infections 159	5.8 (5.2–6.5)	2.1 (2.0–2.3)
Acute cerebrovascular disease 109	0.04 (0.03–0.05)	3.8 (3.5–4.2)
Other upper respiratory infections 126	4.0 (3.6–4.5)	2.1 (1.9–2.3)
Pneumonia (excluding tuberculosis or sexually transmitted diseases) 122	2.5 (2.2–2.8)	2.4 (2.2–2.6)
Encephalitis (excluding tuberculosis or sexually transmitted diseases) 77	0.8 (0.7–1.0)	2.4 (2.2–2.6)
Spondylosis; intervertebral disc disorders; other back problems 205	0.6 (0.5–0.8)	1.9 (1.7–2.0)

Table cells reflect column totals and 95% confidence intervals.

CCS = Clinical Classification Software

<https://doi.org/10.1371/journal.pone.0208622.t003>

In highlighting the national number of procedures, our epidemiologic observations illuminate opportunities for improving the care of patients requiring LPs. LP is a complex, invasive procedure with potentially serious complications and challenges in the face of difficult patient anatomy such as obesity or distorted spinal anatomy. Novel LP techniques include the use of adjunctive fluoroscopy, CT, and ultrasonography to facilitate LP performance. Our observations suggest that a considerable number of patients may benefit from these approaches. Novel biomarkers may identify the early stages of cerebral infection and neurodegenerative diseases. [17–19] For example, some complement activation components and the soluble membrane attack complex show promise as biomarkers of bacterial meningitis and shunt infection in patients with hydrocephalus. [20, 21] Given that LPs are most commonly used to rule out bacterial meningitis, the potential for the application of these diagnostic biomarkers is large.

Table 4. Disposition of Emergency Department-treated patients by age and lumbar puncture status, 2010.

Emergency Department Disposition	Patients <18 Years Old		Patients ≥18 Years Old	
	Lumbar Puncture (N = 89,106)	No Lumbar Puncture (N = 25,421,964)	Lumbar Puncture (N = 273,612)	No Lumbar Puncture (N = 103,176,378)
Routine discharge	25.6 (23.0–28.3)	93.6 (93.1–94.0)	36.0 (33.8–38.2)	77.3 (76.7–77.9)
Transfer to short term hospital	8.9 (6.9–11.3)	1.2 (1.1–1.3)	1.5 (1.3–1.8)	1.6 (1.4–1.7)
Died in ED	N/A	0.03 (0.03–0.04)	N/A	0.2 (0.2–0.2)
Other	N/A	1.4 (1.1–1.8)	N/A	3.2 (2.9–3.5)
Admit				
Routine discharge	59.2 (54.7–63.5)	3.5 (3.1–3.9)	36.9 (35.6–38.2)	11.2 (10.8–11.5)
Transfer to short term hospital	1.9 (1.5–2.3)	0.08 (0.08–0.09)	2.5 (2.3–2.8)	0.5 (0.5–0.5)
Died on ward	0.2 (0.2–0.3)	0.01 (0.01–0.02)	2.1 (1.9–2.3)	0.5 (0.5–0.5)
Other	3.3 (2.4–4.4)	0.2 (0.15–0.26)	22.2 (20.9–23.6)	6.3 (6.1–6.5)

Table cells denote column percentages and 95% confidence intervals. N/A = not available due to n<11.

<https://doi.org/10.1371/journal.pone.0208622.t004>

Limitations

This study has several limitations. NEDS aggregates patients treated only in the ED as well as those admitted through the ED. Diagnostic LPs performed on an outpatient basis were not included in this study. Among admitted patients, we could not differentiate LPs occurring in the ED vs. those performed later in the hospital course. Thus, our observed number of ED LPs may be an underestimate. As a result of this, we also could not assess the specialty of the provider who performed the LP nor whether the admission was a result of the LP or not. We could not ascertain the use or success of different LPs approaches such as patient positioning or the use of varying LP needles.

We were limited to 2010 data and could not evaluate secular trends. Although the NEDS contain comprehensive data, variations in documentation practices may have led to underestimates of the number of LPs. We did not have data on adverse events (such as post-dural headaches) and, therefore, also could not distinguish whether the discharge diagnosis, such as a headache, was the indication for the LP, or a result of the LP. Finally, although we had CCS for each case, these codes do not indicate the reason for an LP. Diagnoses such as pneumonias or UTIs may have denoted instances of sepsis workups where a diagnostic LP was performed as part of the work-up. Alternatively, the LP may have been done later in response to additional developing symptoms.

Conclusions

Approximately 363,000 LPs are performed each year among ED-treated patients in the United States. These observations highlight the national number of diagnostic LP interventions and the opportunities for improving the care of these patients with potential neurologic emergencies.

Supporting information

S1 Appendix. Patient and hospital characteristics of Emergency Department-treated patients undergoing lumbar puncture, stratified by age.

(DOCX)

S2 Appendix. Disposition of Emergency Department-treated patients by age and lumbar puncture status with additional stratification, 2010.

(DOCX)

Acknowledgments

We acknowledge Robert Liu, MD in providing editorial assistance in the preparation and revision of this paper.

Author Contributions

Conceptualization: Adrienne Vickers, John P. Donnelly, Henry E. Wang.

Formal analysis: John P. Donnelly, Henry E. Wang.

Investigation: Adrienne Vickers, Justin Xavier Moore, Theresa N. Schein.

Methodology: Adrienne Vickers, John P. Donnelly, Henry E. Wang.

Resources: Scott R. Barnum, Theresa N. Schein.

Software: John P. Donnelly.

Supervision: Henry E. Wang.

Writing – original draft: Adrienne Vickers.

Writing – review & editing: John P. Donnelly, Justin Xavier Moore, Scott R. Barnum, Theresa N. Schein, Henry E. Wang.

References

1. Editorial: Lumbar puncture. *British medical journal*. 1975; 1(5948):3. PMID: [1120224](#); PubMed Central PMCID: PMC1671923.
2. Kerrigan S, Fuller G. Lumbar puncture—a practical guide. *The Foundation Years*. 2008; 4(4):167–9.
3. Swartz MN. Bacterial meningitis—a view of the past 90 years. *N Engl J Med*. 2004; 351(18):1826–8. <https://doi.org/10.1056/NEJMp048246> PMID: [15509815](#).
4. Domingo P, Pomar V, de Benito N, Coll P. The spectrum of acute bacterial meningitis in elderly patients. *BMC Infect Dis*. 2013; 13:108. Epub 2013/02/27. <https://doi.org/10.1186/1471-2334-13-108> PMID: [23446215](#); PubMed Central PMCID: PMCPMC3599144.
5. Armon C, Evans RW, Neurology T, of TASotAA. Addendum to assessment: Prevention of post-lumbar puncture headaches: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 2005; 65(4):510–2. <https://doi.org/10.1212/01.wnl.0000173034.96211.1b> PMID: [16116106](#).
6. Joffe AR. Lumbar puncture and brain herniation in acute bacterial meningitis: a review. *J Intensive Care Med*. 2007; 22(4):194–207. <https://doi.org/10.1177/0885066607299516> PMID: [17712055](#).
7. Shetty AK, Desselle BC, Craver RD, Steele RW. Fatal cerebral herniation after lumbar puncture in a patient with a normal computed tomography scan. *Pediatrics*. 1999; 103(6 Pt 1):1284–7. PMID: [10353943](#).
8. Büller HR, Prins MH, Lensin AW, Decousus H, Jacobson BF, Minar E, et al. Oral rivaroxaban for the treatment of symptomatic pulmonary embolism. *N Engl J Med*. 2012; 366(14):1287–97. Epub 2012/03/26. <https://doi.org/10.1056/NEJMoa1113572> PMID: [22449293](#).
9. Ntaios G, Papavasileiou V, Diener HC, Makaritsis K, Michel P. Nonvitamin-K-antagonist oral anticoagulants versus warfarin in patients with atrial fibrillation and previous stroke or transient ischemic attack: An updated systematic review and meta-analysis of randomized controlled trials. *Int J Stroke*. 2017; 12(6):589–96. Epub 2017/03/15. <https://doi.org/10.1177/1747493017700663> PMID: [28730948](#).
10. Lip GY, Lane DA. Stroke prevention with oral anticoagulation therapy in patients with atrial fibrillation—focus on the elderly. *Circ J*. 2013; 77(6):1380–8. Epub 2013/05/09. PMID: [23657131](#).
11. Edwards C, Leira EC, Gonzalez-Alegre P. Residency training: a failed lumbar puncture is more about obesity than lack of ability. *Neurology*. 2015; 84(10):e69–72. <https://doi.org/10.1212/WNL.000000000001335> PMID: [25754807](#).
12. (HCUP) HCUP. HCUP Overview [Web page]. Rockville, MD [updated April 21, 2017February 6, 2016]. Available from: <https://www.hcup-us.ahrq.gov/overview.jsp>.
13. (HCUP) HCUP. NEDS Overview Rockville, MD [updated March 22, 2017February 9, 2016]. Available from: <https://www.hcup-us.ahrq.gov/nedsoverview.jsp>.
14. Shah KH, Richard KM, Nicholas S, Edlow JA. Incidence of traumatic lumbar puncture. *Academic emergency medicine: official journal of the Society for Academic Emergency Medicine*. 2003; 10(2):151–4. Epub 2003/02/08. PMID: [12574013](#).
15. Kroll H, Duszak R Jr., Nsiah E, Hughes DR, Sumer S, Wintermark M. Trends in lumbar puncture over 2 decades: a dramatic shift to radiology. *AJR Am J Roentgenol*. 2015; 204(1):15–9. Epub 2014/12/30. <https://doi.org/10.2214/AJR.14.12622> PMID: [25539231](#).
16. Wang HE, Devereaux RS, Yealy DM, Safford MM, Howard G. National variation in United States sepsis mortality: a descriptive study. *Int J Health Geogr*. 2010; 9:9. Epub 2010/02/17. <https://doi.org/10.1186/1476-072X-9-9> PMID: [20156361](#); PubMed Central PMCID: PMCPMC2831852.
17. Fitzner B, Hecker M, Zettl UK. Molecular biomarkers in cerebrospinal fluid of multiple sclerosis patients. *Autoimmun Rev*. 2015; 14(10):903–13. Epub 2015/06/10. <https://doi.org/10.1016/j.autrev.2015.06.001> PMID: [26071103](#).
18. Berger T, Reindl M. Antibody biomarkers in CNS demyelinating diseases—a long and winding road. *Eur J Neurol*. 2015; 22(8):1162–8. Epub 2015/05/24. <https://doi.org/10.1111/ene.12759> PMID: [26010364](#).
19. Reiber H, Peter JB. Cerebrospinal fluid analysis: disease-related data patterns and evaluation programs. *J Neurol Sci*. 2001; 184(2):101–22. PMID: [11239944](#).

20. Stahel PF, Nadal D, Pfister HW, Paradisis PM, Barnum SR. Complement C3 and factor B cerebrospinal fluid concentrations in bacterial and aseptic meningitis. *Lancet*. 1997; 349(9069):1886–7. Epub 1997/06/28. [https://doi.org/10.1016/S0140-6736\(05\)63877-9](https://doi.org/10.1016/S0140-6736(05)63877-9) PMID: 9217766.
21. Ramos TN, Arynchyna AA, Blackburn TE, Barnum SR, Johnston JM. Soluble membrane attack complex is diagnostic for intraventricular shunt infection in children. *JCI Insight*. 2016; 1(10):e87919. Epub 2016/10/05. <https://doi.org/10.1172/jci.insight.87919> PMID: 27699221; PubMed Central PMCID: PMC5033849.