

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2019

Improvement in patient-reported sleep in type 2 diabetes and prediabetes participants receiving a continuous care intervention with nutritional ketosis

Morgan J. Siegmann
Harvard University

Shaminie Athinarayanan
Virta Health

Sarah J. Hallberg
Indiana University Health Arnett

Amy L. McKenzie
Virta Health

Nasir H. Bhanpuri
Virta Health

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Siegmann, Morgan J.; Athinarayanan, Shaminie; Hallberg, Sarah J.; McKenzie, Amy L.; Bhanpuri, Nasir H.; Campbell, Wayne W.; McCarter, James P.; Phinney, Stephen D.; Volek, Jeff S.; and Van Dort, Christa J., "Improvement in patient-reported sleep in type 2 diabetes and prediabetes participants receiving a continuous care intervention with nutritional ketosis." *Sleep Medicine*. 55, 92-99. (2019).
https://digitalcommons.wustl.edu/open_access_pubs/7739

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Morgan J. Siegmann, Shaminie Athinarayanan, Sarah J. Hallberg, Amy L. McKenzie, Nasir H. Bhanpuri, Wayne W. Campbell, James P. McCarter, Stephen D. Phinney, Jeff S. Volek, and Christa J. Van Dort



Original Article

Improvement in patient-reported sleep in type 2 diabetes and prediabetes participants receiving a continuous care intervention with nutritional ketosis



Morgan J. Siegmann^{a, b, c}, Shaminie J. Athinarayanan^d, Sarah J. Hallberg^{d, e}, Amy L. McKenzie^d, Nasir H. Bhanpuri^d, Wayne W. Campbell^f, James P. McCarter^{d, g}, Stephen D. Phinney^d, Jeff S. Volek^{d, h}, Christa J. Van Dort^{a, b, c, *}

^a Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, MA, USA

^b Division of Sleep Medicine, Harvard Medical School, Boston, MA, USA

^c Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA

^d Virta Health, 535 Mission Street, San Francisco, CA, USA

^e Indiana University Health Arnett, Lafayette, IN, USA

^f Department of Nutrition Science, Purdue University, West Lafayette, IN, USA

^g Department of Genetics, Washington University School of Medicine, St Louis, MO, USA

^h Department of Human Sciences, The Ohio State University, Columbus, OH, USA

ARTICLE INFO

Article history:

Received 10 August 2018

Received in revised form

11 December 2018

Accepted 17 December 2018

Available online 3 January 2019

Keywords:

Type 2 diabetes

Prediabetes

Ketogenic diet

PSQI

Nutritional ketosis

ABSTRACT

Objective: Sleep disruption is frequently associated with type 2 diabetes (T2D) and hyperglycemia. We recently reported the effectiveness of a continuous care intervention (CCI) emphasizing nutritional ketosis for improving HbA1c, body weight and cardiovascular risk factors in T2D patients. The present study assessed the effect of this CCI approach on sleep quality using a subjective patient-reported sleep questionnaire.

Methods: A non-randomized, controlled longitudinal study; 262 T2D and 116 prediabetes patients enrolled in the CCI and 87 separately recruited T2D patients continued usual care (UC) treatment. Patients completed the Pittsburgh Sleep Quality Index (PSQI) questionnaire. A PSQI score of >5 (scale 0 to 21) was used to identify poor sleepers.

Results: Global sleep quality improved in the CCI T2D ($p < 0.001$) and prediabetes ($p < 0.001$) patients after one year of intervention. Subjective sleep quality (component 1), sleep disturbance (component 5) and daytime dysfunction (component 7), also showed improvements in the CCI T2D ($p < 0.01$ for sleep quality and sleep disturbance; and $p < 0.001$ for daytime dysfunction) and prediabetes patients ($p < 0.001$ for all three components); compared to the UC T2D group after one year. The proportion of patients with poor sleep quality was significantly reduced after one year of CCI (T2D; from 68.3% at baseline to 56.5% at one year, $p = 0.001$ and prediabetes; from 77.9% at baseline to 48.7% at one year, $p < 0.001$).

Conclusion: This study demonstrates improved sleep quality as assessed by PSQI in patients with T2D and prediabetes undergoing CCI including nutritional ketosis but not in T2D patients receiving UC. The dietary intervention benefited both sleep quality and the severity of T2D symptoms suggesting that nutritional ketosis improves overall health via multiple mechanisms.

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abbreviations: CCI, continuous care intervention; UC, usual care; T2D, type 2 diabetes; BMI, body mass index; PSQI, Pittsburgh Sleep Quality Index; OSA, obstructive sleep apnea; HbA1c, hemoglobin A1c; CPAP, continuous positive airway pressure; AHI, apnea and hypopnea indices; KD, ketogenic diet; REM, rapid eye movement; SWS, slow wave sleep; BHB, beta-hydroxybutyrate; HOMA-IR, homeostatic model assessment of insulin resistance; hsCRP, high-sensitivity C-reactive protein.

* Corresponding author. Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Division of Sleep Medicine, Harvard Medical School, MGH Building 149 Room 4140, 149 13th Street, Charlestown, MA 02129, USA.

E-mail address: cvandort@mgh.harvard.edu (C.J. Van Dort).

<https://doi.org/10.1016/j.sleep.2018.12.014>

1389-9457/© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Sleep disruption is associated with obesity and type 2 diabetes (T2D), yet the bidirectional relationship between sleep and glucose metabolism is not fully understood. It is linked to increased diabetes prevalence in both experimental [1–4] and epidemiological studies [5–7]. In addition, the severity of hyperglycemia in individuals with diabetes is associated with poor sleep quality [8–11], short sleep duration [8,9,12,13] and a greater tendency to develop sleep disorders including obstructive sleep apnea (OSA) [14,15]. Both the International Diabetes Federation (IDF) and American Diabetes Association (ADA) recommend evaluating T2D patients for sleep breathing problems especially OSA and strongly encourage treatment when found [16,17].

Weight loss has demonstrated effectiveness to improve sleep quality, quantity [18,19] and to treat OSA in obese patients. Lifestyle intervention induced weight loss showed significant reduction in the apnea and hypopnea indices (AHI) in conjunction with a decrease in hemoglobin A1c (HbA1c) levels in a randomized controlled trial of obese OSA patients with comorbid diabetes [20]. Further, weight loss following bariatric surgery is effective at improving glycemic control and improving AHI in OSA patients [21]. Intervention studies specifically targeting sleep disruption in OSA patients without any effect on weight, such as continuous positive airway pressure (CPAP) treatment, have shown contradictory results for glycemic control. Most CPAP intervention studies in T2D reported no glycemic benefit from the treatment [22,23], but one study demonstrated a slight reduction in HbA1c [24]. In contrast, CPAP studies on prediabetic OSA patients showed improvements in insulin sensitivity and glucose tolerance [25,26]. It is not clear from these studies whether improvement of glycemic control in conjunction with weight loss improves sleep quality or vice-versa.

A few studies have investigated the impact of dietary macronutrient composition on sleep duration and quality. Two studies reported reduction of slow wave sleep (SWS) and elevation of rapid eye movement (REM) sleep in individuals consuming higher carbohydrates (600 g carbohydrate or 80% energy from carbohydrate) [27,28]. Another study reported the effect of a high carbohydrate (56% energy from carbohydrate) diet in reducing sleep onset latency when compared to a control diet [29]. Studies investigating low carbohydrate diets showed the opposite effect; reduced REM [30], increased REM onset latency [31] and increased SWS [30], even after 4 h of administering a very low carbohydrate meal [30]. Collectively, these findings signify dietary carbohydrate content as an important factor in modulating sleep architecture, but extrapolation from these studies is limited since they were conducted in experimentally controlled conditions with small numbers of healthy individuals in a short time-span and with diets administered at specific time points.

Population and intervention-based studies on the overall impact of carbohydrate intake on sleep indices or sleep quality are very limited. Katagiri et al., showed reduced sleep quality in individuals consuming more carbohydrates as measured by a subjective sleep measure, the Pittsburgh Sleep Quality Index (PSQI) [32]. Studies investigating the effect of ketogenic diet (KD) in children with sleep problems showed improvement in daytime sleepiness [33,34] as well as positive changes in sleep architecture [34,35]. However, in one of these studies, sleep improvements were suggested to be due to weight loss rather than the KD [35]. Despite restricted carbohydrate intake concurrent with sleep improvement in these children, SWS decreased [35] and REM increased [34,35] which contradicts studies on carbohydrate intake and sleep architecture in adults [27,28,30]. Carbohydrate restriction and ketogenic diets are widely used in the clinical management of obesity and diabetes,

but studies assessing the effect of this diet on sleep are currently limited. We recently demonstrated a continuous remote care treatment for T2D including nutritional ketosis significantly improved glycemic control, weight, and cardiovascular disease risk factors and reduced diabetes medication use at one year [36–38].

The purpose of this study was to assess the effect of the intervention by time-interval on the global PSQI and its seven component scores as well as compared its changes with different intervention and disease categories. We also assessed the relationship between changes in the sleep parameters versus key biochemical parameters, and also investigated the correlation of pain, circadian rhythm disruption and CPAP usage versus patient-perceived sleep status. We hypothesized that the global sleep indexes would improve analogously, as improvement in other key biochemical parameters observed in the intervention.

2. Materials and methods

2.1. Study participants and design

This study is part of a clinical trial (*Clinical trials.gov identifier*: NCT02519309) that was approved by the Franciscan Health Lafayette Institutional Review Board. Patients between age 21 and 65 years with either a diagnosis of T2D and a BMI >25 kg/m² or prediabetes and a BMI >30 kg/m² were included in this study. Detailed study design including the inclusion and exclusion criteria were previously reported [36,37]. Briefly, the trial was an open-label, non-randomized, controlled, longitudinal study with patients divided into three groups. The T2D and pre-diabetes patients in the continuous care intervention (CCI) regimen self-selected either on-site (CCI-onsite) or web-based (CCI-web) education delivery. Educational content and medical treatment was the same for both CCI-onsite and CCI-web. As there were no significant differences in outcomes including PSQI scores, between educational groups, they are combined for further analysis [36,37]. Both T2D and prediabetes CCI patients had access to a mobile health application (app) that enabled them to communicate and be continuously monitored by a team of healthcare professionals including a personal health coach and physician or nurse practitioner. Patients received individualized guidance in achieving nutritional ketosis, typically including restriction of daily dietary carbohydrates to less than 30 g. Patients were encouraged to measure and input weight, blood glucose and blood beta-hydroxybutyrate (BHB) concentrations daily in the app. These measurements were used by the health care team for monitoring the patient's condition (weight and glucose) and assessing carbohydrate restriction (BHB).

Separately recruited usual care (UC) T2D patients were participants in a local diabetes education program including care by their primary care physician or endocrinologist and counseling by registered dietitians; no modification to their care was made for the study. This group was observed at baseline and one year as reference for typical disease treatment and progression within the same geography and health system. UC patients were informed that the trial had an intervention arm and could participate in that group if they chose to do so.

2.2. Demographic and clinical variables

Patient demographic and clinical data were collected at baseline, 70 days and one year. Laboratory measures were assessed at a Clinical Laboratory Improvement (CLIA) certified laboratory. These data were initially analyzed to evaluate the safety and effectiveness of the CCI in improving diabetes status (glycemic control and medication use), weight and other metabolic factors in T2D [36,37] and prediabetes patients [38] (unpublished data, manuscript in

preparation). Some of the clinical variables – weight, fasting blood glucose, HbA1c, homeostatic model assessment of insulin resistance (HOMA-IR), BHB and high sensitivity C-reactive protein (hsCRP) – were included for further analyses in this study. Usual care T2D patients were not continuously monitored for weight, blood glucose, or BHB; clinical and laboratory measures were obtained for this group only at baseline and one year.

2.3. Pittsburgh Sleep Quality Index (PSQI)

CCI patients were administered a set of questionnaires, including the PSQI, during visits at baseline, 70 days and one year; UC participants completed questionnaires at baseline and one year. The PSQI consists of 19 validated questions assessing sleep quality and efficiency [39]. The global PSQI score is calculated from seven component scores on subjective sleep quality (component 1), sleep latency (component 2), sleep duration (component 3), habitual sleep efficiency (component 4), sleep disturbances (component 5), use of sleep medication (component 6) and daytime dysfunction (component 7). Each question within the component is scored on a 4-point Likert scale of 0–3, with 3 indicating worse outcomes and the mean was calculated for each component score. The sum of the component score means generates the global PSQI score that ranges from 0 to 21. Higher global PSQI scores indicate poorer sleep. A patient with a global PSQI score ≤ 5 is considered a “good sleeper” and >5 is categorized as a “poor sleeper” [40]. Change in the PSQI score over time was calculated using the formula below:

$$\text{Delta PSQI} = \frac{(\text{Post} - \text{intervention PSQI} - \text{Baseline PSQI})}{\text{Baseline PSQI}}$$

2.4. Pain, shifted sleep chronotype and CPAP usage

Patients were classified into “pain” and “non-pain” groups based on their response to pain-related questions in both the PSQI (question 5i) and a separate questionnaire used to calculate the knee injury and osteoarthritis outcome score (KOOS). Overall KOOS results will be reported in a separate publication. Classification of patients under circadian rhythm “disrupted” and “non-disrupted” groups was based on the wake time and bedtime responses for PSQI questions 1 and 3 for compilation of component 4 (sleep efficiency). Patients were classified as having a shifted wake-up time if they reported typically waking between 11am and 2am, while those with bedtimes between 12am and 6pm were bedtime shifted. These arbitrary bedtime and wake time cut-off ranges were selected based on evening and night shift workers schedule (second shift – 3pm to 11pm and third shift- 11pm to 7am); which causes these workers to have sleep patterns that deviate from a normal chronotype. Patients were also surveyed regarding CPAP usage and discontinuation, however detailed usage information such as CPAP pressure settings and usage compliance were not obtained making it difficult to interpret the patients OSA treatment status.

2.5. Statistical analyses

The questionnaires were administered by research personnel and completed by patients on paper. Paper questionnaires were scanned and responses were transcribed in duplicate by an independent contract data entry firm. The patterns of missing data were assessed using Little's MCAR test [41] and were found to be missing at random (MAR). Missing data were imputed by Multivariate Imputation by Chained Equations (MICE) [42], and Intent to treat (ITT) analyses were performed. Normality of the global PSQI and component scores

was evaluated using Lilliefors test. Even after transformation, the data failed the normality test (ie there was a skew toward lower PSQI scores and a long tail of higher scores) (Supplemental Fig. 1A–C); therefore, nonparametric tests were used for analyses of PSQI scores. Results from continuous variables were expressed as mean \pm standard deviation. Comparisons between groups were performed using the Kruskal–Wallis test, and comparisons within groups were performed using the Wilcoxon Sign Rank test. Tukey's honest significant difference test was used to analyze pairwise differences among significant results from omnibus tests. McNemar's test was used for assessing statistical significance of transitioning between ‘good’ and ‘poor’ sleeper among the CCI and UC cohorts.

Adjusted Pearson's and Spearman correlations were calculated between changes from baseline in global PSQI and changes in metabolic-parameters. Adjusted correlations were performed while controlling for age, gender and BMI at baseline. All participants in the CCI group were stratified by sleep improvement status based on their baseline and one year global PSQI scores. Patients that were initially considered “poor sleepers” with a baseline PSQI >5 but whose score after one year decreased to at or below the threshold of 5 were classified as improved. Those patients who were considered “good sleepers” at both baseline and one year were classified as maintained. Finally, those patients whose 1 year PSQI score was >5 (regardless of their baseline score) were classified as not improved. Stepwise analyses of covariance (ANCOVA) were performed between the three different CCI sleep status groups at one year with the change of the glucose-related, ketone and inflammatory markers, while controlling by age, gender and years living with diabetes. Statistical tests were performed with MATLAB R2017b using the Statistics and Machine Learning Toolbox [43] and the R statistical program version 3.5.0 [44].

3. Results

3.1. Baseline participant characteristics

Details on the recruitment and extensive baseline characteristics of the CCI and UC T2D patients were previously published [36,37]. The demographic, glycemic, inflammatory and sleep baseline characteristics of the participants that were included for assessments of sleep are presented in Table 1. One-hundred forty-three (54.6%) CCI T2D, 61 (54%) CCI prediabetes, and 53 (62.3%) UC T2D patients completed the PSQI at all expected time points. The number of patients who completed the trial at one year were slightly higher than those who completed the PSQI questionnaires. Some of the patients completed the study period and laboratory analysis but were unable to attend the clinic for their 70-days and one-year follow-up visits, where they are required to complete their corresponding questionnaires. The proportion of missing PSQI data were similar across the three groups with 77.61% of CCI T2D, 79.06% of CCI prediabetes and 79.24% of UC T2D completed the PSQI in all expected time points. There were no significant differences between completers and non-completers on baseline characteristics for either group at one year of the intervention (Supplemental Table 1). The global PSQI and component scores did not differ significantly among the groups (CCI T2D, CCI prediabetes and UC T2D) at baseline. The proportion of participants with overall poor sleep quality was higher in the CCI prediabetes group (77.9%) compared to the CCI T2D (68.3%) and UC T2D (68.2%) groups.

3.2. Effect of intervention on sleep

3.2.1. Global PSQI and component scores

Overall sleep quality as assessed by the global PSQI score, improved in CCI T2D (median change from 7 to 6; $p < 0.001$) and

Table 1

Baseline characteristics of participants included in the study. Baseline data were calculated using intent-to-treat (ITT) data.

Patient Cohorts	CCI Type 2 Diabetes	CCI Prediabetes	UC Type 2 Diabetes
Starters, Completers, PSQI Available (n)	262, 218, 143	116, 113, 61	87, 78, 53
	mean (S.D.)	mean (S.D.)	mean (S.D.)
Age (years)	53.8 (±8.4)	51.9 (±9.4)	52.7 (±9.3)
Male/female (ratio)	87/175 (1:2)	29/84 (1:3)	35/50 (2:3)
Body weight (kg)	116.4 (±26.1)	109.9 (±23.6)	108.3 (±25.1)
BMI (kg/m ²)	40.4 (±8.9)	38.8 (±7.1)	38.2 (±9.1)
Fasting glucose (mg/dL)	160.78 (±61.32)	109.58 (±15.20)*	157.08 (±72.48)
HbA1c (%)	7.60 (±1.50)	5.91 (±0.24)*	7.67 (±1.77)
HOMA-IR	11.8 (±13.1)	7.1 (±7.4)*	13.7 (±17.8)
high sensitivity C-reactive protein (nmol/L)	9.31 (±19.31)	7.46 (±7.51)	9.34 (±9.10)
Beta-hydroxybutyrate (mmol/L)	0.17 (±0.15)	0.14 (±0.13)	0.15 (±0.12)
Global PSQI Score	7.72 (±3.72)	7.96 (±3.43)	7.92 (±3.85)
Subjective sleep quality	1.18 (±0.75)	1.22 (±0.73)	1.25 (±0.79)
Sleep latency	1.09 (±0.93)	1.33 (±0.958)	1.05 (±0.89)
Sleep duration	1.23 (±0.92)	1.27 (±0.96)	1.14 (±0.94)
Habitual sleep efficiency	0.68 (±0.99)	0.61 (±0.89)	0.71 (±1.04)
Sleep disturbances	1.64 (±0.63)	1.66 (±0.68)	1.75 (±0.74)
Use of sleep medication	0.69 (±1.16)	0.66 (±1.11)	0.85 (±1.26)
Daytime dysfunction	1.22 (±0.77)	1.21 (±0.76)	1.17 (±0.86)
Poor sleepers N (%)	179 (68.3)	88 (77.9)	58 (68.2)
Good sleepers N (%)	83 (31.7)	25 (22.1)	27 (31.8)

Note. Subjective sleep quality, component 1; sleep latency, component 2; sleep duration, component 3; habitual sleep efficiency, component 4; sleep disturbances, component 5; use of sleep medication, component 6, and daytime dysfunction, component 7.

*p-value <0.001.

prediabetes (median change from 7 to 5; $p < 0.001$) groups after one year of the intervention (Fig. 1). No significant change in the global PSQI score was observed in UC T2D (median change from 7 to 8, $p = 0.245$). At one year, global PSQI scores in the CCI T2D ($p < 0.001$) and prediabetes ($p < 0.01$) were significantly lower than in the UC T2D, whereas no differences were observed at baseline (Supplementary Fig. 2A). Among patients characterized as poor sleepers at baseline (global PSQI >5), one year global PSQI score was lower in the CCI T2D ($p < 0.001$) and prediabetes ($p < 0.001$) than in the UC T2D (Supplementary Fig. 2B). Greater reduction in the global PSQI score was observed in CCI T2D (median change of -1 , $p < 0.01$)

and CCI prediabetes groups (median change of -2 , $p < 0.001$) compared to the UC T2D group (Supplementary Fig. 3). Further assessment of the PSQI component scores revealed three of the seven components showed significant change at one year for CCI T2D and prediabetes groups. Subjective sleep quality ($p < 0.01$ CCI T2D; $p < 0.001$ CCI prediabetes), sleep disturbance ($p < 0.01$ CCI T2D; $p < 0.001$ CCI prediabetes) and daytime dysfunction ($p < 0.001$ CCI T2D; $p < 0.001$ CCI prediabetes) scores were lower in the CCI T2D and prediabetes patients compared to the UC T2D group at one year (Fig. 2 A–C).

3.2.2. Resolution of poor sleep quality

There were 179 (68.3%) T2D and 88 (77.9%) prediabetes patients categorized as “poor sleepers” in the CCI at baseline. The proportions of “poor sleepers” in the CCI were reduced after one year of the intervention, with 56.5% of T2D ($p = 0.001$) and 48.7% ($p < 0.001$) of prediabetes patients categorized as “poor sleepers” at one year. In the UC cohort, the proportion of patients categorized as “poor sleepers” did not change after one year (68.2% at baseline to 69.4% at one year).

3.2.3. Association within the CCI group between changes in global PSQI with metabolic and inflammatory markers

Table 2 shows correlations between changes in the global PSQI score with changes in glucose-related, ketone and inflammatory markers in the CCI. In the prediabetes group, changes in fasting glucose ($r = 0.23$, $p = 0.02$) and HOMA-IR ($r = 0.32$, $p < 0.001$) were correlated to changes in PSQI scores after controlling for baseline age, sex and weight. Increased ketone concentrations in the prediabetes participants were also associated with reduction of global PSQI scores ($r = -0.242$, $p = 0.01$). These correlations observed in the prediabetes group were not present in the CCI T2D group and changes in the HbA1c and hsCRP did not correlate with changes in global PSQI scores in either group. Change in mean weight ($p = 0.04$) and HOMA-IR ($p = 0.01$) were the only variables independently and significantly associated between the three different sleep status (improved, maintained and not improved sleep status) at one year of the intervention. No statistically significant differences were found in weight loss changes between patients with




Fig. 1. Distribution of global PSQI scores at baseline and 365 days in CCI T2D, CCI PreD and UC T2D. Global PSQI score was significantly reduced in the CCI T2D and CCI PreD groups but not in the UC T2D group after 365 days. Boxplot descriptors (Figs. 1 and 2; Supplementary Figs. 3 and 4) Horizontal line within the box indicates median; upper and lower boundaries of the box represent the 25th and 75th percentiles; whiskers of the box indicate the range and “+++” signs represent outlier values. ** p-value <0.01; *** p-value <0.001.




Fig. 2. Distribution of PSQI components subjective sleep quality, sleep disturbances and daytime dysfunction in CCI T2D, CCI PreD and UC T2D groups at three different timepoints (0, 70 and 365 days). Subjective sleep quality (A), sleep disturbances (B), and daytime dysfunction (C) were significantly lower in the CCI T2D and CCI PreD groups when compared to UC T2D group at 365 days. Boxplot descriptors (Figs. 1 and 2; Supplementary Figs. 3 and 4) Horizontal line within the box indicates median; upper and lower boundaries of the box represent the 25th and 75th percentiles; whiskers of the box indicate the range and “+++” signs represent outlier values. ** p-value <0.01; *** p-value <0.001.

Table 2

Correlation analyses between change in the global PSQI score and change in metabolic parameters after one year of CCI.

Variable	CCI T2D Cohort N = 262				CCI Prediabetes Cohort N = 113			
	rho	P value*	Adjusted r	P value ⁺	rho	P value*	Adjusted r	P value ⁺
Δ Fasting glucose (mg/dl)	0.032	0.60	0.008	0.90	0.240	0.01	0.226	0.018
Δ HbA1c (%)	-0.037	0.55	-0.049	0.44	-0.024	0.80	-0.032	0.74
Δ HOMA-IR	-0.060	0.34	-0.069	0.27	0.314	0.0008	0.323	0.0006
Δ BHB	-0.003	0.96	-0.044	0.49	-0.297	0.002	-0.242	0.011
Δ hsCRP	-0.067	0.29	-0.008	0.90	-0.022	0.82	-0.032	0.74

*Spearman and ⁺Adjusted Pearson's correlations. Adjustments while controlling for age, sex and baseline weight.

improved, maintained and not improved sleep status. Patients who maintained sleep showed highest reductions of HOMA-IR (-6.94 ± 0.86), with statistically significant difference than those who did not improve sleep, after one year of the intervention ($p = 0.02$). Improvements in HOMA-IR among patients in the improved sleep (-4.17 ± 0.86) and not improved sleep status (-4.24 ± 0.55) did not differ significantly.

3.3. Effect of persistent pain on sleep improvement

We further assessed the effect of pain on sleep improvement in the CCI by classifying the patient's pain status using response retrieved from questions specifically related to pain in the sleep and knee (KOOS) questionnaires. As illustrated in Supplementary Fig. 4, patients with pain had higher global PSQI scores, indicating poorer sleep, compared to those categorized under “non-pain” group at all three time points. Both patients in the “non-pain” (Supplementary Fig. 5A, $p < 0.001$), and “pain” group (Supplementary Fig. 5B, $p < 0.01$) had reductions in their global PSQI score at 70 days and one year.

3.4. Effect of shifted sleep chronotype on sleep improvement

We also assessed the effect of shifted sleep chronotype on the global PSQI score improvement. Patients were classified as having shifted sleep chronotype based on their self-reported wake-up times and bedtimes as defined in the methods. There were 18, 27, and 96 patients in the CCI cohort classified as both wake-up time and bedtime shifted, wake-up time shifted only or bedtime shifted only respectively. Patients with shifted bedtimes, had reduced global PSQI scores ($p < 0.01$), as did those with normal chronotype ($p < 0.001$) (Supplementary Fig. 6A and B). However, those patients with shifted wake-up times (Supplementary figures 6C) did not show a change in their global PSQI score after one year of the intervention. Those with both shifted wake-up times and bedtimes

also did not show a change in their global PSQI score after one year of the intervention.

3.5. Effect of CPAP usage on sleep improvement

At baseline, there were a total of 140 participants in both CCI and UC treatment groups with CPAP equipment prescribed for sleep. Among CPAP users, 91 were in the CCI T2D group, 31 in the CCI prediabetes and 18 in the UC T2D group. Fifteen (13 CCI T2D and two UC T2D) of the 140 participants discontinued using CPAP at one year. Only six (46%) of the 13 CCI T2D participants discontinued due to patient-reported improvement in sleep quality from the CCI and reduction of weight; the remaining seven reported discontinuation due to discomfort or personal choice. Global PSQI scores among the CPAP users at baseline and one year did not show a significantly different distribution pattern than what was observed in the full cohort of participants.

4. Discussion

This study is one of the first designed to assess the effect of carbohydrate restriction and nutritional ketosis on sleep quality in individuals with hyperglycemia and insulin resistance. Improved patient-reported sleep quality as assessed by global PSQI suggests that CCI including nutritional ketosis benefited sleep quality in both patients with T2D and prediabetes. The proportion of patients categorized as “poor sleepers” at one year was significantly reduced in the CCI groups but not in the UC group. Furthermore, these results demonstrate that the sleep quality improvement observed in the whole intervention population was due in part to 17% of baseline “poor sleepers” being reclassified as “good sleepers” at one year. Our results are consistent with previous findings that showed improved overall sleep quality in children consuming ketogenic diets [33,34].

Improvement in the global PSQI score of patients undergoing the CCI was mainly due to significant changes in three PSQI components: subjective sleep quality, sleep disturbance and daytime dysfunction. Both objective and subjective sleep quality impairment are frequently reported in diabetes patients and positively associated with severity of hyperglycemia [8–11]. Likewise, correlation between poor sleep quality and increased carbohydrate intake [32] has also been previously reported. These observed patterns of association between sleep quality with hyperglycemia and carbohydrate intake may explain why this carbohydrate restriction intervention improved subjective sleep quality. The sleep disturbance component of the global PSQI score is associated with poor glycemic control among T2D patients [45]. One study reported a significant correlation between sleep disturbance and HbA1c level [46]. Night time sleep disturbance in T2D patients can be related to a wide range of conditions such as nocturnal polyuria, pain, and breathing problems, especially in those with OSA. In our study, we also showed that patients encountering persistent pain, including knee pain, had a higher median global PSQI score, while one year of the intervention effectively improved global PSQI scores in these patients despite the persistence of reported pain in some patients. It is possible that improvement in the sleep disturbance of the CCI patients contributed to the glycemic control improvement in these patients. The effectiveness of the intervention in improving sleep in those with pain, further emphasizes its' applicability in alleviating sleep disturbance.

Furthermore, there was a significant improvement in the daytime dysfunction component of the global PSQI score in the CCI group. Excessive daytime sleepiness and dysfunction are reported commonly in T2D [47,48], and weight loss through bariatric surgery has a positive resolving effect on daytime dysfunction and sleepiness [49,50]. In the present investigation, the majority of CCI patients achieved weight loss of $\geq 10\%$, which could have contributed to the significant improvement observed in daytime function. In addition, we also evaluated the effect of the intervention on a subcohort of patients with a self-reported pattern of shifted non-standard bedtimes and wake-up times that were not aligned to the light dark cycle, which likely affects daytime functioning. Circadian rhythm disruption is frequently associated with metabolic alterations, especially in an insulin resistant state [51,52]. While patients with a normal sleep chronotype benefited the most, the intervention also improved the sleep of patients with time shifted bedtimes. A similar advantage of the intervention was not observed in patients with shifted wake-up times, though this may be due to the limited number of patients in this subgroup ($n = 27$).

The improvement in the global PSQI score observed in CCI patients occurred concurrently with weight reduction and glycemic control improvement [36,37]. Martin et al., [53] reported a direct correlation between degree of weight loss and global PSQI score improvement in healthy nonobese adults receiving an energy restricted diet, while Chaput et al., [54] reported an improvement in global PSQI score following the initial 5-kg weight loss, but no additional improvement with subsequent weight loss. A study using a ketogenic diet in children alleviated abnormal sleep architecture; however, weight loss was suggested as the main determinant of improved sleep [35]. These studies collectively imply a direct association between weight loss and improved PSQI score. Likewise, long-term maintenance of weight loss was associated with better sleep quality and quantity [18] while the degree of weight loss reduction is directly correlated with OSA improvement [19]. Alternately, some studies also demonstrate the efficacy of anti-glycemic medications for improving PSQI score concurrent with improved glycemic control [55]. This study identified associations between HOMA-IR and weight reductions with stratification of patients' sleep status in the full CCI cohort even though there

were no significant differences in weight loss and insulin resistance reduction levels between those who had improved sleep and those who did not. Patients with good sleep quality at the beginning of the intervention benefited the most in reducing insulin resistance. Improvement in fasting glucose and HOMA-IR were only positively associated with improved PSQI score in prediabetes patients.

It is not clear if nutritional ketosis achieved by substantial carbohydrate restriction augmented the effect of the intervention on sleep or if weight loss and/or improved glycemic control generated from the intervention contributed to sleep quality improvements. We showed a significant correlation between blood beta-hydroxybutyrate (BHB) levels and PSQI improvement in the prediabetes cohort. While the effect of and mechanism of BHB in sleep are not clear, a positive correlation between blood BHB levels and carbon dioxide (CO₂) response was previously reported in patients with obesity related hypoventilation syndrome that had reduced CO₂ response [56]. A continuous state of ketosis through carbohydrate restriction and fat intake also induces the postprandial release of a satiety hormone, cholecystokinin (CCK) [30,57,58]. When administered in rats, CCK was shown to promote slow wave activity and NREM sleep [59]. CCK was also shown to induce sleep when administered in diabetic rats [60]. Therefore, it is possible that one mechanism of improved sleep with a ketogenic diet that increases BHB levels is through CCK induction.

There are several limitations of our study. The study was designed mainly to assess the impact of the CCI on glycemic control, medication use, weight, and cardiovascular disease risk factors. Patient-reported outcomes for quality of life measures including sleep were included as secondary endpoints. It is difficult to determine the causality among the intervention, related to improvement in primary outcomes and improvement in sleep from this study. A major limitation of this study is the use of subjective sleep measures as self-reported sleep assessment is subject to limited self-knowledge of sleep behavior and inconsistency in reporting. Changes in architecture were not included in the study. Therefore, future studies that use randomized controlled trial designs and objective sleep measures are needed to confirm our results. In addition, patients with an established diagnosis of a sleep disorder such as OSA were not separated in the analysis since complete records of their CPAP usage were not collected in the questionnaire. Patient compliance with CPAP usage is essential for making interpretations about the status of their OSA treatment and its effect on sleep and glycemic control. The study also lacked recruitment of prediabetes patients in the UC group for direct comparison of the treatment effect between UC and CCI on sleep in these patients.

In conclusion, these results demonstrate that overall sleep quality significantly improved in T2D and prediabetes patients undergoing remote CCI including nutritional ketosis but not in T2D patients in the UC group. The sleep improvement was concurrent with weight reduction and glycemic control improvement. The PSQI components that improved were sleep quality, sleep disturbance and daytime dysfunction. These results suggest that nutritional ketosis benefits overall health through improved glycemic control as well as improved sleep quality.

Financial support

The work reported in this study was funded by Virta Health..

Author contributions

S.J.A, M.S, C.J.V and J.P.M drafted the manuscript. A.L.M, N.H.B, S.J.H and S.J.A participated in data acquisition and compiling. M.S and S.J.A analyzed the data. C.J.V supervised this particular analysis,

J.P.M, A.L.M, S.J.H, N.H.B, W.W.C, S.D.P and J.S.D edited the manuscript. W.W.C. proposed measuring subjective sleep quality as part of the parent Continuous Care Intervention clinical trial. All authors approved the final version of the manuscript.

Conflict of interest

SJA, SJH, ALM, NHB, JPM, and SDP are employed by Virta Health Corp and were offered company's stock options. SDP and JSV are founders of Virta Health Corp. CJV, MJS and WWC have no conflict of interest to declare.

The ICMJE Uniform Disclosure Form for Potential Conflicts of Interest associated with this article can be viewed by clicking on the following link: <https://doi.org/10.1016/j.sleep.2018.12.014>.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.sleep.2018.12.014>.

References

- [1] Kuhn E, Brodan V, Brodanova M, et al. Metabolic reflection of sleep deprivation. *Act Nerv Super* 1969;11:165–74.
- [2] Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999;354:1435–9.
- [3] Spiegel K, Knutson K, Leproult R, et al. Sleep loss: a novel risk factor for insulin resistance and type-2 diabetes. *J Appl Physiol* 2005;99:2008–19.
- [4] Nedeltcheva AV, Kessler L, Imperial J, et al. Exposure to recurrent sleep restriction in the setting of high caloric intake and physical inactivity results in increased insulin resistance and reduced glucose tolerance. *J Clin Endocrinol Metab* 2009;94:3242–50.
- [5] Ayas NT, White DP, Al-Delaimy WK, et al. A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes Care* 2003;163:205–9.
- [6] Nilsson PM, Roost M, Engstrom G, et al. Incidence of diabetes in middle-aged men is related to sleep disturbances. *Diabetes Care* 2004;27:2464–9.
- [7] Kawakami N, Takatsuka N, Shimizu H. Sleep disturbance and onset of type 2 diabetes. *Diabetes Care* 2004;27:282–3.
- [8] Lee SW, Ng KY, Chin WK. The impact of sleep amount and sleep quality on glyemic control in type 2 diabetes: a systematic review and meta-analysis. *Sleep Med Rev* 2017;31:91–101.
- [9] Knutson KL, Ryden AM, Mander BA, et al. Role of sleep duration and quality in the risk and severity of type 2 diabetes mellitus. *Arch Intern Med* 2006;166:1768–74.
- [10] Sakamoto R, Yamakawa T, Takahashi K, et al. Association of usual sleep quality and glyemic control in type 2 diabetes in Japanese: a cross sectional study. *Sleep and Food Registry in Kanagawa (SOREKA)*. *PLoS One* 2017. <https://doi.org/10.1371/journal.pone.0191771>.
- [11] Tsai YW, Kann NH, Tung TH, et al. Impact of subjective sleep quality on glyemic control in type 2 diabetes mellitus. *Family Prac* 2012;29:30–5.
- [12] Trento M, Broglio F, Riganti F, et al. Sleep abnormalities in type 2 diabetes may be associated with glyemic control. *Acta Diabetol* 2008;45:225–9.
- [13] Gozashti MH, Eslami N, Radfar MH, et al. Sleep pattern, duration and quality in relation with glyemic control in people with type 2 diabetes mellitus. *Iran J Med Sci* 2016;41:531–8.
- [14] Resnick HE, Redline S, Shafar E, et al. Diabetes and sleep disturbances findings from the sleep heart health study. *Diabetes Care* 2003;26(3):702–9.
- [15] Foster GD, Sander MH, Millman R, et al. Obstructive sleep apnea among obese patients with type 2 diabetes. *Diabetes Care* 2009;32:1017–9.
- [16] American Diabetes Association. 3. Comprehensive medical evaluation and assessment of comorbidities: standards of medical care in diabetes. *Diabetes Care* 2018;41:S28–37.
- [17] The IDF Consensus statement on sleep apnoea and type 2 diabetes. 2008. p. 1–24.
- [18] Yannakoulia M, Anastasiou CA, Karfapoulou E, et al. Sleep quality is associated with weight loss maintenance status: the MedWeight Study. *Sleep Med* 2017;34:242–5.
- [19] Xanthopoulos MS, Berkowitz R, Tapia IE. Effects of obesity therapies on sleep disorders. *Metabolism* 2018;84:109–17.
- [20] Foster GD, Borradaile KD, Sanders MH, et al. A randomized study on the effect of weight loss on obstructive sleep apnea among obese patients with type 2 diabetes: the Sleep AHEAD study. *Arch Intern Med* 2009;169:1619–26.
- [21] Greenburg DL, Lettieri CJ, Eliasson AH. Effects of surgical weight loss on measures of obstructive sleep apnea: a meta-analysis. *Am J Med* 2009;122:535–42.
- [22] Shaw JE, Punjabi NM, Naughton MT, et al. The effect of treatment of obstructive sleep apnea on glyemic control in type 2 diabetes. *Am J Respir Crit Care Med* 2016;194:486–92.
- [23] West SD, Nicoll DJ, Wallace TM, et al. Effect of CPAP on insulin resistance and HbA1c in men with obstructive sleep apnoea and type 2 diabetes. *Thorax* 2017;62:486–92.
- [24] Martinez-Ceron E, Barquiel B, Bezos AM, et al. Effect of continuous positive airway pressure on glyemic control in patients with obstructive sleep apnea and type 2 diabetes. A randomized clinical trial. *Am J Respir Crit Care Med* 2016;194:476–85.
- [25] Pamidi S, Wroblewski K, Stephen M, et al. Eight hours of nightly continuous positive airway pressure treatment of obstructive sleep apnea improves glucose metabolism in patients with prediabetes. A randomized controlled trial. *Am J Respir Crit Care Med* 2015;192:96–105.
- [26] Weinstock TG, Wang X, Rueschman M, et al. A controlled trial of CPAP therapy on metabolic control in individuals with impaired glucose tolerance and sleep apnea. *Sleep* 2012;35:617–25.
- [27] Phillips F, Chen CN, Crisp AH, et al. Isocaloric diet changes and electroencephalographic sleep. *Lancet* 1975;2:723–5.
- [28] Yajima K, Seya T, Iwayama H, et al. Effects of nutrient composition of dinner on sleep architecture and energy metabolism during sleep. *J Nutr Sci Vitaminol* 2014;60:114–21.
- [29] Lindseth G, Lindseth P, Thompson M. Nutritional effects on sleep. *West J Nurs Res* 2013;35:497–513.
- [30] Afaghi A, O'Connor H, Chow CM. Acute effects of the very low carbohydrate diet on sleep indices. *Nutr Neurosci* 2008;11:146–54.
- [31] Kwan RM, Thomas S, Mir MA. Effects of a low carbohydrate isoenergetic diet on sleep behavior and pulmonary functions in healthy female adult humans. *J Nutr* 1986;116:2393–402.
- [32] Katagiri R, Asakura K, Kobayashi S, et al. Low intake of vegetables, high intake of confectionary, and unhealthy eating habits are associated with poor sleep quality among middle-aged female Japanese workers. *J Occup Health* 2014;56:359–68.
- [33] Husain AM, Yancy Jr WS, Carwile ST, et al. Diet therapy for narcolepsy. *Neurology* 2004;62:2300–2.
- [34] Hallbrook T, Lundgren J, Rosen I. Ketogenic diet improves sleep quality in children with therapy-resistant epilepsy. *Epilepsia* 2007;48:59–65.
- [35] Willi SM, Oexmann MJ, Wright NM, et al. The effects of high-protein, low-fat, ketogenic diet on adolescents with morbid obesity, body composition, blood chemistries and sleep abnormalities. *Pediatrics* 1998;101:61–7.
- [36] Hallberg SJ, McKenzie AL, Williams PT, et al. Effectiveness and safety of a novel care model for the management of type 2 diabetes at 1 year: an open-label, non-randomized, controlled study. *Diabetes Ther* 2018;9:583–612.
- [37] Bhanpuri NH, Hallberg SJ, Williams PT, et al. Cardiovascular disease risk factor responses to a type 2 diabetes care model including nutritional ketosis induced by sustained carbohydrate restriction at 1 year: an open label, non-randomized, controlled study. *Cardiovasc Diabetol* 2018;17. <https://doi.org/10.1186/s12933-018-0698-8>.
- [38] McKenzie AL, Hallberg SJ, Bhanpuri NH, et al. Continuous remote care model utilizing nutritional ketosis improves type 2 diabetes risk factors in patients with prediabetes. *Diabetes* 2018; 67 (Supplement 1), <https://doi.org/10.2337/db18-293-OR>.
- [39] Buysse DJ, Reynolds CF, Monk TH, et al. The Pittsburgh Sleep Quality Index (PSQI): a new instrument for psychiatric research and practice. *Psychiatrist Res* 1989;28:193–213.
- [40] Smith MT, Wegener ST. Measures of sleep: the insomnia severity index, medical outcomes study (MOS) sleep scale, Pittsburgh sleep diary (PSD), and Pittsburgh sleep quality index (PSQI). *Arthritis Rheum* 2003;49: S184–9.
- [41] Alexander BA. *BaylorEdPsych: R package for baylor university educational psychology quantitative courses*. R package version 0.5. 2012. <https://CRAN.R-project.org/package=BaylorEdPsych>.
- [42] van Buuren S, Groothuis-Oudshoorn K. Mice: multivariate imputation by chained Equations in R. *J Stat Software* 2011;45:1–67.
- [43] MATLAB and Statistics Toolbox release. Natick, Massachusetts, United States: The MathWorks, Inc.; 2012b.
- [44] R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2018. <https://www.R-project.org/>.
- [45] Song Y, Ye X, Ye L, et al. Disturbed subjective sleep in Chinese females with type 2 diabetes on insulin therapy. *PLoS One* 2013;8(1):e54951. <https://doi.org/10.1371/journal.pone.0054951>.
- [46] Suarez EC. Self-reported symptoms of sleep disturbance and inflammation, coagulation, insulin resistance and psychosocial distress: evidence for gender disparity. *Brain Behav Immun* 2008;22:960–88.
- [47] Chasens ER, Korytkowski M, Sereika SM, et al. Effect of poor sleep quality and excessive daytime sleepiness on factors associated with diabetes self management. *Diabetes Educ* 2013;39:74–82.
- [48] Telford O, Diamantidis CJ, Bosworth HB, et al. The relationship between Pittsburgh Sleep Quality Index subscales and diabetes control. *Chronic Illn* 2018. <https://doi.org/10.1177/1742395318759587>.
- [49] Mello M, Vasques ACJ, Pareja JC, et al. Effect of biliopancreatic diversion on sleep quality and daytime sleepiness in patients with obesity and type 2 diabetes. *Arch Endocrinol Metab* 2017;61:623–7.
- [50] Dilektaşlı E, Dilektaşlı AG. Laparoscopic sleeve gastrectomy improves excessive daytime sleepiness and sleep quality 6 months following surgery: a prospective cohort study. *Adv Ther* 2016;33:774–85.

- [51] Pan A, Schernhammer ES, Sun Q, et al. Rotating night shift work and risk of type 2 diabetes: two prospective cohort studies in women. *PLoS Med* 2011;8: e1001141.
- [52] Reutrakul S, Hood MM, Crowley SJ, et al. Chronotype is independently associated with glycemic control in type 2 diabetes. *Diabetes Care* 2013;36: 2523–9.
- [53] Martin CK, Bhapkar M, Pittas AG, et al. Effect of calorie restriction on mood, quality of life, sleep and sexual function in healthy nonobese adults: the CALERIE 2 Randomized Clinical Trial. *JAMA Intern Med* 2016;176:743–52.
- [54] Chaput JP, Drapeau V, Hetherington M, et al. Psychobiological impact of a progression weight loss program in obese men. *Physiol Behav* 2005;86:224–32.
- [55] Sakamoto Y, Oyama J, Ikeda H, et al. Effects of sitagliptin beyond glycemic control: focus on quality of life. *Cardiovasc Diabetol* 2013;12:35. <https://doi.org/10.1186/1475-2840-12-35>.
- [56] Fried PI, McClean PA, Phillipson EA, et al. Effect of ketosis on respiratory sensitivity to carbon dioxide in obesity. *N Engl J Med* 1976;294:1081–6.
- [57] Hayes MR, Miller CK, Ulbrecht JS, et al. A carbohydrate-restricted diet alters gut peptides and adiposity signals in men and women with metabolic syndrome. *J Nutr* 2007;137:1944–50.
- [58] Chearskul S, Delbridge E, Shulkes A, et al. Effect of weight loss and ketosis on postprandial cholecystokinin and free fatty acid concentrations. *Am J Clin Nutr* 2008;87:1238–46.
- [59] Kapás L, Obal Jr F, Alfoldi P, et al. Effects of nocturnal intraperitoneal administration of cholecystokinin in rats: simultaneous increase in sleep, increase in EEG slow-wave activity, reduction of motor activity, suppression of eating, and decrease in brain temperature. *Brain Res* 1988;438:155–64.
- [60] Kapás L, Obal F, Farkas I, et al. Cholecystokinin promotes sleep and reduces food intake in diabetic rats. *Physiol Behav* 1991;50:417–20.