

2019

Effect of changing urine testing orderables and clinician order sets on inpatient urine culture testing: Analysis from a large academic medical center

Satish Munigala
Washington University School of Medicine in St. Louis

Rebecca Rojek
Barnes-Jewish Hospital, St Louis

Helen Wood
Barnes-Jewish Hospital, St Louis

Melanie L. Yarbrough
Washington University School of Medicine in St. Louis

Ronald R. Jackups
Washington University School of Medicine in St. Louis

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Please let us know how this document benefits you.

Recommended Citation

Munigala, Satish; Rojek, Rebecca; Wood, Helen; Yarbrough, Melanie L.; Jackups, Ronald R.; Burnham, Carey-Ann D.; and Warren, David K., "Effect of changing urine testing orderables and clinician order sets on inpatient urine culture testing: Analysis from a large academic medical center." *Infection Control & Hospital Epidemiology*. 40, 3. 281-286. (2019).
https://digitalcommons.wustl.edu/open_access_pubs/7865


This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Satish Munigala, Rebecca Rojek, Helen Wood, Melanie L. Yarbrough, Ronald R. Jackups, Carey-Ann D. Burnham, and David K. Warren

Original Article

Effect of changing urine testing orderables and clinician order sets on inpatient urine culture testing: Analysis from a large academic medical center

Satish Munigala MBBS, MPH¹, Rebecca Rojek MPH², Helen Wood RN, MA, CIC², Melanie L. Yarbrough PhD³, Ronald R. Jackups Jr MD, PhD³, Carey-Ann D. Burnham PhD³ and David K. Warren MD, MPH¹ 

¹Division of Infectious Diseases, Department of Medicine, Washington University School of Medicine, St Louis, Missouri, ²Department of Hospital Epidemiology and Infection Prevention, Barnes-Jewish Hospital, St Louis, Missouri and ³Department of Pathology and Immunology, Washington University School of Medicine, St Louis, Missouri

Abstract

Objective: To evaluate the impact of changes to urine testing orderables in computerized physician order entry (CPOE) system on urine culturing practices.

Design: Retrospective before-and-after study.

Setting: A 1,250-bed academic tertiary-care referral center.

Patients: Hospitalized adults who had ≥ 1 urine culture performed during their stay.

Intervention: The intervention (implemented in April 2017) consisted of notifications to providers, changes to order sets, and inclusion of the new urine culture reflex tests in commonly used order sets. We compared the urine culture rates before the intervention (January 2015 to April 2016) and after the intervention (May 2016 to August 2017), adjusting for temporal trends.

Results: During the study period, 18,954 inpatients (median age, 62 years; 68.8% white and 52.3% female) had 24,569 urine cultures ordered. Overall, 6,662 urine cultures (27%) were positive. The urine culturing rate decreased significantly in the postintervention period for any specimen type (38.1 per 1,000 patient days preintervention vs 20.9 per 1,000 patient days postintervention; $P < .001$), clean catch (30.0 vs 18.7; $P < .001$) and catheterized urine (7.8 vs 1.9; $P < .001$). Using an interrupted time series model, urine culture rates decreased for all specimen types ($P < .05$).

Conclusions: Our intervention of changes to order sets and inclusion of the new urine culture reflex tests resulted in a 45% reduction in the urine cultures ordered. CPOE system format plays a vital role in reducing the burden of unnecessary urine cultures and should be implemented in combination with other efforts.

(Received 21 September 2018; accepted 11 December 2018)

Urinalysis and urine culture are commonly ordered tests among hospitalized patients suspected of urinary tract infection (UTI). However, these tests are often ordered for patients without clinical suspicion of UTI, leading to unnecessary testing and increased hospital costs.^{1–3} Positive urine cultures are a major driver for antibiotic treatment.^{4–11} Several studies have reported that the treatment of asymptomatic bacteriuria (ASB) does not affect patient outcomes and leads to unnecessary antibiotic use, increasing the prevalence of antibiotic-resistant organisms and *Clostridium difficile* infection.^{12–14} Despite Infectious Disease Society of America and other professional societies' recommendations to avoid antibiotic prescriptions for asymptomatic bacteriuria,^{14–17} its treatment is still common.

Previous interventions to prevent unnecessary urine testing have included provider education, use of pocket cards, antimicrobial stewardship efforts, reflex urine culture cancellation and 2-step urine culture ordering.^{6,7,12,13,18–22} However, data on the effect of changes in electronic order sets and its role on inpatient urine testing practices are limited.

In this study, we evaluated the impact of changes to the inpatient urine orders in a computer physician order entry (CPOE) system on the urine culturing practices of a large urban, academic medical center.

Methods

Setting

This retrospective before-and-after study included patients admitted to Barnes-Jewish Hospital (BJH), a 1,250-bed teaching hospital, from January 1, 2015, to August 31, 2017, who had ≥ 1 urine culture ordered during their stay. Patients who were admitted during the

Author for correspondence: David K. Warren, Email: dwarren@wustl.edu

Cite this article: Munigala S, *et al.* (2019). Effect of changing urine testing orderables and clinician order sets on inpatient urine culture testing: Analysis from a large academic medical center. *Infection Control & Hospital Epidemiology*, 40: 281–286, <https://doi.org/10.1017/ice.2018.356>

Table 1. Urine Order Set Definitions During the Study Period

Preintervention (January 2015–April 2016)		Postintervention (May 2016–August 2017)	
Test Name(s)	Definition	Test name(s)	Definition
		UA Reflex to Microscopy WITH Culture	If urinalysis is positive for nitrites OR leukocyte esterase, then microscopy and urine culture will automatically be performed.
Urine Flex (Urine Macroscopic UA Flex) UA Flex C/S (Urine Macroscopic UA Flex) ^a	Perform urine dipstick. If positive for any protein >trace, blood, nitrite, or leukocyte esterase, then proceed to microscopy and culture.	UA Reflex for Neutropenic Patients	If urinalysis is positive for protein (>trace), blood, nitrites, OR leukocyte esterase, then microscopy and urine culture will automatically be performed.
UA Reflex UA W/Reflexed Microscopic (UA Reflex) ^b	Perform urine dipstick. If positive for any protein >trace, blood, nitrite, or leukocyte esterase, then proceed to microscopy.	UA Reflex to Microscopy WITHOUT Culture	If urinalysis is positive for protein (>trace), blood, nitrites, OR leukocyte esterase, then microscopy will automatically be performed.
Urine Macro Urinalysis UA Macro UA Dip Macroscopic ^c	Macroscopic dipstick urinalysis only	Urine Macro Urinalysis UA Macro UA Dip Macroscopic	Macroscopic dipstick urinalysis only
UA Microscopy UA Micro ^d	Urine sediment examination only	UA Microscopy UA Micro	Urine sediment examination only
Urine culture, X (X = aerobic, fungal, mycobacterial)		Urine culture, X (X = aerobic, fungal, mycobacterial)	

Note. UA, urinalysis.

^aUrine Flex and Urine Flex C/S were the same orders with different names during the preintervention period;

^bUA Reflex and UA W/Reflexed were the same orders with different names during the preintervention period.

^cUrine Macro, Urinalysis, UA Macro and UA Dip Macroscopic were the same orders with different names during the study period.

^dUA Microscopy and UA Micro were the same orders with different names during the study period.

study period but did not have a urine culture ordered during their stay and patients who had their urine cultures obtained at an outpatient settings or the emergency department (ED) were excluded.

Intervention

A staged intervention was performed to clarify test names and to reduce the number of reflex urine cultures performed for nonspecific indications (eg, isolated proteinuria) by making changes to the urine reflex test panel at BJH (Table 1). This intervention was initiated in the CPOE system on January 28, 2016. E-mail notification to providers with the new urine reflex tests was sent prior to initiation. The inclusion of the new reflex tests in commonly used order sets within the CPOE system (eg, medical intensive care unit admission orders) was completed on April 19, 2016; therefore, April 2016 was used as the intervention month. January 2015 through April 2016 was the preintervention period and May 2016 through August 2017 was the postintervention period.

Data collection

Patient and laboratory data were abstracted from the hospital medical informatics database. Data included patient demographics (ie, age, race, and sex), laboratory test results (ie, urinalysis, microscopic exam and urine culture), and discharge disposition (ie, home, other facility, etc). For urine cultures with accompanying urinalysis or microscopy, the time between the urine culture and urinalysis and/or microscopy was calculated. Type of urine culture specimen was also noted (ie, clean catch, catheterized, or procedure related) as indicated by the ordering clinician.

For patients with multiple urine cultures during an admission, each sample was treated as an independent observation.

Definitions

Urine cultures with growth of $\geq 100,000$ colony-forming units (CFU)/mL for a clean-catch specimen and $\geq 10,000$ CFU/mL for a catheterized specimen were considered positive results. Urine cultures that were negative for significant growth or contaminated were considered negative for this analysis. Leukocyte esterase ≥ 1 identified on urinalysis and >5 white blood cells per high-power field on urine microscopy were treated as abnormal and positive test results. We defined an isolated urine culture as a culture without an associated urinalysis and/or urine microscopy performed within 1 calendar day before or after the culture was performed. Catheter-associated urinary tract infection (CAUTI) surveillance was independently conducted by the hospital infection prevention department during the study period. A CAUTI was defined according to National Healthcare Safety Network definitions²³ as a UTI where an indwelling urinary catheter was in place for >2 calendar days on the date of event, with day of device placement being day 1, and an indwelling urinary catheter was in place on the date of event or the day before. If an indwelling urinary catheter was in place for >2 calendar days and then removed, the UTI criteria must have been fully met on the day of discontinuation or the next day.

Cost assessment

Unit cost of a urine culture was obtained from the Medicare Clinical Laboratory Fee Schedule using national median Medicare payment

Table 2. Characteristics of Patients With ≥ 1 Urine Cultures Obtained From Inpatients

Variable	Study Cohort ^a (N = 18,954), No. (%)	Preintervention (N = 11,780), No. (%) ^b	Postintervention (N = 7,174), No. (%) ^c	P Value
Age, median (IQR)	62 (49–72)	61 (48–72)	62 (49–72)	.015
Race				
White	13,043 (68.8)	8,098 (68.7)	4,945 (68.9)	Reference
Black	4,791 (25.3)	2,926 (24.8)	1,865 (26.0)	.217
Other	1,120 (5.9)	756 (6.4)	364 (5.1)	<.001
Sex				
Male	9,040 (47.7)	5,702 (48.4)	3,338 (46.5)	.012
Female	9,914 (52.3)	6,078 (51.6)	3,836 (53.5)	Reference
Median urine culture per admission (range)	1 (1–12)	1 (1–12)	1 (1–12)	
Discharge status^d				
Discharged to home	12,336 (65.6)	7,762 (66.6)	4,574 (63.9)	<.001
Discharged to other facility	4,805 (25.5)	2,890 (24.8)	1,915 (26.7)	Reference
Other	1,670 (8.9)	999 (8.6)	671 (9.4)	.815

Note. IQR, interquartile range.

^aEach admission is treated as an observation for this analysis purpose. Study cohort includes patients admitted to hospital whose urine was tested (≥ 1) for culture at the hospital during the study period (does not include urine cultures performed at the emergency department, a 24/7 clinic, or in any outpatient setting). Overall, 18,954 patient admissions contributed to 24,569 urine culture tests during the study period (median, 1; IQR, 1–1).

^bJanuary 2015 to April 2016.

^cMay 2016 to August 2017.

^dOverall: missing (n = 143); other: still in hospital (n = 9); died (n = 1,532); left against advice (n = 106); unknown (n = 23). Preintervention: missing (n = 129); still in hospital (n = 5); admitted to hospital (n = 12); died (n = 910); left against advice (n = 72); Postintervention: missing (n = 14); still in hospital (n = 4); died (n = 622); left against advice (n = 34); unknown (n = 11).

Table 3. Comparison of Urine Culture Testing Practices Before and After the Intervention

Variable	Total (n = 24,569)	Preintervention (n = 15,746) ^a	Postintervention (n = 8,823) ^b	P Value
Positive cultures, No. (%) ^c	6,642 (27.0)	4,021 (25.5)	2,621 (29.7)	<.001
Isolated cultures, No. (%) ^c	6,240 (25.4)	4,101 (26.0)	2,139 (24.2)	.002
Urine cultures per 1,000 patient days ^d	29.4	38.1	20.9	<.001
Catheterized urine cultures per 1,000 patient days	4.8	7.8	1.9	<.001
Catheter associated UTI, No. (%)	250 (1.0)	125 (0.8)	125 (1.4)	
CAUTI per 1,000 patient days	0.30	0.30	0.30	.871
CAUTI per 1,000 catheter days	1.26	1.25	1.27	.899

Note: UTI, urinary tract infection; CAUTI, catheter associated urinary tract infection.

^aJanuary 2015 to April 2016.

^bMay 2016 to August 2017.

^cSee methods for definitions.

^dBased upon 413,137 patient days preintervention and 421,714 patient days postintervention for all patients admitted to the hospital during the study period (see Methods).

rate of \$15.00 per urine culture (not adjusted to inflation).²⁴ Total laboratory charges for urine cultures during the preintervention and postintervention periods were calculated, and cost difference was estimated.

Statistical analysis

Patient demographics and characteristics are reported on a per-admission basis. Urine cultures rates are reported per 1,000 patient days (ie, the total patient days for all patients admitted during the study period). CAUTI rates are reported per 1,000 patient days and catheter days. Demographic characteristics and urine culture data were compared for the preintervention period and the postintervention period using the Wilcoxon rank-sum test, χ^2 test, or univariable logistic regression where appropriate. An interrupted time-series model was used to analyze the impact of the intervention on urine culture rates during the study period. Data were

analyzed using SAS version 9.3 software (SAS Institute, Cary, NC). The Washington University Human Research Protection Office approved this study.

Results

Patient characteristics

During the study period, 18,954 patients had ≥ 1 urine culture ordered during their hospital stay (11,780 during the preintervention vs 7,174 during the postintervention period) (Table 2). The median age of the patients was 62 years; ~69% of patients were white and 52.3% were female. Also, ~66% of these patients were routinely discharged home and 25.5% were discharged to or transferred to other facilities. Patients in the preintervention period were slightly younger (61 years preintervention vs 62 years postintervention; $P = .015$), were predominantly male (48.4% vs




Fig. 1. Urine culture rate by specimen type. **P* value for clean-catch and catheterized cultures. Note. The preintervention period was January 2015 to April 2016 and the postintervention period was May 2016 to August 2017.

46.5%; $P = .012$), and were routinely discharged home (66.6% vs 63.9%; $P < .001$) compared to the postintervention period.

Urine culture characteristics

A total of 24,569 urine cultures were ordered (during 18,954 admissions at the rate of 29.4 cultures per 1,000 patient days; median, 1 urine culture per admission) during the study period. Of these, 70.7% had an associated urinalysis and 70.4% had an associated microscopy (25.4% of urine cultures were deemed to be isolated). Overall, 6,642 urine cultures (27%) were positive. The proportion of positive urine cultures increased in the postintervention period (25.5% preintervention vs 29.7% postintervention; $P < .001$), whereas the proportion of isolated urine cultures decreased (26.0% preintervention vs 24.2% postintervention; $P = .002$) (Table 3).

Urine culture rates by specimen type

Urine culture decreased by 45.1% in the postintervention period (38.1 per 1,000 patient days preintervention vs 20.9 per 1,000 patient days postintervention; $P < .001$) (Table 3). This decrease was observed for clean catch (30.0 per 1,000 patient days preintervention vs 18.7 per 1,000 patient days postintervention; $P < .001$) and catheterized urine cultures (7.8 per 1,000 patient days preintervention vs 1.9 per 1,000 patient days postintervention; $P < .001$), whereas procedure-related urine cultures remained stable at 0.3 per 1,000 patient days (Fig. 1).

When adjusted for impact of the intervention using an interrupted time series model, urine culture rates decreased significantly for overall ($P < .001$), catheterized ($P < .001$), and isolated cultures ($P = .027$), respectively (Fig. 2).

Catheter associated urinary tract infections (CAUTI)

Overall, 250 CAUTIs were identified during the study period (0.30 per 1,000 patient days); however, after the intervention there was no significant change in the CAUTI rates (0.30 per 1,000 patient days preintervention vs 0.30 per 1,000 patient days postintervention;

$P = .871$; 1.25 per 1,000 catheter days preintervention vs 1.27 per 1,000 catheter days postintervention; $P = .899$) (Table 3).

Effect of intervention on laboratory costs

Our intervention resulted in a \$6,490 reduction in the mean monthly laboratory cost during the postintervention period, with an estimated total cost savings of \$103,345 for inpatient urine culture laboratory costs in the postintervention period (\$236,190 preintervention vs \$132,345 in the postintervention period).

Discussion

In this retrospective study, we observed a 45.1% unadjusted decrease in the rate of inpatient urine cultures performed because of changes to electronic orders in the computer physician order entry system. The reduction in the urine culture rate was most marked for the catheterized (75.6%) compared to a clean-catch specimens (37.8%). We also noticed a 16.4% increase in the proportion of positive urine cultures and a 6.9% decrease in the proportion of isolated urine cultures obtained. Overall, our intervention resulted in an estimated reduction of \$103,845 in laboratory charges to patients.

Unnecessary ordering of urine cultures and inappropriate antimicrobial use for asymptomatic bacteriuria remain common among clinicians.^{13,15-17,25-27} Lack of familiarity with the recommendations, excessive testing in patients with comorbidities, and certain practice patterns among physicians are some of the common factors driving this clinical practice.^{9,28} Moreover, a urine culture result is often difficult for clinicians to ignore and drives antimicrobial therapy regardless of symptoms.²⁹

Several prior efforts to prevent treatment of asymptomatic bacteriuria have included educational sessions,^{6,30} pocket cards with diagnostic algorithms with audit and feedback for training clinicians,¹³ and antimicrobial stewardship efforts. Recently, Hartley et al⁴ replicated these interventions in hospitalist-based service in 3 different hospitals and observed a 24% reduction in ASB treatment rates, resulting in fewer days of antimicrobial therapy. Other recent interventions have included focus groups interviews for identifying factors that affect nurse initiated urine culture ordering and collection practices,³¹ reflex urine culture cancellation,²¹ and 2-step urine culture ordering in the emergency department.²² Although several of these upstream interventions are aimed toward eliminating unnecessary ordering and downstream interventions are aimed toward reducing treatment of asymptomatic bacteriuria, knowledge on the role of CPOE in reducing the burden of unnecessary ordering in the inpatient setting is limited.

Because of our intervention, we also noticed a significant increase in the proportion of urine cultures that were positive during the postintervention period. This finding may indicate increased clarity of reflex algorithm test names and a change in the behavior of ordering clinicians (eg, urine cultures are more likely to be ordered in patients with a higher pretest probability). The postintervention period had a significantly higher proportion of positive urine cultures with an associated abnormal or positive urinalysis (1,896 of 2,621 [72.3%] vs 2,442 of 4,021 [60.7%]; $P < .001$) and a significantly lower proportion of positive urine culture results with an associated negative urinalysis (122 of 2,621 [4.7%] vs 479 of 4,021 [11.9%]; $P < .001$). These findings suggests that a chance of an important UTI having been missed due to the decreased rate of urine culture following the intervention was less unlikely. Although we noticed a significant but small (6.9%) decrease in the isolated urine culture and substantial decrease




Fig. 2. Inpatient urine culturing practices from January 1, 2015 to August 31, 2017. The intervention time point is noted by a dashed line. $P < .001$ for urine cultures and catheterized cultures; $P = .027$ for isolated cultures, using interrupted time series analysis.


(75.6%) in the rate of catheterized urine cultures per 1,000 patient days, there was no significant change in the CAUTI rate postintervention. Given that we previously reported that isolated urine cultures were more likely to be ordered on catheterized patients and patients with prolonged hospital stays,³² we evaluated the proportion of CAUTIs associated with isolated urine cultures. We found no significant difference between study periods in the proportion of CAUTIs that were identified based on isolated urine cultures (39 of 125 [31.2%] preintervention vs 26 of 125 [20.8%] postintervention; $P = .06$). These findings suggest that for patients in whom a clinical suspicion of CAUTI existed, clinicians were ordering diagnostic tests and detecting it in both intervention periods. Therefore, additional infection prevention efforts may be required in this study cohort to prevent CAUTIs.

Our intervention resulted in an estimated cost savings of ~\$104,000 for inpatient laboratory costs after implementation. This represents a fraction of the total costs and does not reflect the costs saved based on the medical decisions (eg, delayed hospital discharge) and antimicrobial therapy.²⁷ In an era of reducing reimbursement for clinical laboratory testing,³³ the prudent use of common diagnostic tests in patient care is increasingly important.

The limitations of our study include a retrospective design, the absence of chart review for test indication, and lack of data on antibiotic use for assessment of antimicrobial therapy. We were unable to assess asymptomatic bacteriuria because data on clinical symptoms or signs were not collected. In addition, this is a single academic medical center and may not be generalizable to other settings. Our medical informatics database does not include orders; therefore, we were unable to directly evaluate the frequency of urinalysis reflex to microscopy with culture and types of urine culture orders. We attempted to address this limitation by examining urine cultures that were performed along with urinalysis and/or microscopy, but we would not be able to identify how much our intervention reduced the proportion of urinalysis that reflexed to culture. The median number of urine cultures for the preintervention and postintervention periods were the same (including demographic characteristics patients who had >1 urine culture); therefore, we did not make any adjustments for the repeat observations. We were unable to directly assess whether antibiotic use changed in patients with urinary testing because of the intervention, and its subsequent effect on antimicrobial resistance among urinary pathogens. Strengths of our study include using data from

a large academic medical center and electronic order sets for the intervention. The use of CPOE for such intervention requires relatively little ongoing intervention effort compared with other diagnostic stewardship efforts, which require constant monitoring. Our study results complement a similar CPOE intervention conducted in the emergency department of the same hospital, where we observed a 47% decrease in urine cultures ordered when only “urinalysis with reflex to microscopy” was retained in the frequently ordered list of laboratory tests.³⁴ A similar study of urine diagnostics reported that the elimination of reflexed microscopy examination for inpatient locations resulted in a 95% reduction in the urine microscopy performed.³⁵

In summary, a staged intervention to clarify test names and inclusion of new reflex tests resulted in a 45% reduction in the urine cultures ordered with an estimated cost savings of \$104,000. Further studies are needed to evaluate the role of CPOE in combination with education sessions for ordering physicians and antimicrobial stewardship efforts in reducing the incidence of unnecessary urine cultures. Future research should also focus on reducing isolated urine cultures and CAUTIs.

Author ORCIDs. David K. Warren,  0000-0001-8679-8241

Acknowledgements. None.

Financial support. This study was funded in part by Centers for Disease Control and Prevention Epicenters Program (grant no. 1U54CK000482-01).

Conflicts of interest. All authors report no conflict of interest related to this manuscript.

References

1. Trautner BW. Asymptomatic bacteriuria: when the treatment is worse than the disease. *Nat Rev Urol* 2011;9:85–93.
2. McMaster-Baxter NL, Musher DM. Clostridium difficile: recent epidemiologic findings and advances in therapy. *Pharmacotherapy* 2007;27:1029–1039.
3. Saint S. Clinical and economic consequences of nosocomial catheter-related bacteriuria. *Am J Infect Control* 2000;28:68–75.
4. Hartley SE, Kuhn L, Valley S, et al. Evaluating a hospitalist-based intervention to decrease unnecessary antimicrobial use in patients with asymptomatic bacteriuria. *Infect Control Hosp Epidemiol* 2016;37:1044–1051.

5. Cope M, Cevallos ME, Cadle RM, Darouiche RO, Musher DM, Trautner BW. Inappropriate treatment of catheter-associated asymptomatic bacteriuria in a tertiary care hospital. *Clin Infect Dis* 2009;48:1182–1188.
6. Pavese P, Saurel N, Labarere J, *et al*. Does an educational session with an infectious diseases physician reduce the use of inappropriate antibiotic therapy for inpatients with positive urine culture results? A controlled before-and-after study. *Infect Control Hosp Epidemiol* 2009;30:596–599.
7. Kelley D, Aaronson P, Poon E, McCarter YS, Bato B, Jankowski CA. Evaluation of an antimicrobial stewardship approach to minimize overuse of antibiotics in patients with asymptomatic bacteriuria. *Infect Control Hosp Epidemiol* 2014;35:193–195.
8. Yin P, Kiss A, Leis JA. Urinalysis orders among patients admitted to the general medicine service. *JAMA Intern Med* 2015;175:1711–1713.
9. Hartley S, Valley S, Kuhn L, *et al*. Overtreatment of asymptomatic bacteriuria: identifying targets for improvement. *Infect Control Hosp Epidemiol* 2015;36:470–473.
10. Grein JD, Kahn KL, Eells SJ, *et al*. Treatment for positive urine cultures in hospitalized adults: a survey of prevalence and risk factors in three medical centers. *Infect Control Hosp Epidemiol* 2016;37:319–326.
11. Trautner BW, Bhimani RD, Amspoker AB, *et al*. Development and validation of an algorithm to recalibrate mental models and reduce diagnostic errors associated with catheter-associated bacteriuria. *BMC Med Inform Decis Mak* 2013;13:48.
12. Linares LA, Thornton DJ, Strymish J, Baker E, Gupta K. Electronic memorandum decreases unnecessary antimicrobial use for asymptomatic bacteriuria and culture-negative pyuria. *Infect Control Hosp Epidemiol* 2011;32:644–648.
13. Trautner BW, Grigoryan L, Petersen NJ, *et al*. Effectiveness of an antimicrobial stewardship approach for urinary catheter-associated asymptomatic bacteriuria. *JAMA Intern Med* 2015;175:1120–1127.
14. Gupta K, Hooton TM, Naber KG, *et al*. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin Infect Dis* 2011;52:e103–e120.
15. Nicolle LE, Bradley S, Colgan R, Rice JC, Schaeffer A, Hooton TM. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis* 2005;40:643–654.
16. Nicolle LE. Asymptomatic bacteriuria: review and discussion of the IDSA guidelines. *Int J Antimicrob Agents* 2006;28(suppl 1):S42–S48.
17. Choosing Wisely: an initiative of the ABIM Foundation. 2015. <http://www.choosingwisely.org/doctor-patient-lists/infectious-diseases-society-of-america/>. Accessed March 3, 2018.
18. Bonnal C, Baune B, Mion M, *et al*. Bacteriuria in a geriatric hospital: impact of an antibiotic improvement program. *J Am Med Dir Assoc* 2008;9:605–609.
19. Chowdhury F, Sarkar K, Branche A, *et al*. Preventing the inappropriate treatment of asymptomatic bacteriuria at a community teaching hospital. *J Community Hosp Intern Med Perspect* 2012;2:(pii)17814.
20. Leis JA, Palmay L, Elligsen M, Walker SA, Lee C, Daneman N. Lessons from audit and feedback of hospitalized patients with bacteriuria. *Am J Infect Control* 2014;42:1136–1137.
21. Jones CW, Culbreath KD, Mehrotra A, Gilligan PH. Reflect urine culture cancellation in the emergency department. *J Emerg Med* 2014;46:71–76.
22. Stagg A, Lutz H, Kirpalaney S, *et al*. Impact of two-step urine culture ordering in the emergency department: a time series analysis. *BMJ Qual Saf* 2018;27:140–147.
23. Catheter-associated urinary tract infection (CAUTI) event. National Health Safety Network Patient Safety Manual; 2018. Centers for Disease Control and Prevention website. https://www.cdc.gov/nhsn/pdfs/validation/2018/pcsmanual_2018-508.pdf. Published 2018. Accessed March 8, 2018.
24. Clinical laboratory fee schedule, 2017. Centers for Medicare and Medicaid Services website. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/ClinicalLabFeeSched/Clinical-Laboratory-Fee-Schedule-Files-Items/17CLAB.html?DLPage=1&DLEntries=1008&DLSort=2&DLSortDir=descending>. Updated January 2017. Accessed December 2018.
25. Tomford JW, Hershey CO. The i.v. therapy team: impact on patient care and costs of hospitalization. *NITA* 1985;8:387–389.
26. Gupta K, Hooton TM, Naber KG, *et al*. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin Infect Dis* 2011;52:e103–e120.
27. Forsman RW. Why is the laboratory an afterthought for managed care organizations? *Clin Chem* 1996;42:813–816.
28. Trautner BW, Petersen NJ, Hysong SJ, Horwitz D, Kelly PA, Naik AD. Overtreatment of asymptomatic bacteriuria: identifying provider barriers to evidence-based care. *Am J Infect Control* 2014;42:653–658.
29. Leis JA, Rebick GW, Daneman N, *et al*. Reducing antimicrobial therapy for asymptomatic bacteriuria among noncatheterized inpatients: a proof-of-concept study. *Clin Infect Dis* 2014;58:980–983.
30. Irfan N, Brooks A, Mithoowani S, Celetti SJ, Main C, Mertz D. A controlled quasi-experimental study of an educational intervention to reduce the unnecessary use of antimicrobials for asymptomatic bacteriuria. *PLoS One* 2015;10:e0132071.
31. Redwood R, Knobloch MJ, Pellegrini DC, Ziegler MJ, Pulia M, Safdar N. Reducing unnecessary culturing: a systems approach to evaluating urine culture ordering and collection practices among nurses in two acute care settings. *Antimicrob Resist Infect Control* 2018;7:4.
32. Carlson AL, Munigala S, Russo AJ, *et al*. Inpatient urine cultures are frequently performed without urinalysis or microscopy: findings from a large academic medical center. *Infect Control Hosp Epidemiol* 2017;38:455–460.
33. Information regarding the final CY 2018 private payor rate-based clinical laboratory fee schedule (CLFS) payment rates. Centers for Medicare and Medicaid website. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/ClinicalLabFeeSched/Downloads/CY2018-CLFS-HCPCS-Median-Calculations.pdf>. Published 2018. Accessed December 2018.
34. Munigala S, Jackups RR Jr, Poirier RF, *et al*. Impact of order set design on urine culturing practices at an academic medical centre emergency department. *BMJ Qual Saf* 2018;27:587–592.
35. Chen M, Eintracht S, MacNamara E. Successful protocol for eliminating excessive urine microscopies: quality improvement and cost savings with physician support. *Clin Biochem* 2017;50:88–93.