

Washington University School of Medicine

Digital Commons@Becker

Open Access Publications

2020

Anellovirus dynamics are associated with primary graft dysfunction in lung transplantation

Joshua A. Blatter

Tsuyoshi Takahashi

Brigitte Mittler

Ruben G. Nava

Varun Puri

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/open_access_pubs

Authors

Joshua A. Blatter, Tsuyoshi Takahashi, Brigitte Mittler, Ruben G. Nava, Varun Puri, Daniel Kreisel, and David Wang

after allograft implantation in PGD cases when compared with controls.¹¹ In pediatric patients, alphatorquevirus levels at 2 weeks posttransplant have already been shown to predict acute cellular rejection (ACR),¹² another key short-term adverse outcome in lung transplant recipients. In the current study, we sought to examine whether these dynamic patterns of alphatorquevirus associated with PGD and ACR in previous studies could be validated in a cohort of adult lung transplant recipients.

MATERIALS AND METHODS

Participant Description

The participants in this study were recruited as part of the lung transplant virome cohort. From July 2017 to August 2018, a total of 64 consecutive patients (19 to 75 y of age, average 58 y) were enrolled at the Washington University/Barnes-Jewish Transplant Center in Saint Louis, Missouri. Induction immunosuppression was standardized, with 62 patients receiving basiliximab and methylprednisolone, and 2 patients with evidence of donor-specific antibodies receiving thymoglobulin and methylprednisolone. Maintenance immunosuppression, including tacrolimus, mycophenolate, and prednisone, was applied per protocol, as was antiviral prophylaxis. This study was approved by the Washington University in St. Louis Institutional Review Board (ID 201706125).

Anellovirus PCR Analysis

Nucleic acid was extracted from whole blood samples using the COBAS AmpliPrep Total Nucleic Acid Isolation Kit (Roche Diagnostics), according to manufacturer's instructions. Before extraction, 425 μ L of whole blood was mixed with an equal volume of the COBAS Specimen Pre-Extraction Reagent. Samples were eluted in 75 μ L of buffer.

Alphatorquevirus levels in extracted samples were quantified using TaqMan quantitative real-time PCR, targeting a conserved segment of the viral untranslated region.^{12,13} The 3 μ L extracted sample, TaqMan Fast Advanced Master Mix (Thermo Fisher), 0.9 μ M AMTS forward primer (5' GTGCCGIAGGTGAGTTTA 3'), 0.9 μ M AMTS reverse primer (5' AGCCCGGCCAGTCC 3'), and 250 nM AMPTU probe (5' TCAAGGGGCAATTCGGGCT 3', 6-FAM/ZEN/3'IBFQ) were combined in 25 μ L total reaction volume. Cycling conditions were 50°C for 2 minutes and 95°C for 20 seconds, followed by 40 cycles of 95°C for 1 second, and 60°C for 20 seconds. Betatorquevirus levels in extracted samples were likewise quantified using TaqMan quantitative real-time PCR, targeting a short segment near the ORF 2.^{12,14} The 3- μ L extracted sample, TaqMan Fast Advanced Master Mix, 0.5 μ M forward primer (5' AGTTTATGCCGCCAGACG 3'), 0.5 μ M reverse primer (5' CCCTAGACTTCGGTGGTTTC 3'), and 250 nM probe (5' ACTCACCTHCGGCACCCGC 3', 6-FAM/TAMRA). Cycling conditions were 95°C for 20 seconds, followed by 40 cycles of 95°C for 1 second and 60°C for 20 seconds. The ViiA 7 Real-Time PCR System (Applied Biosystems) was used with default ViiA 7 Software v1.2 settings. A plasmid control for each assay was used to generate a standard curve (alphatorquevirus: slope -3.67 , R^2 0.99; betatorquevirus: slope -3.83 , R^2 0.99), and the detection limit was 30 and 3 copies for alphatorquevirus and betatorquevirus,

respectively. Samples below the detection limit were assigned a value at the bottom of rank order.

Assignment of Clinical Metadata

The participants' pretransplant listing diagnoses were assigned to Group A (obstructive disease), Group B (pulmonary vascular disease), Group C (cystic fibrosis), and Group D (restrictive disease) per current United Network of Organ Sharing policy.¹⁵ The diagnosis of PGD was based on consensus guidelines developed by The International Society for Heart & Lung Transplantation Working Group on Primary Graft Dysfunction.³ High-grade PGD was assessed as Grade 2 or 3 disease at any time point within the first 72 hours following transplant. ACR was diagnosed using transbronchial biopsies obtained at 1, 2, and 3 months following transplant. The definition of ACR was likewise based on International Society for Heart & Lung Transplantation consensus guidelines,¹⁶ and presence of ACR within the first 3 months (ie, "ever ACR") was defined as Grade 1 disease or above. If patients had insufficient data (ie, <2 monthly transbronchial biopsy specimens obtained within the first 3 mo), no assignment of "ever ACR" was made.

Statistical Analyses

Alphatorquevirus and betatorquevirus levels before and after transplant were compared with Spearman correlation. Pairwise comparison of PGD incidence among the pretransplant diagnoses was made with Fisher exact testing. Bivariate analyses of alphatorquevirus copy number and outcomes of interest were assessed with Mann-Whitney testing. The rate of rise of alphatorquevirus was assessed as a copy number ratio between later and earlier time points, and ratios were compared by Mann-Whitney testing. Statistical analysis was performed with GraphPad Prism version 8.

RESULTS

Study Participants

The characteristics of lung transplant virome study participants are summarized in Table 1. The most common diagnoses were Group A and Group D. High-grade PGD was more prevalent among patients with restrictive than obstructive disease ($P < 0.003$). Of the 64 patients included in the study,




FIGURE 1. Levels of recipient blood alphatorquevirus, 1 day before transplant (pretransplant) and 1–2 days following transplant (posttransplant).

TABLE 1.
Characteristics of LTV cohort

Characteristic	Prevalence (n [%]), or mean (SD)	PGD incidence (n [%])
Female sex	35 (55%)	17 (48%)
Age at transplant	57.9 (14.0)	n/a
Ever primary graft dysfunction (PGD)	27 (42%)	n/a
Ever high-grade PGD (2 or above)	18 (28%)	n/a
Ever ACR	25 (49%)	9 (36%)
Pretransplant diagnosis		
Obstructive disease (Group A)	29 (45%)	6 (21%)
Pulmonary vascular disease (Group B)	3 (5%)	2 (67%)
Cystic fibrosis (Group C)	6 (9%)	2 (33%)
Restrictive disease (Group D)	26 (41%)	16 (62%)

ACR, acute cellular rejection; PGD, primary graft dysfunction; SD, standard deviation.

52 patients had blood samples needed for the assessment of anellovirus rise in the first week posttransplant. Of the 64 patients included in the study, 51 had sufficient data for “ever ACR” assessment. Five of 64 patients died in the first year following their transplant.

Alphatorquevirus and Betatorquevirus Levels in the First Week Following Transplant

Figure 1 shows levels of recipient blood alphatorquevirus 1 day before transplant and 1–2 days after transplant. There is a significant correlation ($P < 0.0001$) between alphatorquevirus levels at these timepoints. Betatorquevirus levels were likewise correlated ($P < 0.0001$, analysis not shown). In Figure 2, the rate of alphatorquevirus rise is expressed as a ratio of 1 week

posttransplant, when compared with 1 day before transplant. Patients with “ever-high PGD” (Grade 2+) had a significantly slower rate of rise in alphatorquevirus when compared with patients without high-grade PGD ($P = 0.031$). We generated a logistic regression model to assess effect size and confounding, but when we applied this technique, there was no significant association between alphatorquevirus and PGD. The association between ever-high PGD and betatorquevirus was not significant ($P = 0.64$). There was also no association between alphatorquevirus or betatorquevirus rate of rise and the subsequent development of ACR (alphatorquevirus, $P = 0.12$; betatorquevirus, $P = 0.59$).

Alphatorquevirus Levels in the Subsequent Month Following Transplant

Because our prior study had shown an association between 2 weeks posttransplant blood alphatorquevirus levels and the subsequent development of ACR, we investigated that relationship in this cohort, but there was not a significant association ($P = 0.55$). Since there was a statistical association between immediate posttransplant alphatorquevirus rate of rise and PGD, we evaluated anellovirus expansion over the first 5 weeks following transplant, and evaluated dynamic trends. Patients with prior PGD had a nonsignificant rise in alphatorquevirus level between 1 and 3 weeks posttransplant (Figure 3). Patients with a more rapid rise in alphatorquevirus level between 1 and 5 weeks posttransplant also had a nonsignificant increase in the incidence of “ever ACR” in the first 3 months following transplant (Figure 4).

DISCUSSION

We determined that blood anellovirus levels are associated with PGD in lung transplant recipients. The finding that

Ratio of 1wk post-/pre-transplant alphatorquevirus, by ever-highPGD (2+) status




FIGURE 2. Levels of recipient blood alphatorquevirus, expressed as a ratio of 1 week posttransplant, when compared with 1 day before transplant, among patients without and with high-grade PGD (Grade 2+). PGD, primary graft dysfunction.

Ratio of 3 vs. 1wk post-transplant alphatorquevirus, by everPGD status




FIGURE 3. Levels of recipient blood alphatorquevirus, expressed as a ratio of 3 week posttransplant, when compared with 1 week after transplant, among patients without and with PGD. PGD, primary graft dysfunction.

Ratio of 5 vs. 1-wk post-transplant alphatorquevirus, by everACR status




FIGURE 4. Levels of recipient blood alphatorquevirus, expressed as a ratio of 5 week posttransplant, when compared with 1 week after transplant, among patients without and with PGD. ACR, acute cellular rejection; PGD, primary graft dysfunction.

a relatively rapid rise in peritransplant alphatorquevirus is associated with protection from PGD is analogous to existing literature that has found a comparably rapid rise in BAL.¹¹ The relatively low rate of PGD in patients with obstructive disease has likewise been noted previously.¹⁷ Nevertheless, this study is the first to identify a viral marker in the blood that is associated with the development of PGD in lung transplant recipients. These results need to be interpreted with caution due to small sample size, and because there is not a significant association between PGD and blood anellovirus when a multivariable analysis is applied.

We did not replicate the finding in our pediatric cohort, in whom a blood alphatorquevirus level drawn at 2 weeks following transplant could be used to predict subsequent episodes of ACR over the first 3 months following transplant.¹² While age-specific alterations in the virome are well-described in the human gut,¹⁸ there are little data about age-dependency of the virome in the human lung. Nevertheless, anellovirus prevalence has been established to change with the age of the human host,⁸ suggesting that there may be age-dependent changes in the human immune milieu that impact a host's anellovirus population. It is also possible that we did not see this alphatorquevirus association with ACR in this adult cohort due to the small sample size.

While there were not significant associations between early anellovirus levels and the subsequent development of ACR in this adult cohort, we evaluated trends in anellovirus temporal dynamics. The now established lag in anellovirus expansion after transplant,¹⁹ rather than reflecting an uninterpretable flux until a steady state population is established, may itself hold clues as to the immune receptivity of a transplant recipient. An immediate rapid rise in anellovirus after transplant is associated with protection from PGD, while there is a nonsignificant tendency toward subsequent slowing in anellovirus expansion in ACR-free patients. The significant association between rapid anellovirus rise and the protection from PGD did not hold up in a multivariable analysis; however, so further study of anellovirus dynamics in the immediate post-transplant period may be needed.

What is not clear from the present study is whether anelloviruses have a direct effect on peritransplant host immunity, or whether they are reflective of a protectively immunosuppressed

dysbiosis. Alphatorqueviruses encode miRNAs that inhibit interferon response,²⁰ and some anelloviruses can inhibit IL-6 and IL-8 transcription by interfering with NF-kappaB signaling.²¹ Due to the lack of understanding regarding the anellovirus life cycle, it is not yet clear why anellovirus dynamics have associations with key clinical outcomes. Nevertheless, given the importance of toll-like receptor signaling in PGD,²² it is possible that that anellovirus levels reflect the state of host innate immunity.

There are potentially confounding factors at the time of transplant, such as blood transfusion, that add to the complexity of delineating the role of anellovirus. We were unable to assess effect size and confounding in the current study since multivariable modeling showed no association between anellovirus and PGD. Given that the association present by bivariate analysis was not present in a multivariable analysis, it is possible that a larger cohort will be needed to evaluate posttransplant anellovirus dynamics, or that there are, in fact, confounding variables limiting the applicability of these findings. The conclusions that can be drawn from this study are limited by sample size, and it is possible that a larger cohort would be needed to identify significant trends in anellovirus expansion following transplant. Nevertheless, anellovirus has now been associated with several key outcomes in lung transplant, including PGD,¹¹ ACR,¹² and chronic lung allograft dysfunction,²³ suggesting that it is a vital biomarker even if it does not impact human health. The development of an animal model for anellovirus infection²⁴ could help us to interpret host-pathogen dynamics.

ACKNOWLEDGMENTS

We thank the patients who have contributed to this study.

REFERENCES

1. Rana A, Gruessner A, Agopian VG, et al. Survival benefit of solid-organ transplant in the United States. *JAMA Surg.* 2015;150:252–259.
2. Christie JD, Sager JS, Kimmel SE, et al. Impact of primary graft failure on outcomes following lung transplantation. *Chest.* 2005;127:161–165.
3. Christie JD, Carby M, Bag R, et al; ISHLT Working Group on Primary Lung Graft Dysfunction. Report of the ISHLT Working Group on Primary Lung Graft Dysfunction part II: definition. A consensus statement of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant.* 2005;24:1454–1459.
4. Christie JD, Kotloff RM, Pochettino A, et al. Clinical risk factors for primary graft failure following lung transplantation. *Chest.* 2003;124:1232–1241.
5. Bharat A, Kuo E, Steward N, et al. Immunological link between primary graft dysfunction and chronic lung allograft rejection. *Ann Thorac Surg.* 2008;86:189–95; discussion 196.
6. Tiriveedhi V, Gautam B, Sarma NJ, et al. Pre-transplant antibodies to α 1 tubulin and collagen-V in lung transplantation: clinical correlations. *J Heart Lung Transplant.* 2013;32:807–814.
7. Xiao J, Peng Z, Liao Y, et al. Organ transplantation and gut microbiota: current reviews and future challenges. *Am J Transl Res.* 2018;10:3330–3344.
8. Ninomiya M, Takahashi M, Nishizawa T, et al. Development of PCR assays with nested primers specific for differential detection of three human anelloviruses and early acquisition of dual or triple infection during infancy. *J Clin Microbiol.* 2008;46:507–514.
9. Young JC, Chehoud C, Bittinger K, et al. Viral metagenomics reveal blooms of anelloviruses in the respiratory tract of lung transplant recipients. *Am J Transplant.* 2015;15:200–209.
10. De Vlaminck I, Khush KK, Strehl C, et al. Temporal response of the human virome to immunosuppression and antiviral therapy. *Cell.* 2013;155:1178–1187.
11. Abbas AA, Diamond JM, Chehoud C, et al. The perioperative lung transplant virome: torque teno viruses are elevated in donor lungs and show divergent dynamics in primary graft dysfunction. *Am J Transplant.* 2017;17:1313–1324.

12. Blatter JA, Sweet SC, Conrad C, et al. Anellovirus loads are associated with outcomes in pediatric lung transplantation. *Pediatr Transplant.* 2018;22.
13. Maggi F, Pifferi M, Fornai C, et al. TT virus in the nasal secretions of children with acute respiratory diseases: relations to viremia and disease severity. *J Virol.* 2003;77:2418–2425.
14. Garcia-Álvarez M, Berenguer J, Alvarez E, et al. Association of torque teno virus (TTV) and torque teno mini virus (TTMV) with liver disease among patients coinfecting with human immunodeficiency virus and hepatitis C virus. *Eur J Clin Microbiol Infect Dis.* 2013;32:289–297.
15. Wille KM, Harrington KF, deAndrade JA, et al. Disparities in lung transplantation before and after introduction of the lung allocation score. *J Heart Lung Transplant.* 2013;32:684–692.
16. Stewart S, Fishbein MC, Snell GI, et al. Revision of the 1996 working formulation for the standardization of nomenclature in the diagnosis of lung rejection. *J Heart Lung Transplant.* 2007;26:1229–1242.
17. Barr ML, Kawut SM, Whelan TP, et al; ISHLT Working Group on Primary Lung Graft Dysfunction. Report of the ISHLT Working Group on Primary Lung Graft Dysfunction part IV: recipient-related risk factors and markers. *J Heart Lung Transplant.* 2005;24:1468–1482.
18. Lim ES, Zhou Y, Zhao G, et al. Early life dynamics of the human gut virome and bacterial microbiome in infants. *Nat Med.* 2015;21:1228–1234.
19. Görzer I, Jaksch P, Kundi M, et al. Pre-transplant plasma torque teno virus load and increase dynamics after lung transplantation. *Plos One.* 2015;10:e0122975.
20. Kincaid RP, Burke JM, Cox JC, et al. A human torque teno virus encodes a microRNA that inhibits interferon signaling. *Plos Pathog.* 2013;9:e1003818.
21. Zheng H, Ye L, Fang X, et al. Torque teno virus (SANBAN isolate) ORF2 protein suppresses NF-kappaB pathways via interaction with ikappaB kinases. *J Virol.* 2007;81:11917–11924.
22. Diamond JM, Wigfield CH. Role of innate immunity in primary graft dysfunction after lung transplantation. *Curr Opin Organ Transplant.* 2013;18:518–523.
23. Görzer I, Jaksch P, Strassl R, et al. Association between plasma torque teno virus level and chronic lung allograft dysfunction after lung transplantation. *J Heart Lung Transplant.* 2017;36:366–368.
24. Nishiyama S, Dutia BM, Stewart JP, et al. Identification of novel anelloviruses with broad diversity in UK rodents. *J Gen Virol.* 2014;95(Pt 7):1544–1553.