

SUPPLEMENTARY TABLE S1. CLINICAL, IMMUNOLOGICAL, AND VIROLOGICAL BASELINE CHARACTERISTICS AND DEMOGRAPHICS OF CHRONICALLY HIV-INFECTED STUDY PARTICIPANTS

<i>Characteristics</i>	<i>Cognitively normal (NL)</i> (n=23)	<i>Cognitively impaired (HAND)</i> (n=27)
Median age (range), years	36 (25 to 48)	35 (22 to 54)
Gender, male:female	10:13	10:17
Neurological diagnosis at baseline, <i>n</i> (%)		
NL	23 (100)	NA
HAND	NA	27 (100)
Neurological diagnosis at 12 months, <i>n</i> (%)		
NL	23 (100)	8 (30)
HAND	NA	19 (70)
Median plasma HIV RNA at baseline (range), log ₁₀ copies/mL	4.7 (3.1 to 5.9)	4.8 (3.3 to 5.9)
Subjects not controlling HIV RNA at 12 months, <i>n</i> (%)	1 (4)	4 (15)
Median CD4 count at baseline (range), cell/mm ³	191 (19 to 343)	182 (14 to 304)
Median CD4 count at 12 months (range), cell/mm ³	393 (130 to 849)	356 (5 to 768)
NPZ _{global} score at baseline (range)	0.12 (-0.45 to 1.10)	-0.71 (-2.6 to 0.06)
NPZ _{global} score at 12 months (range)	0.35 (-0.33 to 1.14)	-0.86 (-2.3 to 0.01)

HAND, HIV-associated neurocognitive disorders; NA, not applicable; NL, cognitively normal.