

Supplementary Data

SUPPLEMENTARY TABLE S1. SONOGRAPHIC FEATURES ASSOCIATED WITH PREDICTING THYROID CANCER

Analysis of effects

<i>Effect</i>	<i>Wald chi-square</i>	<i>Pr > ChiSq</i>
Composition	69.4029	<0.0001
Echogenicity	66.0343	<0.0001
Calcification	107.9542	<0.0001
Margins	57.0829	<0.0001
Doppler	4.6631	0.1982
Location	12.8478	0.0050

SUPPLEMENTARY TABLE S2. MEAN MAXIMUM SIZE AS CALCULATED IN FOUR SEPARATE ANALYSIS OF VARIANCE ANALYSES

<i>Location</i>	<i>Maximum size</i>	<i>Maximum size and Dx</i>
Isthmus	20.24	18.47
Low	23.86	22.58
Middle	25.43	24.00
Upper	18.86	17.77

SUPPLEMENTARY TABLE S3. COMPARISONS OF MAXIMUM SIZE BETWEEN LOCATIONS IN FOUR SEPARATE ANALYSIS OF VARIANCE ANALYSES

<i>Location comparison</i>	<i>Maximum size</i>	<i>Maximum size and Dx</i>
Isthmus vs. middle	<0.0001	0.0002
Isthmus vs. upper	0.6306	0.9697
Isthmus vs. low	0.0019	0.0170
Middle vs. upper	<0.0001	0.0003
Middle vs. low	0.0098	0.3682
Upper vs. low	<0.0001	<0.0001