

2012

# Interethnic differences in $\beta$ -2 agonist treatment of acute asthma on electrolyte/glucose concentrations

Ian Ferguson

*Washington University School of Medicine in St. Louis*

Kazuko Matsuda

*Medicinova Inc.*

Stacey House

*Washington University School of Medicine in St. Louis*

John Schneider

*Washington University School of Medicine in St. Louis*

Kristen Aubuchon

*Washington University School of Medicine in St. Louis*

*See next page for additional authors*

Follow this and additional works at: [http://digitalcommons.wustl.edu/em\\_conf](http://digitalcommons.wustl.edu/em_conf)

## Recommended Citation

Ferguson, Ian; Matsuda, Kazuko; House, Stacey; Schneider, John; Aubuchon, Kristen; Lewis, Lawrence, "Interethnic differences in  $\beta$ -2 agonist treatment of acute asthma on electrolyte/glucose concentrations" (2012). Conference Abstracts and Posters. Paper 13. [http://digitalcommons.wustl.edu/em\\_conf/13](http://digitalcommons.wustl.edu/em_conf/13)

This Presentation Paper is brought to you for free and open access by the Division of Emergency Medicine/Emergency Care Research Section at Digital Commons@Becker. It has been accepted for inclusion in Conference Abstracts and Posters by an authorized administrator of Digital Commons@Becker. For more information, please contact [engeszer@wustl.edu](mailto:engeszer@wustl.edu).

---

**Authors**

Ian Ferguson, Kazuko Matsuda, Stacey House, John Schneider, Kristen Aubuchon, and Lawrence Lewis

# Interethnic Differences in $\beta$ -2 Agonist Treatment of Acute Asthma on Electrolyte/Glucose Concentrations

Ian Ferguson<sup>1</sup>

Kazuko Matsuda<sup>2</sup> MD PhD, Stacey House<sup>1</sup> MD PhD, John Schneider<sup>1</sup>,  
Kristen Aubuchon<sup>1</sup>, and Lawrence Lewis<sup>1</sup> MD

<sup>1</sup>Washington University in St. Louis, St. Louis, MO

<sup>2</sup>Medicinova Inc., San Diego, CA



# Objective

- ◆ Is there an ethnic disparity regarding the effect of plasma albuterol on:
  - ◆ Potassium
  - ◆ Magnesium
  - ◆ Bicarbonate
  - ◆ Phosphate
  - ◆ Glucose

# $\beta$ 2-Agonists and Electrolyte Abnormalities

- ◆ Potassium
  - ◆ Hypokalemia through increased cellular uptake
- ◆ Glucose
  - ◆ Gluconeogenesis
- ◆ Phosphate
  - ◆ Hypophosphatemia may be associated with ventilatory failure
- ◆ Bicarbonate
  - ◆ Increase in lactic acidosis (Type B) in asthma

# Ethnic Differences in Asthma

- ◆ Higher morbidity and mortality in African Americans compared to Caucasians with asthma
- ◆ Higher rates of ED visits (350%) and hospitalizations (240%)
- ◆ Reports of interethnic differences in  $\beta_2$  treatment for acute asthma
  - ◆ SABA overuse and tolerance mechanism
  - ◆ Genetic differences at  $\beta_2$ -receptor

# Study Methods

## **MN-221 Trial**

- ◆ Sub-group analysis of large, multi-center RCT evaluating a novel IV  $\beta$ 2-agonist for acute asthma exacerbation

## **Select Exclusion Criteria**

- ◆ <18 and >55 years old
- ◆ At risk for increased  $\beta$ 1 response
- ◆ Significant smoking history
- ◆ COPD, or other chronic lung disease or respiratory dysfunction

# Methods Continued

## Subject Inclusion

- ◆ Serum Mg, K, bicarbonate, and glucose were measured at “baseline” and “1.25hrs post-baseline”
- ◆ 70 placebo subjects with plasma albuterol (R- and S-) data

## Analysis

- ◆ Percent of lab values outside normal range
- ◆ Pearson correlations for albuterol to electrolyte/glucose concentration

# Results

- ◆ 43 African-Americans and 27 Caucasians
- ◆ Significant correlations observed in 4/5 measures, all ethnic specific
  - ◆ Largest effect size with potassium and bicarbonate
  - ◆ Smaller effect size with magnesium and glucose
  - ◆ No effect with Phosphate, however large % low values

# Correlation Results

Pearson r [95% CI] p value, R <sup>2</sup>	Total albuterol: Caucasians	Total albuterol: African-Americans
Magnesium (mEq/L)	-0.15 [-0.43, -0.15] p= 0.32	0.36 [0.13, 0.56] p<0.01*, 0.13
Potassium (mmol/L)	-0.55 [-0.7, -0.31] p< 0.01*, 0.31	-0.08 [-0.31, -0.14] p=0.44
Glucose (mg/dL)	0.34 [0.07, 0.56] p=0.015*, 0.11	0.10 [-0.11, 0.32] p=0.33
Phosphate (mg/dL)	-0.07 [-0.35, 0.21] p= 0.62	0.15 [-0.07, 0.36] p= 0.17
Bicarbonate (mmol/L)	-0.55 [-0.72, -0.32] p<0.01*, 0.30	-0.15 [-0.35, 0.07] p= 0.18

# Metabolic Abnormalities

<b>Value [Lab Cut-Off]</b>	<b>Median [IQR]</b>	<b>N (%) AA ONR</b>	<b>N (%) CA ONR</b>
Potassium [<3.6] mmol/L	3.8 [3.5-4.1]	14 (32.5%)	11 (46%)
Magnesium [>2.1] mEq/L	1.7 [1.5-1.8]	3 (8.8%)	0 (0%)
Phosphate [<2.4] mg/dL	2.7 [2.3-3.2]	17 (40.5%)	10 (40%)
Bicarbonate [<20] mEq/L	22 [20-24]	10 (23%)	9 (36%)
Glucose [>140] mg/dL	120 [107-152]	22 (51%)	11 (42%)

# Conclusions

- ◆ Ethnic-specific correlations between plasma albuterol and serum K, Bicarb, Mg, and Glu.
- ◆ Specific to Caucasian population:
  - ◆ Sig negative correlation with K and Bicarb
  - ◆ Sig positive correlation with Glucose
- ◆ Specific to African American population:
  - ◆ Sig positive correlation with Mg
- ◆ High % hypophosphatemia observed for both populations

# Potential Clinical Implications

- ◆ When repeated  $\beta$ 2-agonist treatments given for acute asthma, closer ED monitoring may be required for:
  - ◆ Low K, low Bicarb and high Glucose in Caucasians
  - ◆ High magnesium in African Americans
- ◆ Future research implications
  - ◆ These data suggest that combining ethnic groups may mask true effect size in one group

# Limitations

- ◆ Clinical relevance depends on magnitude of change—generally small effect size observed
- ◆ Relatively small sample size due to sub-group analysis

Questions?

## Exclusion Criteria

- Administration of parenteral (intravenous or subcutaneous) beta agonist within 6 hours prior to randomization.
- A current or prior diagnosis or suspected diagnosis of COPD or other chronic lung disease other than asthma, the presence of pneumonia, or presence of significant other respiratory dysfunction such as pneumothorax, pneumomediastinum, or pulmonary edema.
- Known or suspected vocal cord dysfunction syndrome or the presence of aspirated foreign body (known or suspected).
- History of any current clinical evidence suggesting cardiomyopathy or congestive heart failure, history or presence of tachyarrhythmia, with the exception of sinus tachycardia, or a heart rate  $\geq 140$  bpm.
- Hypokalemia, defined as potassium level  $\leq 2.8$  mEq/L obtained at Screening.
- Significant cardiac, renal, hepatic, endocrine, metabolic, neurologic or other systemic disease.
- Self-reported history of greater than 15 packer year smoking history.
- Fever  $\geq 102.0$  °F (38.9°C).
- Uncontrolled hypertension defined as blood pressure  $\geq 170/100$  mm Hg.
- Need for immediate intubation, mechanical ventilation, or non-invasive positive pressure ventilation as determined by the Investigator.
- Pregnant or lactating females.
- Participating in another clinical study with an investigational drug within 30 days of randomization.
- Positive urine drug screen for cocaine, methamphetamine or PCP.
- Any subject with a known allergy components of the MN-221 drug product or other beta agonists.
- Previous exposure to MN-221
- Use of theophylline, beta blockers, diuretics, digoxin, MAO inhibitors, or tricyclic antidepressants, within 2 weeks prior to randomization.

# Plasma/ Nebulized Albuterol Correlation

Plasma Albuterol	Mean	Nebulized Dose (mg)
Total (R+S)	7.0 ng/mL	12.1 mg